

Genetic wealth, population health: Major histocompatibility complex variation in captive and wild ring-tailed lemurs (*Lemur catta*)

Kathleen E. Grogan^{1,2}  | Michelle L. Sauter³ | Frank P. Cuzzo⁴ | Christine M. Drea^{1,2,5}

¹University Program in Ecology, Duke University, Durham, NC, USA

²Department of Evolutionary Anthropology, Duke University, Durham, NC, USA

³Department of Anthropology, University of Colorado-Boulder, Boulder, CO, USA

⁴Lajuma Research Centre, Louis Trichardt (Makhado) 0920 South Africa

⁵Department of Biology, Duke University, Durham, NC, USA

Correspondence

Kathleen E. Grogan, O. Wayne Rollins Research Center, Emory University, Atlanta, GA, USA.
Email: kathleen.e.grogan@gmail.com

Funding information

Duke University; International Primatological Society; Primate Conservation; University of Colorado Boulder; University of North Dakota; National Science Foundation, Grant/Award Number: BCS 0922465, BCS-1232570 and IOS-071900; Margot Marsh Biodiversity Foundation; St. Louis Zoo; American Society of Primatologists

Abstract

Across species, diversity at the major histocompatibility complex (MHC) is critical to individual disease resistance and, hence, to population health; however, MHC diversity can be reduced in small, fragmented, or isolated populations. Given the need for comparative studies of functional genetic diversity, we investigated whether MHC diversity differs between populations which are open, that is experiencing gene flow, versus populations which are closed, that is isolated from other populations. Using the endangered ring-tailed lemur (*Lemur catta*) as a model, we compared two populations under long-term study: a relatively “open,” wild population ($n = 180$) derived from Beza Mahafaly Special Reserve, Madagascar (2003–2013) and a “closed,” captive population ($n = 121$) derived from the Duke Lemur Center (DLC, 1980–2013) and from the Indianapolis and Cincinnati Zoos (2012). For all animals, we assessed MHC-DRB diversity and, across populations, we compared the number of unique MHC-DRB alleles and their distributions. Wild individuals possessed more MHC-DRB alleles than did captive individuals, and overall, the wild population had more unique MHC-DRB alleles that were more evenly distributed than did the captive population. Despite management efforts to maintain or increase genetic diversity in the DLC population, MHC diversity remained static from 1980 to 2010. Since 2010, however, captive-breeding efforts resulted in the MHC diversity of offspring increasing to a level commensurate with that found in wild individuals. Therefore, loss of genetic diversity in lemurs, owing to small founder populations or reduced gene flow, can be mitigated by managed breeding efforts. Quantifying MHC diversity within individuals and between populations is the necessary first step to identifying potential improvements to captive management and conservation plans.

KEYWORDS

conservation genetics, genetic diversity, immunogenetics, inbreeding, MHC-DRB, strepsirrhine primate

1 | INTRODUCTION

In taxa as diverse as fish, amphibians, reptiles, birds, and mammals, diversity and/or the specific alleles present at the major histocompatibility complex (MHC) have become important measures of genetic health. Polymorphism at MHC loci has been linked to many proxy measures of quality (e.g., disease resistance, ornamentation, life span) or fitness (e.g., mating success, offspring production) in individuals, populations, and species (Bernatchez & Landry, 2003; Sommer, 2005). For conservation biologists, these measures reflect the potential for adaptation to emerging zoonosis, habitat fragmentation, and climate change (Ujvari & Belov, 2011). Therefore, understanding whether MHC diversity differs between wild and captive populations could be significant to the development of captive management strategies, as well as to coordinating the protection and recovery of wild populations. Here, we compare diversity at the MHC-DRB gene between a “closed,” captive population and a more “open,” wild population. This comparison allows investigation of the influences of genetic isolation and balancing selection on MHC diversity, as well as the interplay between these forces and captive management.

Genetic diversity, defined as the number of genetic variants within a population or between individuals, allows future generations to adapt to environmental change. Loss of genetic diversity can be especially detrimental to populations that are small, fragmented, or isolated (reviewed in Frankham, 2005a, 2005b; Hedrick & Kalinowski, 2011). Such populations are generally less genetically diverse than are larger, more contiguous populations (Madsen et al., 2000; Sutton, Robertson, & Jamieson, 2015; reviewed in Ouborg, Angeloni, & Vergeer, 2010), owing to population bottlenecks, founder effects, genetic drift, or inbreeding—mechanisms that may act simultaneously and synergistically. Similarly, captive populations are often less genetically diverse than their wild counterparts. They are often founded by only a few individuals and, once established, can be considered “closed” because they experience little additional influx, either from the wild or from other captive groups (reviewed in Ivy, 2016; Fernández-de-Mera et al., 2009; Yao, Zhu, Wan, & Fang, 2015; but see Newhouse & Balakrishnan, 2015). Compared to captive populations, all but the most geographically isolated of wild populations are typically more “open” owing to greater gene flow. Thus, relative to wild populations, captive populations may experience more inbreeding and genetic drift and consequent loss of genetic diversity. For these reasons, captive populations can serve as important proxies for fragmented, wild populations, specifically for examining the consequences of founder effects and decades of isolation on genetic diversity. Assessments of the genetic diversity of captive populations are also critical if these populations are to serve as potential sources for reintroduction efforts for endangered species.

At the individual level, decreases in genetic diversity across the genome and at specific loci typically reduce quality or fitness (i.e., individuals that have decreased genetic diversity have poorer health, shorter life spans, and produce fewer offspring than do individuals that have greater genetic diversity: Crnokrak & Roff, 1999; Frankham, 2005a; Keller & Waller, 2002). At the population level, loss of genetic diversity

reduces adaptive potential, thereby increasing the risk of extinction. The loss of genetic diversity from population decline, genetic drift, or inbreeding is thus one of the major threats to endangered species (Frankham, 2005a, 2005b). Nonhuman primates and other long-lived species are particularly vulnerable to the negative effects of genetic loss because their slow life histories and low rate of reproduction can severely hinder the recovery of genetic diversity (Charpentier, Widdig, & Alberts, 2007). From a theoretical perspective, quantifying the existing genetic diversity within populations is essential for (1) identifying the influence of genetic diversity on quality, fitness, and adaptive potential, (2) elucidating the evolutionary pressures influencing current levels of biological variation, and (3) understanding the specific mechanisms of action. From an applied perspective, understanding the relationship between genetic diversity and fitness has become a task of increasing urgency for recovery and conservation of endangered species.

Because almost half of all nonhuman primate species are classified as “threatened” (IUCN 2010), the threat from loss of genetic diversity is of particular importance for developing successful primate conservation plans. Although transgenerational health and reproductive data are logistically difficult to obtain for long-lived species, in a few semi-free-ranging primates, loss of genetic diversity has been correlated with increased health risks, including greater ectoparasite prevalence and burden, as well as decreased immunocompetence (Charpentier, Williams, & Drea, 2008; Noble, Chesser, & Ryder, 1990; Van Coillie et al., 2008). These health threats, however, did not translate into shorter life spans for less heterozygous individuals (Charpentier, Williams, & Drea, 2015). Decreasing genetic diversity is often measured using neutral heterozygosity, nH_0 , calculated from microsatellite data (Hartl & Clark, 1997; Reed & Frankham, 2001). Nonetheless, these correlations denote an indirect relationship between genetic diversity and health risks because differences in microsatellites do not change protein sequences. Therefore, nH_0 does not equate to functional genetic diversity that reflects the adaptive potential of a population. Relative to nH_0 , genetic diversity at protein coding or regulatory regions is better estimates of evolutionary potential because they are directly associated with factors that influence individual fitness, population viability, and the ability to respond to environmental change (Hedrick, 2001; Ouborg et al., 2010; Väli, Einarsson, Waits, & Ellegren, 2008).

We could gain a better understanding of how an individual's genetic makeup affects its quality or fitness by examining genetic variation at critical functional genes, such as those of the MHC. The MHC is the most polymorphic gene family within vertebrates and is responsible for the activation of the adaptive immune system. MHC genes encode proteins that distinguish between “self” and “non-self” peptides (Piertney & Oliver, 2006). MHC products bind and present “non-self” peptides to immune system cells, initiating the body's immune response (Piertney & Oliver, 2006). Because each MHC molecule “recognizes” a particular subset of pathogens, both overall MHC diversity and specific MHC alleles have been linked to resistance or susceptibility to pathogens (Biedrzycka, Kloch, Buczek, & Radwan, 2011; Carrington & Bontrop, 2002; Evans & Neff, 2009; Oliver, Telfer,



FIGURE 1 Photo of ring-tailed lemur by CM Drea

& Piertney, 2009; Savage & Zamudio, 2011; Wegner, Kalbe, Kurtz, Reusch, & Milinski, 2003; reviewed in Bernatchez & Landry, 2003 and Sommer, 2005). In some species, the links between MHC and disease are independent of the impact of nH_e , providing evidence that parasite load is specifically MHC-mediated, rather than a result of nH_e (Schwensow, Fietz, Dausmann, & Sommer, 2007; Westerdahl et al., 2005). Beyond the direct link between MHC and health, MHC diversity is also linked to differential lifetime survival (Huchard, Knapp, Wang, Raymond, & Cowlshaw, 2010) and reproductive success (Kalbe et al., 2009; Knapp, Ha, & Sackett, 1996; Ober, 1999; Sauermann et al., 2001; Setchell, Charpentier, Abbott, Wickings, & Knapp, 2010; Smith et al., 2010; Thoß, Ilmonen, Musolf, & Penn, 2011).

Given the correlations between MHC diversity and proxies of quality or fitness, such as health, individual and offspring survival, and reproductive success, measuring MHC diversity should be a fundamental component of wildlife conservation and captive-breeding efforts (reviewed in Ujvari & Belov, 2011). Individuals in small or fragmented populations and populations that have undergone a bottleneck often show reduced MHC-DRB diversity when compared to individuals in large or connected populations and in populations prior to size reduction (Agudo et al., 2011; Bollmer, Ruder, Johnson, Eimes, & Dunn, 2011; Sutton, Nakagawa, Robertson, & Jamieson, 2011; Sutton et al., 2015). A decline in genetic diversity can be further compounded by the more pronounced effects of genetic drift on smaller populations. Thus, small or fragmented populations are subject to a double-pronged threat.

As an endangered primate (Andriaholinirina et al., 2014), the ring-tailed lemur (*Lemur catta*; Figure 1) is currently facing many anthropogenic stressors, including habitat fragmentation, increased contact with humans and domestic animals, and climate change (Schwitzer et al., 2013). Evaluated by the IUCN in 2014, the population density of

ring-tailed lemurs in Madagascar was found to be low and restricted to isolated fragments. Habitat loss, hunting pressure, and the illegal pet trade are estimated to have produced at least a 50% reduction in the wild population over the last few decades (Andriaholinirina et al., 2014). Population estimates from areas known to historically contain ring-tailed lemurs indicate a dramatic population reduction to as few as 2,500 wild animals spread across southwestern Madagascar (Gould & Sauther, 2016; LaFleur et al., 2016), increasing the probability of loss of genetic diversity due to genetic drift. Because of these threats, ring-tailed lemurs might be representative of vulnerable species worldwide. Moreover, the extensive, long-term study of large populations in the wild and in captivity (Gould, Sussman, & Sauther, 2003; Jolly, 1966; Sauther, Gould, Cuozzo, & O'Mara, 2015; Sussman et al., 2012; Zehr et al., 2014), coupled with confirmed susceptibility to inbreeding depression (Charpentier et al., 2008), make the ring-tailed lemur a suitable model for testing if functional genetic diversity at the MHC differs between captive and wild populations or if MHC diversity declines in captivity.

We used two measures of MHC diversity to compare wild and captive individuals: absolute MHC allelic diversity and the diversity of MHC supertypes, which are groups of MHC alleles with similar motifs in their binding pockets despite nucleotide differences (Doytchinova & Flower, 2005; Huchard, Weill, Cowlshaw, Raymond, & Knapp, 2008; Sepil, Lachish, Hinks, & Sheldon, 2013). These alleles therefore likely bind similar or identical pathogen peptides and can be grouped according to functional binding properties. Using these data, we evaluated the effect of breeding management on the more “closed,” captive population by estimating if its diversity (1) differs from that of the more “open,” wild population and (2) has changed across decades. We also used previously published microsatellite data to assess the relationship between nH_e and functional genetic diversity (or MHC-DRB diversity: Charpentier et al., 2008; Grogan, McGinnis, Sauther, Cuozzo, & Drea, 2016; Pastorini et al., 2015), as well as to estimate effective population sizes.

2 | METHODS

2.1 | Subjects

Our subjects were 301 ring-tailed lemurs, including 121 (67 females; 51 males; three infants of unknown sex) captive animals from three institutions that are part of the Association of Zoos and Aquariums (AZA) in the USA and 180 (86 females; 94 males) wild animals from Bezà Mahafaly Special Reserve (BMSR), in the region of Atsimo-Andrefana, Madagascar.

The wild ring-tailed lemurs at BMSR have been studied since 1987 and have been collared for identification and monitored monthly since 2003 (for more details on BMSR and the monthly census of the wild population, see Sussman et al., 2012). From 1987 to 2013, this population has generally averaged approximately 95 adult, ring-tailed lemurs annually (range = 58–120; Gould, Sussman, & Sauther, 1999; Gould et al., 2003; Sauther & Cuozzo unpublished data). Our BMSR population represents the majority (>90%) of adults collared from 2003 to 2013. We consider this population to be “open,” owing to the

known immigration and emigration of animals (Sauther, Sussman, & Gould, 1999; Sussman, 1991, 1992).

The captive ring-tailed lemurs comprised 52 individuals residing, from 2012 to 2013, at the Duke Lemur Center (DLC; $n = 32$) in Durham, NC, the Indianapolis Zoo ($n = 18$), in Indianapolis, IN, and the Cincinnati Zoo ($n = 2$), in Cincinnati, OH, as well as an additional 69 individuals that had previously resided at the DLC, from 1980 to 2012, for which DNA samples could be obtained. These captive animals included individuals born at any of the three institutions, as well as individuals that had transferred between these and other institutions, either to address a housing constraint or to satisfy a breeding recommendation from the AZA Species Survival Plan (SSP) that originated in 1995. All individuals were housed in mixed-sex groups, with similar housing and provisioning conditions across the three institutions (for details, see Scordato, Dubay, & Drea, 2007; McKenney, Rodrigo, & Yoder, 2015). In past decades, groups at the DLC ranged from male–female pairs to multimale, multifemale groups as large as 25 individuals. More recently, groups have included a maximum of 12 individuals. We treated the captive animals as a single population, regardless of home institution, because they are managed as such by the AZA; however, we have surveyed only three of over 200 institutions that house ring-tailed lemurs. We also consider them a relatively “closed” population because of the small group of founders and because, each year, only a few individuals breed or transfer between institutions.

2.2 | Sampling

We obtained whole blood or tissue collected from wild and captive lemurs by veterinarians following established protocols (Charpentier et al., 2008; Sauther & Cuzzo, 2008). All of our sampling methods followed approved animal handling guidelines and protocols of the Institutional Animal Care and Use Committees of Duke University, the University of North Dakota, and/or the University of Colorado (most recently, these included Duke University IACUC #A143-12-05, approved 05/25/2012, and University of North Dakota IACUC #0802-2, approved 04/03/08). Sample collection in Madagascar was also approved by Madagascar National Parks and CITES (05US040035/9).

2.3 | DNA extraction

We extracted DNA from the samples following the manufacturers' instructions, using either DNA miniprep kits (Sigma, St. Louis, MO) or DNeasy® Blood and Tissue kits (Qiagen, Valencia, CA). We preserved whole blood from the wild subjects on Whatman FTA® Classic cards (GE Healthcare Life Sciences, Buckinghamshire, UK). From these cards, we extracted DNA and increased its quality and quantity by whole genome amplification (Repli-G Single Cell Kit®, Qiagen, Valencia, CA).

2.4 | Genotyping

To genotype individuals at the MHC-DRB loci, we used parallel-tagged sequencing to pool amplicons of 50–125 individuals, including

two to five replicates per individual, for a total of 845 amplicons in eight PCR pools. We sequenced a 171-bp fragment of the MHC-DRB second exon, containing the antigen-binding site, using three Ion Torrent PGM® 314v2 chips (Life Technologies, Grand Island, NY) and five 454 Titanium® 1/8th lanes (Roche, Nutley, NJ). To distinguish alleles from artifacts, we used a published workflow that relies upon relative frequency of variants within an amplicon and comparisons between replicate PCRs for each sample to identify alleles (for details, see Grogan et al., 2016). Once animals were genotyped, we determined MHC supertype by identifying nucleotide sites under positive selection or positively selected sites (PSS), via the CODEML analysis in PAML Version 4.7 (Yang, 2007). We used the physicochemical properties of PSS amino acids to determine MHC-DRB superotypes of alleles or groups of alleles with similar binding properties at antigen-binding sites (ABS). We used heat mapping to determine superotypes in Genesis Version 1.7.6 (See Table S1 for details regarding MHC superotypes and corresponding MHC alleles: Doytchinova & Flower, 2005; Huchard et al., 2008; Schwensow et al., 2007; Sepil et al., 2013).

2.5 | Statistical analyses

Using an ANOVA, we compared the average MHC-DRB diversity of captive vs. wild individuals. We investigated whether sampling effort affected the number of alleles detected in each population compared to the actual allelic richness present in each population. To do so, we conducted permutation tests by randomly sampling ten individuals per population and counting the number of unique alleles represented in this random sample. After 100 iterations, we calculated the mean and *SD* of a sampling effort of ten. We then repeated this procedure in an incremental process, adding ten to our sampling effort in each step, up to a sample of 100 individuals. We conducted these permutation tests for both the wild population ($n = 180$) and the captive DLC population ($n = 101$), and plotted the average number of MHC-DRB alleles detected per given sampling effort. We performed all analyses in RStudio (Version 3.0.2).

To evaluate the use of nH_o as a proxy for MHC diversity, we used a linear regression to compare MHC-DRB diversity to microsatellite data separately for each population, using data from previously published work (for details of microsatellite genotyping of the captive population, see Charpentier et al., 2008; for details of microsatellite genotyping of the wild population, see Pastorini et al., 2015). Briefly, 73 ring-tailed lemurs from the DLC population were each genotyped at 10–15 microsatellite loci. From BMSR, 130 individuals captured from 2003 to 2005 were genotyped at 10 microsatellite loci. Only microsatellite loci that conformed to Hardy–Weinberg expectations were used to calculate observed nH_o per individual. Within the 73 captive individuals, nH_o ranged from 0.21 to 0.86 (mean \pm SE = 0.56 ± 0.02 ; Charpentier et al., 2008). In contrast, nH_o for the wild population ranged from 0.33 to 1.0 (mean \pm SE = 0.76 ± 0.14 ; Pastorini et al., 2015).

Lastly, to assess the influence of approximately 50 years of captivity and captive management on MHC-DRB diversity in the DLC population, we divided our study period into decades and compared the average MHC-DRB diversity of the infants born during each




FIGURE 2 Genetic diversity in ring-tailed lemurs showing the number of (a) MHC-DRB alleles and (b) MHC-DRB superotypes at the individual level for both captive (green) and wild (black) populations

decade from 1980 to 2013 (see Charpentier et al., 2008). To disentangle the relative influences of the founder effect from genetic drift over generations, we used the microsatellite data and the linkage disequilibrium method (Laurie-Ahlberg & Weir, 1979; Hill, 1981; Waples & Do, 2010) in NeEstimator V2 (Do et al., 2014) to estimate the effective population size for the DLC population ($n = 80$) per decade

TABLE 1 Individual and population level of MHC-DRB diversity across captive animals at three institutions compared to animals from one wild population

	Captive: Cincinnati zoo	Captive: Indianapolis zoo	Captive: DLC	Total captive population	Wild population
Number of individuals genotyped	2	18	101	121	180
Average number of alleles per individual	3.5	2.56	2.0	2.10	2.78
Total number of alleles in population	4	12	16	20	53
Number of private alleles	0	2	1	2	35
Average number of superotypes per individual	3.5	2.11	1.51	1.65	2.6
Total number of superotypes in population	4	9	11	13	24
Number of private superotypes	0	1	1	1	12

in captivity from 1980 to 2007, as well as for the wild population ($n = 130$) from 2003 to 2005. These data are somewhat limited, however, as they do not include every individual residing at the DLC or BMSR during these years, nor every individual that was genotyped for MHC-DRB. Additionally, we lack microsatellite data from individuals born at or transferred to the DLC after 2007, as well as for all individuals residing at the Cincinnati and Indianapolis Zoos and born at BMSR after 2006.

3 | RESULTS

In testing for differences in MHC-DRB diversity between wild and captive populations of ring-tailed lemurs, we found that wild individuals were significantly more diverse than were captive individuals (Figure 2). Specifically, there were significantly more MHC-DRB alleles ($n = 301$, $F = 23.97$, $p < .001$; Figure 2a; Table 1), as well as significantly more MHC-DRB superotypes ($n = 301$, $F = 7.035$, $p < .001$; Figure 2b; Table 1) in wild individuals compared to captive individuals. Wild individuals possessed a mean ($\pm SD$) of 2.78 (± 1.34) alleles (range = 1–7) and 2.60 (± 1.32) superotypes (range = 1–7), whereas captive individuals possessed 2.1 (± 0.86) alleles (range = 1–5) and 1.65 (± 0.83) superotypes (range = 1–5).

The wild population also showed more allelic richness, with 52 unique MHC-DRB alleles and 24 MHC superotypes, compared to 13 superotypes comprised of 20 MHC-DRB alleles in the captive population (Figure 3). The two populations shared 18 alleles (12 superotypes); however, the captive population had two private alleles not found in the wild population and one private MHC supertype. In contrast, the wild population had 35 private alleles and 12 MHC superotypes that were not shared by the captive population (for details of which alleles were shared or were private, see Table S1). Neither sex nor age influenced the distribution of MHC-DRB alleles or superotypes between populations or individuals.

Within each population, MHC-DRB alleles and superotypes were unevenly distributed (Figure 3). Within the captive population, three alleles were present in a majority of the individuals, whereas the remaining 17 alleles were present in only a few individuals (overall range = 0.0–60.3%, overall mean = 10.6%; Figure 3a). The most abundant allele was present in more than 60% of captive individuals, and





FIGURE 3 The frequency distribution of (a) MHC-DRB alleles (Leca-DRB) and (b) MHC-DRB supertypes in captive (green) and wild (black) populations of ring-tailed lemurs




FIGURE 4 The mean number (\pm SE) of unique MHC-DRB alleles detected in captive (green circles) and wild (black circles) populations of ring-tailed lemurs. The unique alleles detected were dependent upon sampling effort, estimated using resampling techniques, and are compared to the number of unique alleles present in the captive (dashed green line) and wild (dashed black line) populations




FIGURE 5 Effective population size (N_e) at the DLC and BMSR, calculated using the linkage disequilibrium method. Genotyped infants from the DLC (in green circles) were grouped according to individuals born within each decade from 1980 through 2010, whereas N_e from BMSR (represented by a black circle) was estimated from animals residing in the reserve between 2005 and 2006. Upper and lower limits indicate 95% confidence intervals calculated using the jackknife method

more strikingly, the most common MHC supertype was present in >90% of captive individuals (range = 1.0%–90.0%, mean = 6.0%). By contrast, the most abundant MHC-DRB allele within the wild population was present in only 38.8% of individuals (range = 0.0–38.8%, mean = 5.2%; Figure 3b), and the most abundant supertype was present in only 51% of individuals (range = 2.0%–51.0%, mean = 9%). Thus,

we found that MHC-DRB variation in the wild population was more evenly spread than was MHC-DRB variation in the captive population.

Next, we calculated the average allelic richness detected for a given sampling effort (Figure 4). In the captive population, the number of unique alleles detected reached a plateau near a sampling effort of 60 individuals or 50% of the captive population sampled. When 60 captive animals were sampled, approximately 75% of 20 possible unique alleles were detected. In the wild population, however, sampling 55% of the population detected only 44% ($n = 23$) of the 52 unique MHC-DRB alleles present. Therefore, the more diverse a population, the greater the sampling effort required to detect the majority of the MHC diversity present.

In our comparison between neutral and functional diversity, we found nH_o to be significantly and positively correlated with the number of MHC-DRB alleles ($n = 71$, $F = 5.884$, $p = .018$) in the captive population; however, variation in nH_o explained little of the variation in the number of MHC-DRB alleles (adjusted $R^2 = 0.065$, slope = 1.26). Conversely, in the more diverse wild population, MHC diversity was not correlated with nH_o ($n = 88$, $F = 0.002$, $p = .962$). Consequently, although average nH_o decreased significantly across the decades from 1980 to 2010 (see Charpentier et al., 2008), MHC-DRB diversity in infants born at the DLC remained constant (Spearman correlation, $n = 83$, $r = -0.11$, $p = .28$). Despite careful management, the estimated effective population sizes decreased from 15.4 individuals during 1980–1989 to 14.4 individuals during 1990–1999, then down to 12.9 individuals during 2000–2009. In contrast, the effective population size from 2003 to 2005 at BMSR was 52.1 individuals (Figure 5). After 2010, however, MHC-DRB diversity in captive-born infants increased significantly (Spearman correlation, $n = 93$, $r = 0.35$, $p < .001$; Figure 6) compared to the MHC-DRB diversity of infants born in the decades prior to 2010. In fact, the MHC-DRB diversity of infants born after 2010 (mean \pm SD = 2.8 ± 0.92) was comparable to the average MHC-DRB diversity of individuals in the wild ($F = 0.002$, $p = .966$).

4 | DISCUSSION

In a comparison of two populations of ring-tailed lemurs, both individual- and population-level MHC-DRB diversity were greater in wild animals than in captive animals. Individuals from among the wild lemurs possessed, on average, one allele more than did individuals from among the captive lemurs. The number of unique MHC-DRB alleles and MHC-DRB supertypes present in the wild population was more than twice the number in the captive population at the three institutions included in this study. In both the wild and captive populations, allelic frequency showed a skewed distribution, as over 50% of MHC-DRB alleles present in each population were found in five or fewer individuals. Just as some microsatellite variation already has been lost




FIGURE 6 Mean MHC-DRB allelic richness (\pm SD) possessed by ring-tailed lemur infants born at the DLC, separated typically by decade of birth from 1980 through the birth season of 2013. The number of infants born per period is shown above each data point

in captivity (Charpentier et al., 2008), these rare MHC-DRB alleles are at greater risk of being lost via genetic drift than they would be in a larger population. In the captive population, the decreased genetic diversity likely reflects effects of 1) a genetic bottleneck, owing to only a few founding individuals having been initially captured from the wild, 2) decades of reduced gene flow between institutions, and 3) genetic drift, especially in small populations. Without the subsequent introduction of new MHC alleles, captive management efforts at the DLC, based on pedigree data, did not counteract the loss of genetic diversity over four decades (Charpentier et al., 2008); however, we found evidence of a genetic rescue occurring via the transfer of new individuals to the DLC after 2010.

Genetic drift may, in fact, have a stronger effect on immune system or functional genes than on neutral genetic diversity (Bateson, Whittingham, Johnson, & Dunn, 2015; Froeschke & Sommer, 2014; Marsden et al., 2012) or may overwhelm the influence of selection in small populations (Luo, Pan, Liu, & Li, 2012; Miller & Lambert, 2004). In some cases, MHC diversity even declines faster than does neutral diversity (Bateson et al., 2015; Bollmer et al., 2011; Ejsmond & Radwan, 2009; Eimes et al., 2011; Sutton et al., 2011, 2015); however, we saw a more rapid decline of neutral diversity compared to MHC diversity in the captive ring-tailed lemur population at the DLC. Thus, although genetic drift has resulted in decreased neutral diversity in the DLC population, selective pressure may be mitigating the loss of immunogenetic diversity in captivity (Ellison et al., 2012; Newhouse & Balakrishnan, 2015). Captivity may even be exerting strong selective pressure through exposure to novel pathogens not previously experienced during millions of years of isolation on Madagascar.

Although limited MHC diversity does not necessarily condemn a species to extinction (e.g., Castro-Prieto, Wachter, & Sommer, 2011; Lau, Jaratlerdsiri, Griffith, Gongora, & Higgins, 2014; Plasil et al., 2016; Weber, Stewart, Schienman, & Lehman, 2004; Weber et al., 2013; Zhu, Ruan, Ge, Wan, & Fang, 2007), low MHC diversity can compromise

population viability over the long term by increasing susceptibility to disease (Belov, 2011; Radwan, Biedrzycka, & Babik, 2010; Segelbacher et al., 2014; Spielman, Brook, Briscoe, & Frankham, 2004). In particular, many species are likely to increasingly encounter novel pathogens in the future because of increased exposure to humans, their livestock, other domesticated animals (e.g., feral dogs and cats), and invasive species (Barrett, Brown, Junge, & Yoder, 2013; Daszak, Cunningham, & Hyatt, 2000, 2001). The negative effects of low MHC diversity on an animal's ability to combat novel pathogens also may be exacerbated by stress from human disturbance or environmental perturbations (Frankham, 2005b). These perturbations to the environment, which often decrease population size (e.g., a drought: Gould et al., 2003), are likely to subject populations to increasing effects of genetic drift, which may further decrease genetic diversity in a negative feedback loop that subsequently renders a population even more susceptible to novel pathogens. Thus, quantifying the current level of MHC diversity, especially in wild or endangered species, is essential to future conservation efforts.

Although previous researchers have estimated genetic diversity using nH_o as a proxy for functional diversity (Höglund, Wengström, Rogell, & Meyer-Lucht, 2015; Segelbacher et al., 2014), we found only a weak correlation between MHC-DRB diversity and nH_o in captivity and no correlation in our wild population. Our MHC and nH_o findings, however, confirm that captive individuals have less overall genetic diversity than do wild individuals (Charpentier et al., 2008; Pastorini et al., 2015). Captive ring-tailed lemurs are more closely related within institutions than between institutions, and also have greater relatedness between individuals at a single institution than do individuals within a social group at the BMSR (Pastorini et al., 2015). Together, the MHC and nH_o data indicate that groups of ring-tailed lemurs at some institutions have historically functioned as independent, small populations, exhibiting less gene flow than typically occurs between wild groups (Pastorini et al., 2015). In other words, there are fewer transfers between captive groups at the institutions examined in this study than there are immigrations between wild groups. Between 1980 and 2013, the AZA SSP included 2,839 ring-tailed lemurs housed at 289 institutions. During this period, 1,168 transfers, including some animals that were transferred multiple times, occurred between 217 institutions, which is a rate of 0.1 transfers per institution per year (Gina Ferrie, personal communication). In contrast, researchers at the BMSR and Berenty Reserve have observed an average of 1.4 and 2.9 transfers, respectively, per social group annually (Koyama, Nakamichi, Ichino, & Takahata, 2002; Sautner, 1991; Sussman, 1992).

In captivity, these transfers occur for several reasons, most importantly as part of the AZA SSP management of diversity of the captive population in the USA. The ring-tailed lemur population including all individuals from all AZA SSP institutions is a large and successfully managed population, considered a Green SSP Program because the population has retained "a minimum of 90% gene diversity at 100 years or 10 generations, and includes at least 50 animals held among at least three AZA member institutions" (AZA 2014). Estimates of genetic diversity are based on the number of founder lineages represented in the population, rather than on molecular estimates via microsatellites

or allelic diversity of SNPs or candidate genes (AZA Population Management Center Population Management Guidelines; Lacy, 1995). Unfortunately, little is known about the origins or relationships of the founding ring-tailed lemurs in the captive population, and molecular work has not confirmed that founders were unrelated to one another. AZA SSP management has prevented some loss of diversity through genetic drift; however, these efforts have not been completely successful at maintaining levels of neutral or adaptive diversity equivalent to the level of genetic diversity seen in the wild. Given our findings regarding the limited correlation between neutral and adaptive diversity, we recommend the inclusion of molecular data, particularly of potentially important candidate genes, into the Breeding and Transfer Plan. We also recommend increased transfer of key individuals possessing unique MHC-DRB alleles or substantial MHC-DRB diversity, particularly between international institutions that may hold unique individuals from different source populations (Frankham, 2015; Ivy, 2016; Pastorini et al., 2015). The inclusion of molecular data and the movement of endangered species across international borders may be financially and logistically constrained, but could increase the diversity of captive offspring and preserve extremely rare MHC-DRB alleles within the captive population.

Additionally, although uneven MHC-DRB allelic frequencies are present in other vertebrate species (Alcaide, Muñoz, Martínez-de la Puente, Soriguer, & Figuerola, 2014; Pechouskova et al., 2015; Weber et al., 2013), the highly skewed distribution of MHC-DRB alleles in ring-tailed lemurs increases the likelihood that rare, potentially important alleles may have been lost, particularly from the captive population, due to genetic drift. The diversity and distribution of MHC-DRB in animals across institutions should be evaluated and that assessment incorporated into breeding recommendations. To gain an accurate picture of MHC-DRB diversity in populations, especially captive ones, sampling should include almost every individual because of the abundance of rare alleles and uneven distribution.

Lastly, we show that captive populations suffer from the effects of small founder population and possible genetic drift, but that careful management, including the influx of new individuals, can negate this potential threat and lead to an increase in the number of MHC-DRB alleles. This kind of genetic “rescue” by a few individuals is one method of increasing genetic diversity, viability, and survivorship (e.g., Adams, Vucetich, Hedrick, Peterson, & Vucetich, 2011; Johnson et al., 2010; Westemeier et al., 1998; reviewed in Frankham, 2015). Without continued introduction of new alleles, however, the increase in individual diversity may be temporary (Hedrick, Peterson, Vucetich, Adams, & Vucetich, 2014). The inclusion of measures of genetic diversity into conservation management plans is a critical step in the preservation of viable populations, both in captivity and in the wild (Witzenberger & Hochkirch, 2011).

MHC-DRB is not the only component of immunogenetic diversity that could be monitored and managed, however. For example, toll-like receptors could provide an informative comparison to the MHC-DRB. Whereas the MHC controls the activation of the adaptive immune system, toll-like receptors are essential to the innate immune response through intracellular signaling (Gonzalez-Quevedo, Phillips,

Spurgin, & Richardson, 2015; Grueber et al., 2015; Morris, Wright, Grueber, Hogg, & Belov, 2015). Alternatively, assessment of functional immunogenetic diversity could be expanded using next-generation Ig-seq technology to explore expressed antibodies (Larsen, Campbell, & Yoder, 2014). This method allows for monitoring of disease status of wild populations and surveillance of novel diseases, an important component of advanced conservation management.

As an endangered strepsirrhine endemic to Madagascar, the ring-tailed lemur is both a flagship conservation species for one of the world's top biodiversity hot spots and a prime example of a species in peril (Gould & Sauther, 2016; LaFleur et al. 2016). Like all nonhuman primates, ring-tailed lemurs face significant anthropogenic threats (Reuter et al. 2017; Schwitzer et al., 2013) and are increasingly susceptible to environmental change through loss of genetic diversity (Frankham, 2005a, 2005b; Charpentier et al., 2008; Spielman et al., 2004). Studies such as this one are key for assessing a species' ability to respond to these threats, including novel pathogens, as well as to adapt to changing climatic conditions. Such studies are also crucial for determining whether captive populations are sufficiently representative of wild populations to serve as a reservoir for the purpose of reintroduction or repopulation.

ACKNOWLEDGMENTS

Funding for sample collection from captive animals and MHC genotyping was provided to CMD by the National Science Foundation (NSF, IOS-071900) and to CMD and KEG by the NSF (BCS-1232570), the Margot Marsh Biodiversity Foundation (MMBF), the Duke Primate Genomics Initiative, and Duke University. Funding for sample collection at the BMSR was provided to MLS and FPC by the NSF (BCS 0922465), the MMBF, the Universities of Colorado and North Dakota, UND SSAC, ND EPSCoR, Primate Conservation Inc., the International Primatological Society, the American Society of Primatologists, and the St. Louis Zoo (FRC 06-1). Funding for publication was provided by the Emory University Open Access Publishing Fund to KEG and Duke University to CMD. We would like to thank Erin Ehmke, Andrea Katz, and Charlie Welch for their input regarding captive management at the DLC, as well as Lynne Villers and Mark Campbell for their assistance in organizing sample collection at the Indianapolis Zoo and Cincinnati Zoo, respectively. The staff of the Duke Lemur Center, as well as the Indianapolis and Cincinnati Zoos, graciously facilitated sample collection. Similarly, we would like to acknowledge the census team at the BMSR, all of the veterinary and field assistants who helped with sample collection in Madagascar for the last two decades, and two anonymous reviewers for their helpful comments. Lastly, we are grateful to Gina Ferrie, the Species Survival Plan[®] Coordinator for the AZA ring-tailed lemur population, who generously provided the information on the entire AZA managed population, as well as relevant comments about the manuscript. This manuscript is DLC publication #1365.

CONFLICT OF INTEREST

The authors declare no conflict of interests.

AUTHOR CONTRIBUTIONS

CMD and KEG conceived the study and participated in study design and sample collection. KEG performed the molecular work and data analyses, as well as drafted the manuscript. CMD critically revised the manuscript drafts. MLS and FPC coordinated and collected the wild samples from Madagascar, as well as edited the manuscript. All authors approved the final manuscript.

DATA ACCESSIBILITY

The genotypes of each ring-tailed lemur, as well as location of origin, have been deposited in Dryad: <https://doi.org/10.5061/dryad.bn7h0>.

REFERENCES

- Adams, J. R., Vucetich, L. M., Hedrick, P. W., Peterson, R. O., & Vucetich, J. A. (2011). Genomic sweep and potential genetic rescue during limiting environmental conditions in an isolated wolf population. *Proceedings of the Royal Society B: Biological Sciences*, 278, 3336–3344.
- Agudo, R., Alcaide, M., Rico, C., Lemus, J. A., Blanco, G., Hiraldo, F., & Donazar, J. A. (2011). Major histocompatibility complex variation in insular populations of the Egyptian vulture: Inferences about the roles of genetic drift and selection. *Molecular Ecology*, 20, 2329–2340.
- Alcaide, M., Muñoz, J., Martínez-de la Puente, J., Soriguer, R., & Figuerola, J. (2014). Extraordinary MHC class II B diversity in a non-passerine, wild bird: The Eurasian Coot *Fulica atra* (Aves: Rallidae). *Ecology and Evolution*, 4, 688–698.
- Andriaholinirina, N., Baden, A., Blanco, M., Chikhi, L., Cooke, A., Davies, N., ... Zaramody, A. (2014). *Lemur catta*. The IUCN Red List of Threatened Species 2014: e.T11496A62260437. Downloaded 17 March 2016.
- Association of Zoos and Aquariums Population Management Center. *Population management guidelines*. Association of Zoos and Aquariums Population Management Center, Lincoln Park Zoo, Chicago, IL. Retrieved from www.lpzoo.org/sites/default/files/pdf/cs/pmc/Population%20Management%20Guidelines_916.pdf.
- Barrett, M. A., Brown, J. L., Junge, R. E., & Yoder, A. D. (2013). Climate change, predictive modeling and lemur health: Assessing impacts of changing climate on health and conservation in Madagascar. *Biological Conservation*, 157, 409–422.
- Bateson, Z. W., Whittingham, L. A., Johnson, J. A., & Dunn, P. O. (2015). Contrasting patterns of selection and drift between two categories of immune genes in prairie-chickens. *Molecular Ecology*, 24, 6095–6106.
- Belov, K. (2011). The role of the Major histocompatibility complex in the spread of contagious cancers. *Mammalian Genome*, 22, 83–90.
- Bernatchez, L., & Landry, C. (2003). MHC studies in nonmodel vertebrates: What have we learned about natural selection in 15 years? *Journal of Evolutionary Biology*, 16, 363–377.
- Biedrzycka, A., Kloch, A., Buczek, M., & Radwan, J. (2011). Major histocompatibility complex DRB genes and blood parasite loads in fragmented populations of the spotted suslik *Spermophilus suslicus*. *Mammalian Biology*, 76, 672–677.
- Bollmer, J. L., Ruder, E. A., Johnson, J. A., Eimes, J. A., & Dunn, P. O. (2011). Drift and selection influence geographic variation at immune loci of prairie-chickens. *Molecular Ecology*, 20, 4695–4706.
- Carrington, M., & Bontrop, R. E. (2002). Effects of MHC class I on HIV/SIV disease in primates. *Aids*, 16(Supplement 4), S105–S114.
- Castro-Prieto, A., Wachter, B., & Sommer, S. (2011). Cheetah paradigm revisited: MHC diversity in the world's largest free-ranging population. *Molecular Biology and Evolution*, 28, 1455–1468.
- Charpentier, M. J. E., Widdig, A., & Alberts, S. C. (2007). Inbreeding depression in non-human primates: A historical review of methods used and empirical data. *American Journal of Primatology*, 69, 1370–1386.
- Charpentier, M. J. E., Williams, C. V., & Drea, C. M. (2008). Inbreeding depression in ring-tailed lemurs (*Lemur catta*): Genetic diversity predicts parasitism, immunocompetence, and survivorship. *Conservation Genetics*, 9, 1605–1615.
- Charpentier, M. J. E., Williams, C. V., & Drea, C. M. (2015). Erratum to: Inbreeding depression in ring-tailed lemurs (*Lemur catta*): Genetic diversity predicts parasitism, immunocompetence, and survivorship. *Conservation Genetics*, 9, 1605–1615.
- Crnokrak, P., & Roff, D. A. (1999). Inbreeding depression in the wild. *Heredity*, 83, 260–270.
- Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging infectious diseases of wildlife—Threats to biodiversity and human health. *Science*, 287, 443–449.
- Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Tropica*, 78, 103–116.
- Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J., & Ovenden, J. R. (2014). NeEstimator V2: Re-implementation of software for the estimation of contemporary effective population size (*N_e*) from genetic data. *Molecular Ecology Resources*, 14(1), 209–214.
- Doytchinova, I. A., & Flower, D. R. (2005). In silico identification of super-types for class II MHCs. *The Journal of Immunology*, 174, 7085–7095.
- Eimes, J. A., Bollmer, J. L., Whittingham, L. A., Johnson, J. A., van Oosterhout, C., & Dunn, P. O. (2011). Rapid loss of MHC class II variation in a bottlenecked population is explained by drift and loss of copy number variation. *Journal of Evolutionary Biology*, 24, 1847–1856.
- Ejmond, M. J., & Radwan, J. (2009). MHC diversity in bottlenecked populations: A simulation model. *Conservation Genetics*, 12, 129–137.
- Ellison, A., Allainguillaume, J., Girdwood, S., Pachebat, J., Peat, K. M., Wright, P., & Consuegra, S. (2012). Maintaining functional major histocompatibility complex diversity under inbreeding: The case of a selfing vertebrate. *Proceedings of the Royal Society B: Biological Sciences*, 279, 5004–5013.
- Evans, M. L., & Neff, B. D. (2009). Major histocompatibility complex heterozygote advantage and widespread bacterial infections in populations of Chinook salmon (*Oncorhynchus tshawytscha*). *Molecular Ecology*, 18, 4716–4729.
- Fernández-de-Mera, I. G., Vicente, J., Pérez de la Lastra, J., Mangold, A. J., Naranjo, V., Fierro, Y., ... Gortázar, C. (2009). Reduced major histocompatibility complex class II polymorphism in a hunter-managed isolated Iberian red deer population. *Journal of Zoology*, 277, 157–170.
- Frankham, R. (2005a). Genetics and extinction. *Biological Conservation*, 126, 131–140.
- Frankham, R. (2005b). Stress and adaptation in conservation genetics. *Journal of Evolutionary Biology*, 18, 750–755.
- Frankham, R. (2015). Genetic rescue of small inbred populations: Meta-analysis reveals large and consistent benefits of gene flow. *Molecular Ecology*, 24, 2610–2618.
- Froeschke, G., & Sommer, S. (2014). Role of selection versus neutral processes determining genetic variation in a small mammal along a climatic gradient in southern Africa. *Evolutionary Ecology*, 28, 1169–1190.
- Gonzalez-Quevedo, C., Phillips, K. P., Spurgin, L. G., & Richardson, D. S. (2015). 454 screening of individual MHC variation in an endemic island passerine. *Immunogenetics*, 67, 149–162.
- Gould, L., & Sauther, M. L. (2016). Going, Going, Gone...Is the iconic ring-tailed lemur (*Lemur catta*) headed for imminent extirpation? *Primate Conservation*, 30, 89–101.
- Gould, L., Sussman, R. W., & Sauther, M. L. (1999). Natural disasters and primate populations: The effects of a 2-year drought on a naturally occurring population of ring-tailed lemurs (*Lemur catta*) in Southwestern Madagascar. *International Journal of Primatology*, 20, 69–84.
- Gould, L., Sussman, R. W., & Sauther, M. L. (2003). Demographic and life-history patterns in a population of ring-tailed lemurs (*Lemur catta*) at

- Beza Mahafaly Reserve, Madagascar: A 15-year perspective. *American Journal of Physical Anthropology*, 120, 182–194.
- Grogan, K. E., McGinnis, G. J., Sauther, M. L., Cuozzo, F. P., & Drea, C. M. (2016). Next-generation genotyping of hypervariable loci in many individuals of a non-model species: Technical and theoretical implications. *BMC Genomics*, 17, 204.
- Grueber, C. E., Knafler, G. J., King, T. M., Senior, A. M., Grosser, S., Robertson, B., ... Jamieson, I. G. (2015). Toll-like receptor diversity in 10 threatened bird species: Relationship with microsatellite heterozygosity. *Conservation Genetics*, 16, 595–611.
- Hartl, D. L., & Clark, A. G. (1997). *Principles of population genetics*. Sunderland, MA: Hedrick.
- Hedrick, P. W. (2001). Conservation genetics: Where are we now? *Trends in Ecology & Evolution*, 16, 629–636.
- Hedrick, P. W., & Kalinowski, S. T. (2011). Inbreeding depression in conservation biology. *Annual Review of Ecology & Systematics*, 31, 139–162.
- Hedrick, P. W., Peterson, R. O., Vucetich, L. M., Adams, J. R., & Vucetich, J. A. (2014). Genetic rescue in Isle Royale wolves: Genetic analysis and the collapse of the population. *Conservation Genetics*, 15, 1111–1121.
- Hill, W. G. (1981). Estimation of effective population size from data on linkage disequilibrium. *Genetical Research*, 38, 209–216.
- Höglund, J., Wengström, Å., Rogell, B., & Meyer-Lucht, Y. (2015). Low MHC variation in isolated island populations of the Natterjack toad (*Bufo calamita*). *Conservation Genetics*, 16, 1007–1010.
- Huchard, E., Knapp, L. A., Wang, J., Raymond, M., & Cowlshaw, G. (2010). MHC, mate choice and heterozygote advantage in a wild social primate. *Molecular Ecology*, 19, 2545–2561.
- Huchard, E., Weill, M., Cowlshaw, G., Raymond, M., & Knapp, L. A. (2008). Polymorphism, haplotype composition, and selection in the Mhc-DRB of wild baboons. *Immunogenetics*, 60, 585–598.
- IUCN (2010). *IUCN red list of threatened species. Version 2010.4*. Retrieved from www.iucnredlist.org. Downloaded on 7 February 2012.
- Ivy, J. A. (2016). Ameliorating the loss of genetic diversity in captive wildlife populations. *Animal Conservation*, 19, 112–113.
- Johnson, W. E., Onorato, D. P., Roelke, M. E., Land, E. D., Cunningham, M., Belden, R. C., ... O'Brien, S. J. (2010). Genetic restoration of the Florida panther. *Science*, 329, 1641–1645.
- Jolly, A. (1966). *Lemur behavior*. Chicago, IL: University of Chicago Press.
- Kalbe, M., Eizaguirre, C., Dankert, I., Reusch, T. B. H., Sommerfeld, R. D., Wegner, K. M., & Milinski, M. (2009). Lifetime reproductive success is maximized with optimal major histocompatibility complex diversity. *Proceedings of the Royal Society B: Biological Sciences*, 276, 925–934.
- Keller, L. F., & Waller, D. M. (2002). Inbreeding effects in wild populations. *Trends in Ecology & Evolution*, 17, 19–23.
- Knapp, L. A., Ha, J. C., & Sackett, G. P. (1996). Parental MHC antigen sharing and pregnancy wastage in captive pigtailed macaques. *Journal of Reproductive Immunology*, 32, 73–88.
- Koyama, N., Nakamichi, M., Ichino, S., & Takahata, Y. (2002). Population and social dynamics changes in ring-tailed lemur troops in Berenty, Madagascar between 1989–1999. *Primates*, 43, 291–314.
- Lacy, R. C. (1995). Clarification of genetic terms and their use in the management of captive populations. *Zoo Biology*, 14, 565–578.
- LaFleur, M., Clarke, T. A., Reuter, K., & Schaeffer, T. (2016). Rapid Decrease in Populations of Wild Ring-Tailed Lemurs (*Lemur catta*) in Madagascar. *Folia Primatologica*, 87, 320–330.
- Larsen, P. A., Campbell, C. R., & Yoder, A. D. (2014). Next-generation approaches to advancing eco-immunogenomic research in critically endangered primates. *Molecular Ecology Resources*, 14, 1198–1209.
- Lau, Q., Jaratlerdsiri, W., Griffith, J. E., Gongora, J., & Higgins, D. P. (2014). MHC class II diversity of koala (*Phascolarctos cinereus*) populations across their range. *Heredity*, 113, 287–296.
- Laurie-Ahlberg, C., & Weir, B. S. (1979). Allozyme variation and linkage disequilibrium in some laboratory populations of *Drosophila melanogaster*. *Genetical Research*, 32, 215–229.
- Luo, M. F., Pan, H. J., Liu, Z. J., & Li, M. (2012). Balancing selection and genetic drift at major histocompatibility complex class II genes in isolated populations of golden snub-nosed monkey (*Rhinopithecus roxellana*). *BMC Evolutionary Biology*, 12, 207.
- Madsen, T., Olsson, M., Wittzell, H., Stille, B., Gullberg, A., Shine, R., ... Tegelström, H. (2000). Population size and genetic diversity in sand lizards (*Lacerta agilis*) and adders (*Vipera berus*). *Biological Conservation*, 94, 257–262.
- Marsden, C. D., Woodroffe, R., Mills, M. G. L., McNutt, J. W., Creel, S., Groom, R., ... Mable, B. K. (2012). Spatial and temporal patterns of neutral and adaptive genetic variation in the endangered African wild dog (*Lycaon pictus*). *Molecular Ecology*, 21, 1379–1393.
- McKenney, E. A., Rodrigo, A., & Yoder, A. D. (2015). Patterns of gut bacterial colonization in three primate species. *PLoS One*, 10, e0124618.
- Miller, H. C., & Lambert, D. M. (2004). Genetic drift outweighs balancing selection in shaping post-bottleneck major histocompatibility complex variation in New Zealand robins (*Petroicidae*). *Molecular Ecology*, 13, 3709–3721.
- Morris, K. M., Wright, B., Grueber, C. E., Hogg, C., & Belov, K. (2015). Lack of genetic diversity across diverse immune genes in an endangered mammal, the Tasmanian devil (*Sarcophilus harrisii*). *Molecular Ecology*, 24, 3860–3872.
- Newhouse, D. J., & Balakrishnan, C. N. (2015). High major histocompatibility complex class I polymorphism despite bottlenecks in wild and domesticated populations of the zebra finch (*Taeniopygia guttata*). *BMC Evolutionary Biology*, 15, 265.
- Noble, S. J., Chesser, R. K., & Ryder, O. A. (1990). Inbreeding effects in captive populations of Ruffed Lemurs. *Human Evolution*, 5, 283–291.
- Ober, C. (1999). Studies of HLA, fertility and mate choice in a human isolate. *Human Reproduction Update*, 5, 103–107.
- Oliver, M. K., Telfer, S., & Pierny, S. B. (2009). Major histocompatibility complex (MHC) heterozygote superiority to natural multi-parasite infections in the water vole (*Arvicola terrestris*). *Proceedings of the Royal Society B: Biological Sciences*, 276, 1119–1128.
- Ouborg, N. J., Angeloni, F., & Vergeer, P. (2010). An essay on the necessity and feasibility of conservation genomics. *Conservation Genetics*, 11, 643–653.
- Pastorini, J., Sauther, M. L., Sussman, R. W., Gould, L., Cuozzo, F. P., Fernando, P., ... Mundy, N. I. (2015). Comparison of the genetic variation of captive ring-tailed lemurs with a wild population in Madagascar. *Zoo Biology*, 34, 463–472.
- Pechouskova, E., Dammhahn, M., Brameier, M., Fichtel, C., Kappeler, P. M., & Huchard, E. (2015). MHC class II variation in a rare and ecological specialist mouse lemur reveals lower allelic richness and contrasting selection patterns compared to a generalist and widespread sympatric congener. *Immunogenetics*, 67, 229–245.
- Pierny, S. B., & Oliver, M. K. (2006). The evolutionary ecology of the major histocompatibility complex. *Heredity*, 96, 7–21.
- Plasil, M., Mohandesan, E., Fitak, R. R., Musilova, P., Kubickova, S., Burger, P. A., & Horin, P. (2016). The major histocompatibility complex in Old World camels and low polymorphism of its class II genes. *BMC Genomics*, 17, 167.
- Radwan, J., Biedrzycka, A., & Babik, W. (2010). Does reduced MHC diversity decrease viability of vertebrate populations? *Biological Conservation*, 143, 537–544.
- Reed, D. H., & Frankham, R. (2001). How closely correlated are molecular and quantitative measures of genetic variation? A meta-analysis. *Evolution; International Journal of Organic Evolution*, 55, 1095–1103.
- Reuter, K. E., LaFleur, M., & Clarke, T. A. (2017). Endangered species: Illegal lemur trade grows in Madagascar. *Nature*, 154, 157.
- Sauermann, U., Nürnberg, P., Bercovitch, F., Berard, J. K., Trefilov, A., Widdig, A., ... Krawczak, M. (2001). Increased reproductive success of MHC class II heterozygous males among free-ranging rhesus macaques. *Human Genetics*, 108, 249–254.
- Sauther, M. L. (1991). Reproductive behavior of free-ranging Lemur catta at Beza Mahafaly Special Reserve, Madagascar. *American Journal of Physical Anthropology*, 84, 463–477.

- Sauther, M. L., & Cuzzo, F. P. (2008). Somatic variation in living, wild ring-tailed lemurs (*Lemur catta*). *Folia Primatologica*, 79, 55–78.
- Sauther, M. L., Gould, L., Cuzzo, F. P., & O'Mara, M. T. (2015). *Ring-tailed lemurs: A species re-imagined*. *Folia Primatologica*, 86, 5–13. Karger Publishers.
- Sauther, M. L., Sussman, R. W., & Gould, L. (1999). The socioecology of the ringtailed lemur: Thirty-five years of research. *Evolutionary Anthropology*, 8, 120–132.
- Savage, A. E., & Zamudio, K. R. (2011). MHC genotypes associate with resistance to a frog-killing fungus. *Proceedings of the National Academy of Sciences*, 108, 1–6.
- Schwensov, N., Fietz, J., Dausmann, K. H., & Sommer, S. (2007). Neutral versus adaptive genetic variation in parasite resistance: Importance of major histocompatibility complex supertypes in a free-ranging primate. *Heredity*, 99, 265–277.
- Schwitzer, C., Mittermeier, R., Davies, N., Johnson, S., Ratsimbazafy, J., Razafindramanana, J., ... Rajaobelina, S. (Eds.) (2013). *Lemurs of Madagascar: A strategy for their conservation 2013–2016*. Bristol, UK: IUCN SSC Primate Specialist Group, Bristol Conservation and Science Foundation, and Conservation International.
- Scordato, E. S., Dubay, G., & Drea, C. M. (2007). Chemical composition of scent marks in the ringtailed lemur (*Lemur catta*): Glandular differences, seasonal variation, and individual signatures. *Chemical Senses*, 32, 493–504.
- Segelbacher, G., Strand, T. M., Quintela, M., Axelsson, T., Jansman, H. A. H., Koelewijn, H. P., & Höglund, J. (2014). Analyses of historical and current populations of black grouse in Central Europe reveal strong effects of genetic drift and loss of genetic diversity. *Conservation Genetics*, 15, 1183–1195.
- Sepil, I., Lachish, S., Hinks, A. E., & Sheldon, B. C. (2013). Mhc supertypes confer both qualitative and quantitative resistance to avian malaria infections in a wild bird population. *Proceedings of the Royal Society B: Biological Sciences*, 280, 20130134.
- Setchell, J., Charpentier, M., Abbott, K., Wickings, E., & Knapp, L. A. (2010). Opposites attract: MHC-associated mate choice in a polygynous primate. *Journal of Evolutionary Biology*, 23, 136–148.
- Smith, S., Mang, T., De Bellocq, J. G., Schaschl, H., Zeitlhofer, C., Hackländer, K., & Suchentrunk, F. (2010). Homozygosity at a class II MHC locus depresses female reproductive ability in European brown hares. *Molecular Ecology*, 19, 4131–4143.
- Sommer, S. (2005). Major histocompatibility complex and mate choice in a monogamous rodent. *Behavioral Ecology and Sociobiology*, 58, 181–189.
- Spielman, D., Brook, B. W., Briscoe, D. A., & Frankham, R. (2004). Does inbreeding and loss of genetic diversity decrease disease resistance? *Conservation Genetics*, 5, 439–448.
- Sussman, R. W. (1991). Demography and social organization of free-ranging *Lemur catta* in the Beza Mahafaly Reserve, Madagascar. *American Journal of Physical Anthropology*, 84, 43–58.
- Sussman, R. W. (1992). Male life history and intergroup mobility among ringtailed lemurs (*Lemur catta*). *International Journal of Primatology*, 13, 395–413.
- Sussman, R. W., Richard, A. F., Ratsirarson, J., Sauther, M. L., Brockman, D. K., Gould, L., ... Cuzzo, F. P. (2012). *Beza Mahafaly Special Reserve: Long-Term Research on Lemurs in Southwestern Madagascar*. In P. M. Kappeler & D. P. Watts (Eds.), *Long-term field studies of primates* (pp. 45–66). Berlin, Germany: Springer.
- Sutton, J. T., Nakagawa, S., Robertson, B. C., & Jamieson, I. G. (2011). Disentangling the roles of natural selection and genetic drift in shaping variation at MHC immunity genes. *Molecular Ecology*, 20, 4408–4420.
- Sutton, J. T., Robertson, B. C., & Jamieson, I. G. (2015). MHC variation reflects the bottleneck histories of New Zealand passerines. *Molecular Ecology*, 24, 362–373.
- Thoß, M., Ilmonen, P., Musolf, K., & Penn, D. J. (2011). Major histocompatibility complex heterozygosity enhances reproductive success. *Molecular Ecology*, 20, 1546–1557.
- Ujvari, B., & Belov, K. (2011). Major histocompatibility complex (MHC) markers in conservation biology. *International Journal of Molecular Sciences*, 12, 5168–5186.
- Väli, U., Einarsson, A., Waits, L., & Ellegren, H. (2008). To what extent do microsatellite markers reflect genome-wide genetic diversity in natural populations? *Molecular Ecology*, 17, 3808–3817.
- Van Coillie, S., Galbusera, P., Roeder, A. D., Schempp, W., Stevens, J. M. G., Leus, K., ... Pereboom, Z. (2008). Molecular paternity determination in captive bonobos and the impact of inbreeding on infant mortality. *Animal Conservation*, 11, 306–312.
- Waples, R. S., & Do, C. (2010). Linkage disequilibrium estimates of contemporary Ne using highly variable genetic markers: A largely untapped resource for applied conservation and evolution. *Evolutionary Applications*, 3, 244–262.
- Weber, D. S., Stewart, B. S., Schienman, J., & Lehman, N. (2004). Major histocompatibility complex variation at three class II loci in the northern elephant seal. *Molecular Ecology*, 13, 711–718.
- Weber, D. S., Van Coeverden De Groot, P. J., Peacock, E., Schrenzel, M. D., Perez, D. A., Thomas, S., ... Desalle, R. (2013). Low MHC variation in the polar bear: Implications in the face of Arctic warming? *Animal Conservation*, 16, 671–683.
- Wegner, K. M., Kalbe, M., Kurtz, J., Reusch, T. B. H., & Milinski, M. (2003). Parasite selection for immunogenetic optimality. *Science*, 301, 24306.
- Westemeier, R., Brawn, J., Simpson, S., Esker, T. L., Jansen, R. W., Walk, J. W., ... Paige, K. N. (1998). Tracking the long-term decline and recovery of an isolated population. *Science*, 282, 1695–1698.
- Westerdahl, H., Waldenström, J., Hansson, B., Hasselquist, D., von Schantz, T., & Bensch, S. (2005). Associations between malaria and MHC genes in a migratory songbird. *Proceedings of the Royal Society B: Biological Sciences*, 272, 1511–1518.
- Witzenberger, K. A., & Hochkirch, A. (2011). Ex situ conservation genetics: A review of molecular studies on the genetic consequences of captive breeding programmes for endangered animal species. *Biodiversity and Conservation*, 20, 1843–1861.
- Yang, Z. (2007). PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, 24, 1586–1591.
- Yao, G., Zhu, Y., Wan, Q. H., & Fang, S. G. (2015). Major histocompatibility complex class II genetic variation in forest musk deer (*Moschus berezovskii*) in China. *Animal Genetics*, 46, 535–543.
- Zehr, S. M., Roach, R. G., Haring, D., Taylor, J., Cameron, F. H., & Yoder, A. D. (2014). Life history profiles for 27 strepsirrhine primate taxa generated using captive data from the Duke Lemur Center. *Scientific Data*, 1, 1–11.
- Zhu, L., Ruan, X. D., Ge, Y. F., Wan, Q. H., & Fang, S. G. (2007). Low major histocompatibility complex class II DQA diversity in the Giant Panda (*Ailuropoda melanoleuca*). *BMC Genetics*, 8, 29.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Grogan KE, Sauther ML, Cuzzo FP, Drea CM. Genetic wealth, population health: Major histocompatibility complex variation in captive and wild ring-tailed lemurs (*Lemur catta*). *Ecol Evol*. 2017;7:7638–7649. <https://doi.org/10.1002/ece3.3317>