



Draft Genome Sequence of *Enterococcus plantarum* Strain TRW2, Isolated from Lettuce

 Inhwan You,^{a,b} Sukjung Choi,^b Thomas R. Williams,^c Maria L. Marco,^c Eun Bae Kim^{a,b,d}

^aDepartment of Animal Life Science, Kangwon National University, Chuncheon, Kangwon-do, Republic of Korea

^bLaboratory of Microbial Genomics and Big Data, College of Animal Life Sciences, Kangwon National University, Chuncheon, Kangwon-do, Republic of Korea

^cDepartment of Food Science and Technology, University of California, Davis, California, USA

^dDivision of Applied Animal Science, Kangwon National University, Chuncheon, Kangwon-do, Republic of Korea

ABSTRACT We report the draft genome sequence for *Enterococcus plantarum* strain TRW2, isolated from the phyllosphere of romaine lettuce. The draft sequence consists of 3,383,441 bp, with a G+C content of 35.8% and 3,218 protein-coding genes. None of the 22,190 known antibiotic resistance genes were detected.

Enterococci, well-known opportunistic pathogens that are associated with numerous human infections, are naturally found in the gastrointestinal tracts of humans and animals in various environments (1). However, plant-associated *Enterococcus* strains have been investigated in a limited number of studies (2, 3). Lettuce is one of the most popular vegetables to be consumed raw. Characterization of microorganisms associated with lettuce is important for food safety, especially unknown species that are associated with plants.

Strain TRW2 was isolated from romaine lettuce grown in the Salinas Valley of California as part of the 100K Pathogen Genome Project (4), and this genome sequence was also part of the 100K Pathogen Genome Project using previously published methods (5–7).

The sonication was performed in sterilized peptone water to obtain bacteria from the lettuce. Genomic DNA was isolated from a single colony from a de Man-Rogosa-Sharp (MRS) agar plate and inoculated into MRS broth overnight at 30°C under anaerobic conditions. DNA was extracted using a whole-genome isolation kit (Qiagen, Valencia, CA). Fragmented genomic DNA was used for the library construction with the KAPA high-throughput (HTP) library preparation kit (catalog number KK8234, Boston, MA) on the Agilent Bravo NGS workstation (Santa Clara, CA). Library quantification was performed using the KAPA library quantification kit (catalog number KK4824).

The paired-end sequencing was performed with the BGI@UCDavis sequencing center (UC Davis, CA) using the Illumina HiSeq 2000 platform with PE100 (San Diego, CA). The 100-bp paired-end reads were assembled after being quality filtered with an in-house Perl script. Briefly, 4,195,682 reads (length \geq 70 bp) representing 262-fold coverage of the genome sequence, in which 95% of bases showed a quality score of 31 (Illumina 1.8+), were selected and assembled using Ray 2.3.1 with a k-mer size of 31 (8). The assembled genome sequence was annotated with the Rapid Annotation using Subsystem Technology server databases and the NCBI Prokaryotic Genome Annotation Pipeline (9, 10).

To identify the species of strain TRW2, 16S rRNA sequences, the *sodA* gene, and the genome sequence were compared with those of other *Enterococcus* species (Fig. 1). The

Citation You I, Choi S, Williams TR, Marco ML, Kim EB. 2019. Draft genome sequence of *Enterococcus plantarum* strain TRW2, isolated from lettuce. *Microbiol Resour Announc* 8:e01428-18. <https://doi.org/10.1128/MRA.01428-18>.

Editor Christina Cuomo, Broad Institute of MIT and Harvard University

Copyright © 2019 You et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Eun Bae Kim, itanimal@kangwon.ac.kr.

Received 21 October 2018

Accepted 17 December 2018

Published 7 February 2019




FIG 1 Overview of the *Enterococcus plantarum* TRW2 genome. 16S rRNA phylogenetic tree based on 16S rRNA gene sequences in other enterococcal species. *Lactobacillus plantarum* ATCC 14917 was included as an outline; T means type strain. Phylogenetic analysis was performed with the neighbor-joining method with MEGA version 7.0 (13), with a 1,000-bootstrap analysis after sequence alignment with ClustalW (14). The TRW2 strain clustered with three other *E. plantarum* strains.

most similarities were found with *E. plantarum* (99.93% for 16S rRNA, 98.7% for the *sodA* gene, and 86.3% for the genome sequence), which indicates that the species of the strain is *E. plantarum*.

The properties of the *E. plantarum* TRW2 draft genome sequence are presented in Table 1. The draft sequence consisted of 132 contigs (≥ 500 bp), including 3,383,441 bp with a G+C content of 35.8% and N_{50} length of 89,597 bp.

Strain TRW2 has none of 22,190 known antibiotic resistance genes which were collected in the Comprehensive Antibiotic Resistance Database (CARD) (11), as well as none of the known enterococcal virulence factor (VF) genes analyzed by a previously reported method (12). These present results contribute to an initial description of the genomes of the *E. plantarum* group.

TABLE 1 Genome sequence statistics for *E. plantarum* TRW2

Attribute	Value
Genome size (bp)	3,383,441
DNA coding (bp)	2,805,327
DNA G+C content (bp)	1,211,271
No. of DNA scaffolds	134
No. of total genes	3,409
No. of protein-coding genes	3,218
No. of RNA genes	60
No. of pseudogenes	131
No. of genes with function prediction	2,484
No. of genes assigned to clusters of orthologous groups	2,951
No. of genes with Pfam domains	2,478
No. of genes with signal peptides	3,413
No. of genes with transmembrane helices	728
No. of CRISPR repeats	2

Data availability. The genome sequence of *E. plantarum* strain TRW2 has been deposited in NCBI GenBank under BioSample number [SAMN07453839](https://www.ncbi.nlm.nih.gov/biosample/SAMN07453839) and BioProject number [PRJNA397329](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA397329). Sequence data have been deposited in the Sequence Read Archive under the accession number [SRP166833](https://www.ncbi.nlm.nih.gov/sra/SRP166833).

ACKNOWLEDGMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2016R1C1B2016246), Brain Korea 21 Plus Project.

We declare that we have no conflict of interest.

REFERENCES

- Gao W, Howden BP, Stinear TP. 2018. Evolution of virulence in *Enterococcus faecium*, a hospital-adapted opportunistic pathogen. *Curr Opin Microbiol* 41:76–82. <https://doi.org/10.1016/j.mib.2017.11.030>.
- Mundt JO. 1963. Occurrence of enterococci on plants in a wild environment. *Appl Microbiol* 11:141–144.
- Müller T, Ulrich A, Ott E-M, Müller M. 2001. Identification of plant-associated enterococci. *J Appl Microbiol* 91:268–278. <https://doi.org/10.1046/j.1365-2672.2001.01373.x>.
- Weimer BC. 2017. 100K Pathogen Genome Project. *Genome Announc* 5:e00594-17. <https://doi.org/10.1128/genomeA.00594-17>.
- Kong N, Thao K, Huang C, Appel M, Lappin S, Knapp L, Kelly L, Weimer B. 2014. Automated library construction using KAPA library preparation kits on the Agilent NGS workstation yields high-quality libraries for whole-genome sequencing on the Illumina platform. *Agilent Technologies*, Santa, Clara, CA.
- Jeannotte R, Lee E, Kong N, Ng W, Kelly L, Weimer B. 2014. High-throughput analysis of foodborne bacterial genomic DNA using Agilent 2200 TapeStation and genomic DNA ScreenTape system. *Agilent Technologies*, Santa, Clara, CA.
- Kong N, Ng W, Cai L, Leonardo A, Kelly L, Weimer B. 2014. Integrating the DNA integrity number (DIN) to assess genomic DNA (gDNA) quality control using the Agilent 2200 TapeStation system. Application note. *Agilent Technologies*, Santa, Clara, CA.
- Boisvert S, Laviolette F, Corbeil J. 2010. Ray: simultaneous assembly of reads from a mix of high-throughput sequencing technologies. *J Comput Biol* 17:1519–1533. <https://doi.org/10.1089/cmb.2009.0238>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJ, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The comprehensive antibiotic resistance database. *Antimicrob Agents Chemother* 57:3348–3357. <https://doi.org/10.1128/AAC.00419-13>.
- Kim EB, Marco ML. 2014. Nonclinical and clinical *Enterococcus faecium* strains, but not *Enterococcus faecalis* strains, have distinct structural and functional genomic features. *Appl Environ Microbiol* 80:154–165. <https://doi.org/10.1128/AEM.03108-13>.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870–1874. <https://doi.org/10.1093/molbev/msw054>.
- Thompson JD, Gibson TJ, Higgins DG. 2002. Multiple sequence alignment using ClustalW and ClustalX. *Curr Protoc Bioinformatics Chapter 2:Unit 2.3*. <https://doi.org/10.1002/0471250953.bi0203s00>.