



ORAL PRESENTATION

Open Access

Expression of the epithelial polymeric immunoglobulin receptor is decreased in allergic rhinitis and eosinophilic rhinosinusitis

Cloé Hupin^{1*}, Philippe Rombaux¹, Marylène Lecocq², Charles Pilette²

From 9th Symposium of Experimental Rhinology and Immunology of the Nose (SERIN 2013) Leuven, Belgium. 21-23 March 2013

Background

Transcytosis of immunoglobulin A (IgA) through polarized bronchial and sinonasal epithelial cells is mediated by the polymeric immunoglobulin receptor (pIgR), which represents the rate-limiting factor for this frontline protective mechanism in the airways. pIgR expression is decreased in COPD, lung cancer and nasopharyngeal carcinoma, while its role in sinonasal chronic inflammatory diseases has not been explored. The aim of this study was thus to assess pIgR expression in sinonasal mucosa of patient with chronic rhinosinusitis with (CRSwNP) or without polyps (CRSsNP) and in allergic rhinitis (AR), as well as IgA and SC (the released soluble part of the pIgR) in nasal secretions.

Methods

Nasal and ethmoidal biopsies, as well as nasal fluid, were collected from patients with CRSwNP, CRSsNP and AR, as compared to control subjects. pIgR expression was analyzed by RT-qPCR and immunohistochemistry. IgA and SC were measured in nasal secretions by ELISA. Quantification of mucosal eosinophils was performed following hematoxylin-eosin staining.

Results

RT-qPCR showed a significant reduction of pIgR expression in ethmoidal biopsies from CRSwNP ($p = 0,01$) and AR ($p = 0,04$). This reduction was confirmed at the protein level by immunohistochemistry, and resulted into reduced levels of SC and trends for reduced IgA in nasal secretions from these patients. Decreased pIgR expression

was mainly observed in patients with increased mucosal eosinophils.

Conclusion

Epithelial pIgR expression is decreased in patients with CRSwNP and AR, results in decreased SC (and IgA) in nasal secretions, and closely relates to Th2-type eosinophilic inflammation. Whether this defect leads to impaired defense of the upper airways against pathogens remains to explore.

Author details

¹Cliniques Universitaires Saint-Luc, Service d'Oto-Rhino-Laryngologie, Brussels, Belgium. ²Université Catholique de Louvain, IREC, Pole de Pneumologie, ORL & Dermatologie, Brussels, Belgium.

Published: 16 July 2013

doi:10.1186/2045-7022-3-S2-O8

Cite this article as: Hupin et al.: Expression of the epithelial polymeric immunoglobulin receptor is decreased in allergic rhinitis and eosinophilic rhinosinusitis. *Clinical and Translational Allergy* 2013 3(Suppl 2):O8.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Cliniques Universitaires Saint-Luc, Service d'Oto-Rhino-Laryngologie, Brussels, Belgium

Full list of author information is available at the end of the article