

Association of curry consumption with blood lipids and glucose levels

Youngjoo Kwon[§]

Department of Food Science and Engineering, Ewha Womans University, 52, Ewhayeodae-gil, Seodaemun-gu, Seoul 03760, Korea

BACKGROUND/OBJECTIVES: Curcumin, an active ingredient in turmeric, is highly consumed in South Asia. However, curry that contains turmeric as its main spice might be the major source of curcumin in most other countries. Although curcumin consumption is not as high in these countries as South Asia, the regular consumption of curcumin may provide a significant health-beneficial effect. This study evaluated whether the moderate consumption of curry can affect blood glucose and lipid levels that become dysregulated with age.

SUBJECTS/METHODS: This study used data obtained from the Korea National Health and Nutrition Examination Survey, conducted from 2012 to 2013, to assess curry consumption frequency as well as blood glucose and blood lipid levels. The levels of blood glucose and lipids were subdivided by age, sex, and body mass index, and compared according to the curry consumption level. The estimates in each subgroup were further adjusted for potential confounding factors, including the diagnosis of diseases, physical activity, and smoking.

RESULTS: After adjusting for the above confounding factors, the blood glucose and triglyceride levels were significantly lower in the moderate curry consumption group compared to the low curry consumption group, both in older (> 45) male and younger (30 to 44) female overweight individuals who have high blood glucose and triglyceride levels.

CONCLUSIONS: These results suggest that curcumin consumption, in an ordinary diet, can have health-beneficial effects, including being helpful in maintaining blood glucose and triglyceride levels that become dysregulated with age. The results should be further confirmed in future studies.

Nutrition Research and Practice 2016;10(2):212-220; doi:10.4162/nrp.2016.10.2.212; pISSN 1976-1457 eISSN 2005-6168

Keywords: Blood glucose, curcumin, curry, triglyceride, turmeric

INTRODUCTION

Curry rice is a popular dish in many countries, including Korea. The distinct yellow color of curry or turmeric, is primarily derived from a polyphenolic compound, curcumin [1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadien-3,5-dione]. The potential health-beneficial effect of turmeric and its active component, curcumin, has been recognized; as such, there is an increasing interest in turmeric and turmeric-enhanced products [1].

A number of preclinical studies have demonstrated that curcumin exerts various biological activities, including anti-inflammatory, lipid-lowering, and anticancer properties [2-7]. In addition, clinical studies indicate that curcumin enhances cardiovascular health and insulin sensitivity [8-12]. However, the dose required to achieve these biological activities may vary, depending on existing disease conditions [1]. For example, the anticancer effect of curcumin may require high concentrations (1.5 to 4 g/day) [13,14], whereas less than 0.5 g of curcumin may be effective in treating some inflammatory conditions [15,16].

Turmeric has been commonly used for flavoring and color in food preparation, and for the treatment of inflammatory conditions in South Asia [17], leading to the high consumption

in these regions. In most other countries, including Korea, where turmeric is not a main ingredient for food preparation, the consumption rate for turmeric is low [1]. In these countries, curry may be the major curcumin-containing food. Previous studies indicate that regular curry consumption may be beneficial especially in the elderly, who have dysregulated physiological functions [10], although the consumption level observed is not as high as in South Asia. Therefore, it might be valuable to evaluate whether curcumin consumption through ordinary diet can exert any health-beneficial effects in a population where turmeric is not a popular ingredient.

In this study, the consumption level of curry, a major curcumin-containing food, was assessed using the Korea National Health and Nutrition Examination Survey (KNHANES) Food Frequency Questionnaire (FFQ). This study evaluated whether the consumption of curry in the typical Korean diet can affect the blood glucose and lipid levels that are dysregulated in older individuals.

SUBJECTS AND METHODS

Study design

This study evaluated the relationship between curry consumption and blood lipid levels, blood glucose levels, and glycated

[§] Corresponding Author: Youngjoo Kwon, Tel. 82-2-3277-3103, Fax. 82-2-3277-4213, Email. Youngjoo.Kwon@ewha.ac.kr

Received: August 20, 2015, Revised: December 7, 2015, Accepted: December 18, 2015

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

hemoglobin (HbA_{1c}), using the data obtained from KNHANES, a nationwide survey program conducted by the Korea Center for Disease Control and Prevention to access the health and nutritional status of Koreans. KNHANES collected information on the socioeconomic status, health-related behavior, quality of life, anthropometric measure, biochemical and clinical profiles of approximately 10,000 individuals, aged one and older, every year [1,18].

Estimation of curry consumption

The curry consumption was estimated using the KNHANES FFQ data conducted in 2012 and 2013 where the food consumption of individuals aged 19 to 64 was collected. Curry rice was the only curry-related food among the food items surveyed. The midpoint of each category was taken from the pre-coded frequency category, and converted into the frequency of consumption per month. The frequency of curry consumption per month was multiplied by the self-reported portion size, to derive the monthly mean intake of curry. According to the frequency of their curry consumption, subjects were divided into three groups: the low consumption group (LC; "almost never", or "once a month"), the moderate consumption group (MC; "2-3 times a month" or "once a week"), and the high consumption group (HC; "2-4 times per week", or "5-6 times per week").

Statistical analysis

SAS software (version 9.3, SAS Institute Inc., Cary, NC) was used for the data preparation and statistical analyses. Two years (2012 and 2013) of KNHANES data were combined to assess the characteristics of individuals, according to the level of curry consumption, and to evaluate the relationship between curry consumption and blood glucose levels or lipid profiles. A multi-stage sampling design was considered for all data generation and analysis. Since only 48 individuals were included in the HC, they were excluded from the statistical analysis. Chi-squared tests and *t*-tests were performed to determine the differences in the demographics and health-related characteristics between the LC and MC groups. The resulting *P*-values less than 0.05 were considered significant.

Individuals aged 30 and over were subjected to a comparison of their health-related characteristics. For the comparison of blood test measurements according to their curry consumption level, individuals were further stratified to control for the known confounding factors of age (young: 30 to 44; old: 45 to 64), sex (male/female), and body mass index (BMI; underweight: < 18.5; normal: 18.5 to 24.9; overweight: ≥ 25). Since the BMI in the underweight category was less (i.e. certain age-sex subgroups contained less than 10 individuals), these individuals were eliminated from the statistical analysis. Potential confounding variables other than age, sex, and BMI were examined using an analysis of covariance; they included dietary supplement use (yes, no), hypertension (yes, no, borderline hypertension), diabetes (yes, no, prediabetes), dyslipidemia (yes, no), instant noodle consumption (low: "almost never", "once a month", or "2-3 times per month"; high: "one time per week", "2-4 times per week", or "5-6 times per week"), tobacco use (never, past, current use), and physical activities (low: "never", or "one day per week";

intermediate: "2", "3", or "4 days per week"; high: "5", "6", or "7 days per week"). Non-responders for the above variables were excluded from the analysis. A multivariate analysis was conducted for the identified confounders, and adjusted means were estimated for each blood test measurement. A significance level was set at *P*-values less than 0.05.

RESULTS

Estimation of curry consumption by FFQ

Curry rice, simply "curry", was the only curry-related food item surveyed. Half of the population almost never consumed curry over the course of a single year (Table 1). Individuals who consumed curry mostly had it once a month, or 2-3 times a month, with very few (< 1%) having it more than once a week (Table 1). Nobody reported consuming curry more than once in a day. The majority of subjects (80%) reported the intake of 1 serving of curry each time (Table 1). The monthly curry consumption was lower at the age of 50 and over (0.51 servings per month) compared to the younger ages (about 1 serving per month) among those individuals who consumed curry (Table 2).

Demographic and health-related characteristics of individuals with different levels of curry consumption

Subjects were divided into three groups according to their frequency of curry consumption. Their demographic characteristics are described in Table 3. The estimated mean curry consumption was 0.33, 2.80, and 13.70 servings per month in the LC, MC, and HC, respectively. These were equivalent to the consumption of less than 0.01, 0.05, and 0.26 g of turmeric per day, respectively. A high frequency of curry consumption occurred in the younger age group; individuals between the

Table 1. The percentage of individuals in each category of the pre-coded frequency and portion size in the curry consumption frequency questionnaire

Frequency category	Total (n = 7,634) %	Frequency per month ¹⁾	Portion size category (serving)	Total (n = 7,634) %
Almost never	50.17	0 (0 × 1)	Half	5.30
1 time per month	25.15	1 (1 × 1)	One	34.25
2-3 times per month	13.63	2.5 (2.5 × 1)	One and half	3.49
1 time per week	3.52	4.3 (1 × 4.3)	None	50.17
2-4 times per week	0.68	12.9 (3 × 4.3)	No response	6.79
5-6 times per week	0.05	23.65 (5.5 × 4.3)		
No response	6.79			

¹⁾The midpoint of each category was taken and converted into the frequency of consumption per month (4.3 weeks in a month).

Table 2. Curry consumption per month in different age groups

Age	Mean monthly intake ¹⁾ (serving)
19-29	1.03 ± 0.07
30-39	1.12 ± 0.05
40-49	0.94 ± 0.04
50-64	0.51 ± 0.03

¹⁾The converted frequency of curry consumption per month (Table 1) was multiplied by the self-reported portion size to derive the monthly mean intake.

Table 3. The demographic characteristics of individuals with different levels of curry consumption

Characteristics	Low (LC), %	Moderate (MC), %	High (HC), %	P-value ¹⁾
Intake (serving/month)	0.33 ± 0.01 ²⁾ (< 0.01 g turmeric /day ³⁾)	2.80 ± 0.03 (0.05 g turmeric /day)	13.70 ± 0.76 (0.26 g turmeric /day)	< 0.0001
Age	41.5 ± 0.22	38.3 ± 0.37	35.4 ± 2.16	< 0.0001
19-29	21.80	23.35	45.20	
30-39	22.11	32.06	21.88	
40-49	25.13	27.54	18.88	
50-64	30.96	17.05	14.05	
Sex				0.0088
Male	51.13	46.61	47.65	
Female	48.87	53.39	52.35	
Residential area				0.0979
Urban	83.25	86.05	89.84	
Rural	16.75	13.95	10.16	
Residential type				0.0017
High-rise apartment	58.50	52.37	60.04	
Non-apartment residence	41.50	47.63	39.96	
Income quartile				0.1939
Lowest	25.61	24.76	43.04	
Medium lowest	26.62	23.62	8.11	
Medium highest	23.99	25.70	30.65	
Highest	23.78	25.93	18.20	
Education ⁴⁾				< 0.0001
Primary school or less	15.36	30.57	11.78	
Middle school	6.82	4.78	7.38	
High school	14.45	10.38	15.53	
University or above	63.37	54.27	65.31	

¹⁾ The significance of the differences in the frequencies and means between the low and moderate curry consumption groups was tested using *t*-test and chi-squared tests, respectively.

²⁾ Mean ± SE

³⁾ Daily turmeric consumption based on about 0.56 g turmeric in one serving of curry rice and 30 days per month.

⁴⁾ Frequency was age-adjusted.

ages of 30 and 39 were frequent in the MC (32.1%), while individuals between the ages of 19 and 29 were frequent in the HC (45.2%). In comparison, these age groups comprised less than 23% of the LC. The HC resembles the MC more than the LC in its demographic characteristics, with the exception of the type of residence and education level (Table 3). Due to the low frequency (only 48 individuals), the HC was eliminated from the statistical analysis. The MC included more females (53.4%) than the LC (48.9%). Individuals in the MC more frequently resided in non-apartment residences (47.6%) and had a lower education (54.3% in the “university or above” category) in comparison to 41.5 and 63.4%, respectively, in the LC. There were no significant differences in the residential area and individual income between the two groups (Table 3).

Subjects aged 30 and older were examined for their health-related characteristics (Table 4). The mean age was significantly lower in the MC (42.7) compared to the LC (46.3). Significantly more individuals had diagnoses of hypertension (24.9), hyper-

Table 4. The health-related characteristics of individuals aged 30 and older with different levels of curry consumption

Characteristics	Low (LC), %	Moderate (MC), %	P-value ¹⁾
Age (Mean ± SE)	46.3 ± 0.20	42.7 ± 0.30	< 0.0001
Hypertension			< 0.0001
No	48.05	57.39	
Borderline hypertension	27.09	23.36	
Yes	24.86	19.25	
Hypercholesterolemia			0.0027
No	85.57	88.75	
Yes	14.43	11.25	
Hypoalphalipoproteinemia			0.1910
No	76.80	79.42	
Yes	23.19	20.58	
Hypertriglyceridemia			0.0010
No	81.37	87.61	
Yes	18.63	12.39	
Diabetes			< 0.0001
No	68.09	76.92	
Prediabetes	23.53	17.25	
Yes	8.34	5.83	
Dietary supplement use ²⁾			0.0022
Yes	47.67	53.66	
No	52.33	46.34	
Vigorous exercise ³⁾			0.2171
Low	74.42	71.25	
Intermediate	19.87	22.14	
High	5.71	6.61	
Moderate exercise ⁴⁾			0.0349
Low	74.10	69.38	
Intermediate	19.42	23.17	
High	6.48	7.46	
Walking ⁵⁾			0.0124
Low	24.36	20.01	
Intermediate	34.53	34.10	
High	41.12	45.89	
Tobacco use ⁶⁾			0.0003
Never	52.33	60.05	
Past	20.65	19.22	
Current	27.02	20.73	
Instant noodle consumption ⁷⁾			< 0.0001
Low	59.83	49.23	
High	40.17	50.77	

¹⁾ The significance of the differences in the frequencies and means between the low and moderate curry consumption group was tested using *t*-test and chi-squared tests, respectively.

²⁾ Dietary supplement use during the last year

³⁻⁵⁾ Low: “never” or “one day per week”; intermediate: 2-4 days per week; high: 5-7 days per week

⁶⁾ Never: never user; past: past user; current: current user

⁷⁾ Low: “almost never”, “once a month”, or “2-3 times per month”; high: “once a week”, “2-4 times a week”, or “5-6 times a week”

cholesterolemia (14.4%), hypertriglyceridemia (18.6%) and diabetes (8.3%) in the LC than the MC (19.3, 11.3, 12.4, and 5.8%, respectively) (Table 4). The MC had a significantly higher frequency of dietary supplement use (53.7%), moderate exercise (30.6%), and walking (80.0%) compared to the LC (47.7, 25.9,

75.7%, respectively), whereas non-tobacco users were more frequent in the MC (60.1%) compared to the LC (52.3%). The frequency of instant noodle consumption was significantly higher in the MC (50.8%) than the LC (40.2%) (Table 4).

Anthropometric and blood test measurements of individuals with different levels of curry consumption

Subjects aged 30 and older were stratified into 12 groups, according to their age, sex, and BMI. Those who were underweight (BMI < 18.5) were eliminated due to the low numbers. Tables 5 and 6 describe the anthropometric and blood test measurements in the remaining groups. The mean age was not significantly different according to the curry consumption level except for the old-female-normal weight subgroup (Table 5). The anthropometric measurements were not significantly different according to the curry consumption level. However, in the old-male-overweight subgroup, individuals in the MC had lower ($P = 0.0591$) waist circumferences than the individuals in the LC (Table 5). The total cholesterol level was significantly higher in the old-female-normal weight subgroup, while the triglyceride (TG) level was significantly lower in the old-male-overweight subgroup, according to curry consumption (Table 6). The level of HbA_{1c} was lower ($P = 0.0548$) according to the curry consumption in the young-male-normal weight subgroup. There were no significant differences in blood levels of glucose and HDL cholesterol according to the curry consumption in any of the subgroups (Table 6). The compliance rate for the measurement of LDL cholesterol was very low; therefore, its statistical significance could not be considered (data not shown).

Since blood levels of glucose and lipids are influenced by disease conditions (e.g. diabetes and dyslipidemia), physical activity, and tobacco use, an analysis of covariance was conducted

to identify potential covariates for each blood measurement. The diagnosis of diabetes and moderate exercise were confounders for both the glucose and HbA_{1c} levels (Table 7). Additionally, glucose levels were confounded by hypertriglyceridemia, and HbA_{1c} levels were modulated by hypercholesterolemia and tobacco use (Table 7). The TG levels were confounded by the diagnoses of all tested disease conditions (hypertension, dyslipidemia, and diabetes) and tobacco use (Table 7). The total cholesterol level was similarly influenced as observed in the TG levels except for the tobacco use (Table 7). The HDL cholesterol level was modulated by the diagnosis of all tested conditions except hypertension, tobacco use, and instant noodle consumption (Table 7). The use of dietary supplements, vigorous exercise, and walking did not significantly affect the levels of any of the tested measurements (Table 7). After the adjustments for the identified confounding variables, the difference in the glucose levels according to curry consumption became significant, while the difference in the TG levels between the two curry consumption groups became insignificant in the old-male-overweight individuals (Table 8). Adjusted means of TG levels were significantly lower according to the curry consumption in the young-female-overweight subgroup (Table 8). HDL levels in the old-female-overweight subgroups were significantly lower in the MC compared to the LC after the adjustment (Table 8). The adjusted P -values for the differences in the blood glucose and HbA_{1c} levels according to the curry consumption were 0.0646 and 0.0490, respectively, in young-male-normal individuals, which are lower than they were when they were unadjusted. On the other hand, the difference in the total cholesterol level in the old-female-normal weight subgroup according to the curry consumption became insignificant after the adjustment (Table 8).

Table 5. Anthropometric measurements of individuals aged 30 and older with different levels of curry consumption

Variable	Age ¹⁾	Sex	BMI ²⁾	Low (LC)		Moderate (MC)		P-value ³⁾
				n	Mean ± SE	n	Mean ± SE	
Intake (serving/month)	Young	Male	Normal	430	0.45 ± 0.03	119	2.96 ± 0.09	
			Over	350	0.43 ± 0.03	82	3.08 ± 0.13	
		Female	Normal	810	0.42 ± 0.02	348	2.74 ± 0.07	
			Over	267	0.37 ± 0.04	71	2.56 ± 0.15	
	Old	Male	Normal	709	0.24 ± 0.02	91	2.80 ± 0.12	
			Over	455	0.25 ± 0.02	56	3.15 ± 0.22	
		Female	Normal	1,126	0.22 ± 0.01	197	2.43 ± 0.09	
			Over	596	0.22 ± 0.02	92	2.69 ± 0.12	
Age	Young	Male	Normal	430	37.1 ± 0.23	119	37.3 ± 0.51	0.8100
			Over	350	37.3 ± 0.03	82	36.8 ± 0.13	0.3916
		Female	Normal	810	37.2 ± 0.21	348	37.4 ± 0.28	0.5684
			Over	267	37.9 ± 0.34	71	37.9 ± 0.55	0.8973
	Old	Male	Normal	709	53.5 ± 0.25	91	52.1 ± 0.57	0.0259
			Over	455	53.1 ± 0.31	56	51.9 ± 0.76	0.1672
		Female	Normal	1,126	53.6 ± 0.21	197	51.5 ± 0.39	< 0.0001
			Over	596	54.1 ± 0.28	92	53.4 ± 0.73	0.3572
Height (cm)	Young	Male	Normal	430	172.9 ± 0.31	119	173.6 ± 0.56	0.3308
			Over	350	173.0 ± 0.34	82	173.9 ± 0.77	0.2589
		Female	Normal	810	160.3 ± 0.21	348	159.7 ± 0.37	0.2120
			Over	267	159.5 ± 0.48	71	160.4 ± 0.60	0.2347

Table 5. continued

Variable	Age ¹⁾	Sex	BMI ²⁾	Low (LC)		Moderate (MC)		P-value ³⁾	
				n	Mean ± SE	n	Mean ± SE		
Weight (kg)	Old	Male	Normal	709	168.9 ± 0.25	91	170.2 ± 0.65	0.0701	
			Over	455	169.1 ± 0.32	56	170.5 ± 0.90	0.1226	
		Female	Normal	1,126	156.6 ± 0.19	197	156.8 ± 0.43	0.6177	
			Over	596	155.4 ± 0.25	92	156.0 ± 0.69	0.4055	
	Young	Male	Normal	430	67.21 ± 0.37	119	67.78 ± 0.72	0.4879	
			Over	350	83.23 ± 0.58	82	83.79 ± 1.20	0.6751	
		Female	Normal	810	55.43 ± 0.21	348	55.27 ± 0.35	0.7049	
			Over	267	71.46 ± 0.74	71	71.93 ± 1.20	0.7441	
	WC (cm) ⁴⁾	Old	Male	Normal	709	64.65 ± 0.28	91	66.04 ± 0.82	0.1128
				Over	455	77.82 ± 0.38	56	78.88 ± 1.06	0.3339
			Female	Normal	1,126	54.92 ± 0.19	197	54.81 ± 0.47	0.8290
				Over	596	67.67 ± 0.47	92	67.41 ± 1.57	0.8761
Young		Male	Normal	430	78.85 ± 0.32	118	78.92 ± 0.58	0.9150	
			Over	350	91.93 ± 0.44	82	90.70 ± 0.94	0.2358	
		Female	Normal	809	72.90 ± 0.23	347	72.80 ± 0.32	0.7968	
			Over	266	88.23 ± 0.61	71	87.83 ± 1.11	0.7523	
BMI (kg/m ²)		Old	Male	Normal	708	81.16 ± 0.30	91	81.90 ± 0.69	0.3242
				Over	455	91.56 ± 0.31	55	90.11 ± 0.72	0.0591
			Female	Normal	1,126	75.91 ± 0.22	197	75.09 ± 0.44	0.0847
				Over	595	89.03 ± 0.48	92	86.97 ± 1.38	0.1561
	Young	Male	Normal	430	22.45 ± 0.09	119	22.47 ± 0.17	0.9065	
			Over	350	27.77 ± 0.14	82	27.66 ± 0.32	0.7425	
		Female	Normal	810	21.58 ± 0.07	348	21.64 ± 0.09	0.5736	
			Over	267	28.05 ± 0.21	71	27.95 ± 0.42	0.8375	
	Old	Male	Normal	709	22.64 ± 0.07	91	22.78 ± 0.21	0.5372	
			Over	455	27.19 ± 0.09	56	27.07 ± 0.21	0.6298	
		Female	Normal	1,126	22.39 ± 0.06	197	22.28 ± 0.14	0.4525	
			Over	596	28.00 ± 0.19	92	27.60 ± 0.47	0.4386	

¹⁾ Young: 30 ≤ age < 45; old: 45 ≤ age < 65

²⁾ Underweight: BMI < 18.5; normal: 18.5 ≤ BMI < 24.9; overweight: BMI ≥ 25

³⁾ The significance of the differences in the means between low and moderate curry consumption was tested using *t*-tests.

⁴⁾ Waist circumference

Table 6. Blood measurements of lipids, glucose, and HbA_{1c} in individuals aged 30 and older with different levels of curry consumption

Variable	Age ¹⁾	Sex	BMI ²⁾	Low (LC)		Moderate (MC)		P-value ³⁾	
				n	Mean ± SE	n	Mean ± SE		
Glucose (mg/dL) ⁴⁾	Young	Male	Normal	420	95.15 ± 1.09	116	92.42 ± 1.02	0.0988	
			Over	339	99.34 ± 1.23	79	96.38 ± 1.38	0.1068	
		Female	Normal	772	91.45 ± 0.52	323	90.44 ± 0.43	0.1332	
			Over	252	99.50 ± 1.64	69	103.46 ± 5.57	0.5073	
	Old	Male	Normal	688	102.62 ± 0.90	88	107.68 ± 4.08	0.2305	
			Over	441	109.09 ± 1.25	50	103.79 ± 3.24	0.1250	
		Female	Normal	1,068	96.68 ± 0.83	191	98.97 ± 2.83	0.4352	
			Over	560	104.96 ± 1.46	89	113.39 ± 6.82	0.2283	
HbA _{1c} (mg/dL) ⁵⁾	Young	Male	Normal	420	5.61 ± 0.04	116	5.49 ± 0.04	0.0548	
			Over	339	5.75 ± 0.04	79	5.71 ± 0.06	0.5097	
		Female	Normal	772	5.48 ± 0.02	323	5.47 ± 0.02	0.8177	
			Over	252	5.80 ± 0.07	68	5.89 ± 0.19	0.6768	
	Old	Male	Normal	688	5.89 ± 0.04	88	5.92 ± 0.13	0.8319	
			Over	441	6.03 ± 0.04	50	5.99 ± 0.12	0.7707	
			Female	Normal	1,068	5.80 ± 0.03	191	5.81 ± 0.09	0.9347
		Old	Female	Normal	1,068	5.80 ± 0.03	191	5.81 ± 0.09	0.9347
				Over	559	6.09 ± 0.05	89	6.33 ± 0.22	0.2738

Table 6. continued

Variable	Age ¹⁾	Sex	BMI ²⁾	Low (LC)		Moderate (MC)		P-value ³⁾
				n	Mean ± SE	n	Mean ± SE	
TG (mg/dL) ⁶⁾	Young	Male	Normal	421	140.43 ± 5.82	116	126.95 ± 12.21	0.3400
			Over	339	219.46 ± 10.89	79	223.23 ± 25.08	0.8950
		Female	Normal	772	92.25 ± 3.06	323	86.24 ± 4.07	0.2084
			Over	253	141.81 ± 9.22	69	119.48 ± 9.34	0.0862
	Old	Male	Normal	688	154.57 ± 4.78	88	181.10 ± 27.37	0.3434
			Over	441	199.71 ± 9.14	50	162.84 ± 12.22	0.0182
	Female	Normal	1,068	121.29 ± 2.90	191	116.21 ± 7.22	0.4200	
	Over	562	148.86 ± 5.06	89	200.47 ± 49.74	0.3034		
Cholesterol (mg/dL) ⁷⁾	Young	Male	Normal	421	188.69 ± 2.03	116	186.90 ± 3.24	0.6278
			Over	339	199.90 ± 2.39	79	202.62 ± 7.05	0.7161
		Female	Normal	772	179.23 ± 1.21	323	176.69 ± 2.07	0.2448
			Over	253	192.50 ± 3.08	69	196.74 ± 6.27	0.5448
	Old	Male	Normal	688	189.72 ± 1.64	88	192.46 ± 4.85	0.6005
			Over	441	192.31 ± 2.20	50	194.10 ± 5.12	0.7498
	Female	Normal	1,068	198.73 ± 1.22	191	205.23 ± 2.95	0.0434	
	Over	562	203.19 ± 1.79	89	204.78 ± 6.00	0.7967		
HDL (mg/dL) ⁸⁾	Young	Male	Normal	421	48.00 ± 0.53	116	49.02 ± 1.14	0.3972
			Over	339	43.20 ± 0.56	79	43.16 ± 1.27	0.9803
		Female	Normal	772	54.67 ± 0.37	323	53.49 ± 0.69	0.1848
			Over	253	48.75 ± 0.72	69	50.52 ± 1.67	0.3158
	Old	Male	Normal	688	46.48 ± 0.50	88	46.55 ± 1.30	0.9595
			Over	441	42.67 ± 0.46	50	43.63 ± 1.44	0.5125
	Female	Normal	1,068	51.82 ± 0.37	191	52.26 ± 0.94	0.6598	
	Over	562	48.73 ± 0.54	89	46.54 ± 1.16	0.0691		

¹⁻³⁾ Refers to Table 5.

⁴⁻⁶⁾ Were measured during a fasting state.

⁶⁾ Triglyceride levels

⁸⁾ High density lipoprotein cholesterol levels

Table 7. The determination of confounding variables for blood measurement by an analysis of covariance

Correlates	Glucose	HbA _{1c}	TG	Cholesterol	HDL
Age ¹⁾	0.0247	< 0.0001	0.8585	< 0.0001	0.2568
Sex	0.8376	0.2166	0.0010	0.3775	< 0.0001
BMI ²⁾	0.0079	0.0002	0.0267	< 0.0001	< 0.0001
Hypertension	0.0928	0.5600	0.0001	< 0.0001	0.4125
Hypercholesterolemia	0.3227	0.0002	0.0043	< 0.0001	< 0.0001
Hypoalphalipoproteinemia	0.4259	0.4130	< 0.0001	< 0.0001	< 0.0001
Hypertriglyceridemia	0.0093	0.7206	< 0.0001	< 0.0001	< 0.0001
Diabetes	< 0.0001	< 0.0001	0.0305	< 0.0001	< 0.0001
Supplement use ³⁾	0.9910	0.8761	0.7448	0.3131	0.2835
Vigorous exercise ⁴⁾	0.1541	0.1244	0.9978	0.0619	0.1368
Moderate exercise ⁵⁾	0.0111	0.0183	0.6914	0.0537	0.0702
Walking ⁶⁾	0.7382	0.4907	0.7037	0.9414	0.5262
Tobacco use ⁷⁾	0.6724	0.0046	0.0026	0.4060	0.0029
Instant noodle consumption ⁸⁾	0.4235	0.4483	0.6620	0.1442	0.0096

¹⁾ Young: 30 ≤ age < 45; old: 45 ≤ age < 65

²⁾ Normal: 18.5 ≤ BMI < 24.9; over: BMI ≥ 25

³⁾ Dietary supplement use during the last year

⁴⁻⁶⁾ Low: "never" or "one day per week"; intermediate: 2-4 days per week; high: 5-7 days per week

⁷⁾ Tobacco use

⁸⁾ Low: "almost never", "once per month" or "2-3 times per month"; high: "once per week", "2-4 times per week", or "5-6 times per week"

Table 8. The blood measurements of lipids, glucose, and HbA_{1c} in individuals aged 30 and older with different levels of curry consumption after adjustments for identified covariates

Variable	Age ¹⁾	Sex	BMI ²⁾	Low (LC)	Moderate (MC)	P-value ³⁾
				Mean ± SE	Mean ± SE	
Glucose (mg/dL) ⁴⁾	Young	Male	Normal	98.58 ± 1.57	95.93 ± 1.08	0.0646
			Over	98.33 ± 0.99	98.64 ± 1.56	0.8404
		Female	Normal	97.95 ± 1.47	97.70 ± 1.47	0.6766
			Over	99.27 ± 0.86	99.61 ± 4.31	0.9373
	Old	Male	Normal	98.25 ± 0.53	98.78 ± 1.90	0.7938
			Over	100.69 ± 0.77	94.58 ± 2.54	0.0289*
		Female	Normal	98.10 ± 0.74	102.05 ± 2.60	0.1125
			Over	98.94 ± 0.89	105.34 ± 4.54	0.1822
HbA _{1c} (%) ⁵⁾	Young	Male	Normal	5.71 ± 0.05	5.63 ± 0.04	0.0490*
			Over	5.68 ± 0.04	5.74 ± 0.05	0.1821
		Female	Normal	5.65 ± 0.05	5.66 ± 0.06	0.6506
			Over	5.82 ± 0.05	5.69 ± 0.13	0.3501
	Old	Male	Normal	5.78 ± 0.04	5.77 ± 0.10	0.9378
			Over	5.84 ± 0.04	5.82 ± 0.09	0.7946
		Female	Normal	5.77 ± 0.06	5.81 ± 0.08	0.4841
			Over	5.94 ± 0.04	6.11 ± 0.14	0.2417
TG (mg/dL) ⁶⁾	Young	Male	Normal	146.74 ± 5.74	145.01 ± 11.41	0.8531
			Over	155.44 ± 8.06	163.68 ± 14.51	0.6812
		Female	Normal	129.51 ± 5.53	124.35 ± 5.33	0.0630
			Over	145.37 ± 6.47	123.54 ± 6.06	0.0020*
	Old	Male	Normal	143.03 ± 5.18	161.02 ± 17.42	0.3224
			Over	142.66 ± 6.37	134.36 ± 10.74	0.4103 [#]
		Female	Normal	131.44 ± 2.83	131.21 ± 4.55	0.9550
			Over	136.32 ± 4.96	141.45 ± 13.15	0.7109
Cholesterol (mg/dL) ⁷⁾	Young	Male	Normal	193.57 ± 2.48	192.27 ± 3.37	0.7338
			Over	197.37 ± 2.43	204.79 ± 6.92	0.3461
		Female	Normal	187.00 ± 3.23	184.44 ± 2.98	0.2115
			Over	198.35 ± 2.12	194.29 ± 3.75	0.3698
	Old	Male	Normal	193.10 ± 1.46	194.55 ± 3.70	0.7262
			Over	195.05 ± 3.13	203.79 ± 5.56	0.1187
		Female	Normal	198.16 ± 1.09	203.78 ± 2.79	0.0646 [#]
			Over	201.55 ± 1.76	198.44 ± 4.78	0.5571
HDL (mg/dL) ⁸⁾	Young	Male	Normal	46.74 ± 0.57	47.41 ± 1.05	0.5468
			Over	44.75 ± 0.73	46.08 ± 1.06	0.2834
		Female	Normal	52.44 ± 0.59	51.22 ± 0.76	0.1107
			Over	50.03 ± 0.80	51.69 ± 1.32	0.1319
	Old	Male	Normal	47.37 ± 0.53	45.75 ± 1.14	0.1333
			Over	46.15 ± 0.65	47.21 ± 1.29	0.3657
		Female	Normal	52.18 ± 0.63	51.98 ± 0.98	0.8241
			Over	49.30 ± 0.67	47.11 ± 0.94	0.0112*

¹⁻⁸⁾ Refer to Table 6.

* The difference between the two levels of curry consumption became significant after adjusting for the covariates

[#] The difference between the two levels of curry consumption became insignificant after adjusting for the covariates

DISCUSSION

Curry is the major curcumin-containing food in the Korean diet [1]. Although, only a few individuals were included in the HC, their mean daily turmeric intake was about 0.26 g (Table 3). The demographic characteristics of the study subjects in the higher curry consumption groups resemble the curcumin consumers retrieved by the 24-hour recall data obtained from the KNHANES of years 2008-2012 [1]. The higher curry consumption groups more frequently included younger individuals and

non-apartment residents (Table 3), as previously reported [1]. Therefore, information on food consumption surveyed through 24-hour recall and FFQ is comparable when the curry consumption is considered.

Compared to the LC, more individuals in the MC reported being in better health; they had less diagnoses of hypertension, dyslipidemia, and diabetes. The MC group also included more individuals who had healthier behaviors, as more of them used dietary supplements, exercised, and were non-smokers (Table 5). Therefore, healthier individuals might consume curry more

often. In contrast, individuals in the MC more frequently consumed instant noodles that are considered an unhealthy food choice (Table 5). This may reflect the more frequent use of convenience foods by individuals in the MC which, correlates with the fact that curry is a convenient food. Alternatively, the health related behaviors observed simply represents the behavior of younger individuals as the mean age of the MC was lower compared to the LC. Since the two groups exhibited different health-related behaviors and diagnostic histories, it was critical to evaluate whether these factors confounded the blood measurements in this study.

All of the blood measurements were confounded by the BMI, which is a known correlate of the blood lipid and glucose levels [19,20]. The blood glucose, HbA_{1c}, and cholesterol levels were influenced by age as well (Table 7). The blood TG and HDL levels were also affected by sex (Table 7), as reported previously [21,22]. Interestingly, diabetes was a confounding factor for TG, total cholesterol, as well as HDL cholesterol (Table 7). Hyperglycaemia is associated with adverse lipid profiles [23], suggesting that the dysregulation of the glucose levels may affect lipid metabolism. Similarly, hypertriglyceridemia but not hypercholesterolemia influenced the blood glucose levels (Table 7).

Curry consumption may be beneficial in overweight (BMI \geq 25) individuals who have high levels of blood glucose or TG. The difference in the glucose and TG levels according to the curry consumption became significant in the old-male-overweight and young-female-overweight subgroups, respectively, after adjusting for the identified confounding variables (Table 8). Blood glucose and TG levels are associated with insulin sensitivity [24]. Therefore, the lower levels of both the blood glucose and the TG in the MC may indicate that individuals with higher consumption of curry may have better insulin sensitivity than individuals in the LC. Unfortunately insulin levels were not assessed in the KNHANES conducted in 2012 and 2013. Unexpectedly, the blood HDL level adjusted for the confounding variable was significantly lower in old-female-overweight individuals in the MC compared to the LC. However, in this subgroup, the difference of the HDL level is less than 3 mg/dL, and the tight standard deviation within this subgroup may contribute to a significant difference.

Previous preclinical studies demonstrated that curcumin exhibits a hypolipidemic effect and enhances insulin sensitivity. Dietary curcumin (less than 0.05%) improves insulin resistance and lowers total cholesterol, free fatty acid, and TG levels in the blood of high-fat fed rodents [25]. Feeding curcumin also reduces the blood glucose levels as well as the lipid levels and enhances insulin sensitivity in diabetic mice [26] and diabetes-induced rats [27]. Therefore, studies indicate that dietary curcumin modulates the lipid and carbohydrate metabolism, subsequently lowering the blood levels of glucose, total cholesterol, and TG. However, this effect only occurs in diabetic mice or high-fat fed animals, where the blood levels of glucose and lipids are elevated. Dietary curcumin does not affect the blood glucose or the glucose-regulating enzyme activities in non-diabetic mice [26,27]. In addition, the cholesterol lowering effect is only observed in old rats with high cholesterol levels [5]. This may be why the effect of curry consumption only occurred in overweight individuals, who had higher levels of

glucose and TG, in this study. However, the total cholesterol level was not altered by the curry consumption in any subgroups, and moderate curry consumption did not affect the blood glucose and lipid levels in young-male-overweight or old-female overweight individuals, who also had high blood levels of glucose and TG (Table 8). This may be because moderate curry consumption might not be enough to reduce the glucose and lipids in all individuals.

Glycated hemoglobin (HbA_{1c}) is used as a marker that estimates average blood glucose levels over a period of months [28]. Moderate curry consumption decreases glucose, but not HbA_{1c} in old-male-overweight individuals. The HbA_{1c} levels were in the normal range (4-5.9%) in all subgroups (Table 8), which might cause no significant difference according to the curry consumption. In contrast, moderate curry consumption significantly decreased the HbA_{1c}, but not the glucose levels in young-male-normal individuals. However, in this subgroup, the difference of the HbA_{1c} is only 0.1% and, more than likely, the very tight standard deviation within this subgroup may be attributed to a significant difference.

Sources of turmeric other than curry were not included in the FFQ conducted by KNHANES in 2012 and 2013. However, the primary source of turmeric in the Korean diet is curry, and other sources of turmeric intake are rare [1]. In addition, other types of curry consumption could not be estimated in this study. However, the usage of curry other than "curry rice" was very rare in spite of its use in batter for frying or sauce [1]. Although curry was not classified by turmeric content in the FFQ, this may not cause misclassification of the curry consumption groups since the turmeric content is similar between products made by different manufacturers, and turmeric enhanced products are less common in the market. The accuracy of the self-reported levels of curry consumption could not be determined in this study; however, curry is usually consumed as a main dish, and a recall bias might be insignificant. The MC might include more individuals with higher total energy consumption. However, this likelihood may not be very high, since difference tests were performed within the defined BMI categories and the self-reported portion size for curry was mostly one serving (Table 1).

A major obstacle in this study was that few individuals consumed curry with high frequency. In addition, the effect of curry consumption could not be evaluated in individuals with a prior history of diabetes or dyslipidemia, due to the limited number of individuals with diabetes or dyslipidemia in each age-sex-BMI subgroup. Nevertheless, it is noteworthy that a moderate level of curry consumption (2 to 4 times per month) is related to lower blood levels of glucose and TG in overweight individuals who have high blood lipid and glucose levels, after estimates were adjusted for the confounding factors. Curry is a major curcumin-containing food in most countries other than South Asia, where turmeric consumption is not high. Although it is possible that healthier individuals may consume curry more often, these results are suggestive of the potential health benefits stemming from the consumption of curcumin through an ordinary diet; this should be confirmed in a longitudinal study in the future.

REFERENCES

1. Kwon Y. Estimation of curcumin intake in Korea based on the Korea National Health and Nutrition Examination Survey (2008-2012). *Nutr Res Pract* 2014;8:589-94.
2. Bagad AS, Joseph JA, Bhaskaran N, Agarwal A. Comparative evaluation of anti-inflammatory activity of curcuminoids, turmerones, and aqueous extract of *Curcuma longa*. *Adv Pharmacol Sci* 2013;2013:805756.
3. Guimarães MR, Leite FR, Spolidorio LC, Kirkwood KL, Rossa C Jr. Curcumin abrogates LPS-induced pro-inflammatory cytokines in RAW 264.7 macrophages. Evidence for novel mechanisms involving SOCS-1, -3 and p38 MAPK. *Arch Oral Biol* 2013;58:1309-17.
4. Hu GX, Lin H, Lian QQ, Zhou SH, Guo J, Zhou HY, Chu Y, Ge RS. Curcumin as a potent and selective inhibitor of 11 β -hydroxysteroid dehydrogenase 1: improving lipid profiles in high-fat-diet-treated rats. *PLoS One* 2013;8:e49976.
5. Kwon Y, Malik M, Magnuson BA. Inhibition of colonic aberrant crypt foci by curcumin in rats is affected by age. *Nutr Cancer* 2004;48:37-43.
6. Peschel D, Koerting R, Nass N. Curcumin induces changes in expression of genes involved in cholesterol homeostasis. *J Nutr Biochem* 2007;18:113-9.
7. Zingg JM, Hasan ST, Meydani M. Molecular mechanisms of hypolipidemic effects of curcumin. *Biofactors* 2013;39:101-21.
8. Gupta SC, Patchva S, Aggarwal BB. Therapeutic roles of curcumin: lessons learned from clinical trials. *AAPS J* 2013;15:195-218.
9. Hatcher H, Planalp R, Cho J, Torti FM, Torti SV. Curcumin: from ancient medicine to current clinical trials. *Cell Mol Life Sci* 2008;65:1631-52.
10. Ng TP, Chiam PC, Lee T, Chua HC, Lim L, Kua EH. Curry consumption and cognitive function in the elderly. *Am J Epidemiol* 2006;164:898-906.
11. Ng TP, Niti M, Yap KB, Tan WC. Curcumins-rich curry diet and pulmonary function in Asian older adults. *PLoS One* 2012;7:e51753.
12. Sugawara J, Akazawa N, Miyaki A, Choi Y, Tanabe Y, Imai T, Maeda S. Effect of endurance exercise training and curcumin intake on central arterial hemodynamics in postmenopausal women: pilot study. *Am J Hypertens* 2012;25:651-6.
13. Carroll RE, Benya RV, Turgeon DK, Vareed S, Neuman M, Rodriguez L, Kakarala M, Carpenter PM, McLaren C, Meyskens FL Jr, Brenner DE. Phase IIa clinical trial of curcumin for the prevention of colorectal neoplasia. *Cancer Prev Res (Phila)* 2011;4:354-64.
14. Dhillon N, Aggarwal BB, Newman RA, Wolff RA, Kunnumakkara AB, Abbruzzese JL, Ng CS, Badmaev V, Kurzrock R. Phase II trial of curcumin in patients with advanced pancreatic cancer. *Clin Cancer Res* 2008;14:4491-9.
15. Chandrasekaran CV, Sundarajan K, Edwin JR, Gururaja GM, Mundkinajeddu D, Agarwal A. Immune-stimulatory and anti-inflammatory activities of *Curcuma longa* extract and its polysaccharide fraction. *Pharmacognosy Res* 2013;5:71-9.
16. Lahiff C, Moss AC. Curcumin for clinical and endoscopic remission in ulcerative colitis. *Inflamm Bowel Dis* 2011;17:E66.
17. Ammon HP, Wahl MA. Pharmacology of *Curcuma longa*. *Planta Med* 1991;57:1-7.
18. Kweon S, Kim Y, Jang MJ, Kim Y, Kim K, Choi S, Chun C, Khang YH, Oh K. Data resource profile: the Korea National Health and Nutrition Examination Survey (KNHANES). *Int J Epidemiol* 2014;43:69-77.
19. Kostis JB, McCrone K, Moreyra AE, Hosler M, Cosgrove N, Kuo PT. The effect of age, blood pressure and gender on the incidence of premature ventricular contractions. *Angiology* 1982;33:464-73.
20. Hu FB, Wang B, Chen C, Jin Y, Yang J, Stampfer MJ, Xu X. Body mass index and cardiovascular risk factors in a rural Chinese population. *Am J Epidemiol* 2000;151:88-97.
21. Magkos F, Mittendorfer B. Gender differences in lipid metabolism and the effect of obesity. *Obstet Gynecol Clin North Am* 2009;36:245-65.
22. Faerch K, Borch-Johnsen K, Vaag A, Jørgensen T, Witte DR. Sex differences in glucose levels: a consequence of physiology or methodological convenience? The Inter99 study. *Diabetologia* 2010;53:858-65.
23. Zhang L, Qiao Q, Tuomilehto J, Hammar N, Janus ED, Söderberg S, Mohan V, Ramachandran A, Dong YH, Lam TH, Pang ZC. Blood lipid levels in relation to glucose status in seven populations of Asian origin without a prior history of diabetes: the DECODA study. *Diabetes Metab Res Rev* 2009;25:549-57.
24. Boden G, Laakso M. Lipids and glucose in type 2 diabetes: what is the cause and effect? *Diabetes Care* 2004;27:2253-9.
25. Jang EM, Choi MS, Jung UJ, Kim MJ, Kim HJ, Jeon SM, Shin SK, Seong CN, Lee MK. Beneficial effects of curcumin on hyperlipidemia and insulin resistance in high-fat-fed hamsters. *Metabolism* 2008;57:1576-83.
26. Seo KI, Choi MS, Jung UJ, Kim HJ, Yeo J, Jeon SM, Lee MK. Effect of curcumin supplementation on blood glucose, plasma insulin, and glucose homeostasis related enzyme activities in diabetic db/db mice. *Mol Nutr Food Res* 2008;52:995-1004.
27. Na LX, Zhang YL, Li Y, Liu LY, Li R, Kong T, Sun CH. Curcumin improves insulin resistance in skeletal muscle of rats. *Nutr Metab Cardiovasc Dis* 2011;21:526-33.
28. Donahue RP, Bean JA, Donahue RD, Goldberg RB, Prineas RJ. Does insulin resistance unite the separate components of the insulin resistance syndrome? Evidence from the Miami Community Health Study. *Arterioscler Thromb Vasc Biol* 1997;17:2413-7.