

POSTER PRESENTATION

Open Access

A central role for Galectin 3 during renal epithelial cell morphogenesis after nephrectomy

F Poirier^{1*}, A Viau², J Magescas¹, M Burtin², F Terzi², D Delacour¹

From First International Cilia in Development and Disease Scientific Conference (2012)
London, UK. 16-18 May 2012

Galectin-3, a member of the multigene family of beta-galactoside binding lectins, is mainly expressed in epithelial cells. The distribution of galectin-3 depends on cell type and cell differentiation. We previously showed that galectin-3 is transiently associated with the centrosome, and more specifically with the basal body at the basis of the primary cilium of MDCK cells. In the present study, we use *galectin-3* null (*Gal3*^{-/-}) mutant mice to investigate the functional consequences of our observations in the context of kidney regeneration after subtotal nephrectomy. Three months after surgery, renal functions were more severely affected in *Gal3*^{-/-} mutant than in wt animals. Upon sacrifice, we found that *Gal3*^{-/-} mutant kidneys were bigger and heavier, and this hypertrophy was associated with large tubular dilatations. Interestingly, in wt mice, galectin 3 intracellular distribution was altered in response to injury, displaying a distinct centrosomal localization in epithelial cells lining the collecting ducts close to the wound, whereas in cells away from the wound galectin-3 remained uniformly cytosolic. The impact of nephron reduction on primary cilium growth was examined by 3D reconstruction of wt and *Gal3*^{-/-} renal tubules stained for acetylated alpha-tubulin. This analysis revealed that mutant primary cilia are irregular, bent, twisted and longer than wt primary cilia, indicating that Galectin-3 is essential for primary cilium growth. This suggests that the failure in kidney regeneration observed in *Gal3*^{-/-} mutant mice may be due to defects in primary cilium biogenesis.

Author details

¹Institut Jacques Monod CNRS Paris Diderot, France. ²INSERM U845, France.

Published: 16 November 2012

* Correspondence: poirier@ijm.univ-paris-diderot.fr

¹Institut Jacques Monod CNRS Paris Diderot, France

Full list of author information is available at the end of the article

doi:10.1186/2046-2530-1-S1-P76

Cite this article as: Poirier et al.: A central role for Galectin 3 during renal epithelial cell morphogenesis after nephrectomy. *Cilia* 2012 **1** (Suppl 1):P76.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

