

Supplementary Online Content

Kim YG, Lee KN, Han KD, et al. Association of depression with atrial fibrillation in South Korean adults. *JAMA Netw Open*. 2022;5(1):e2141772.
doi:10.1001/jamanetworkopen.2021.41772

eTable. Diagnostic codes.

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Diagnostic Codes.

	ICD-10 codes
Depression	F32 (all sub-codes), F33 (all sub-codes)
Atrial fibrillation	I48 (all sub-codes)
Mitral stenosis	I05.0, I05.2, I34.2
Valve surgery	O1781, O1791, O1782, O1792, O1783, O1793, O1799, O1797
Heart failure	I50 (all sub-codes)
Atrial fibrillation	I48 (all sub-codes)
Type 2 diabetes	E11 – E14 (all sub-codes)
Hypertension	I10 – I13, I15 (all sub-codes)
Dyslipidemia	I78 (all sub-codes)
Chronic kidney disease	Based on creatinine checked during national health check-up
Schizophrenia	F20 (all sub-codes)
Bipolar affective disorder	F31 (all sub-codes)
Dementia	G30 (all sub-codes), F02 (all sub-codes), F03 (all sub-codes)
Hypo- or hyperthyroidism	E03 (all sub-codes), E05 (all sub-codes)