

# Rain Amplification of Persistent Organic Pollutants

Gemma Casas, Alicia Martinez-Varela, Maria Vila-Costa, Begoña Jiménez, and Jordi Dachs\*



Cite This: *Environ. Sci. Technol.* 2021, 55, 12961–12972



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

**ABSTRACT:** Scavenging of gas- and aerosol-phase organic pollutants by rain is an efficient wet deposition mechanism of organic pollutants. However, whereas snow has been identified as a key amplification mechanism of fugacities in cold environments, rain has received less attention in terms of amplification of organic pollutants. In this work, we provide new measurements of concentrations of perfluoroalkyl substances (PFAS), organophosphate esters (OPEs), and polycyclic aromatic hydrocarbons (PAHs) in rain from Antarctica, showing high scavenging ratios. Furthermore, a meta-analysis of previously published concentrations in air and rain was performed, with 46 works covering different climatic regions and a wide range of chemical classes, including PFAS, OPEs, PAHs, polychlorinated biphenyls and organochlorine compounds, polybromodiphenyl ethers, and dioxins. The rain–aerosol ( $K_{RP}$ ) and rain–gas ( $K_{RG}$ ) partition constants averaged  $10^{5.5}$  and  $10^{4.1}$ , respectively, but showed large variability. The high field-derived values of  $K_{RG}$  are consistent with adsorption onto the raindrops as a scavenging mechanism, in addition to gas–water absorption. The amplification of fugacities by rain deposition was up to 3 orders of magnitude for all chemical classes and was comparable to that due to snow. The amplification of concentrations and fugacities by rain underscores its relevance, explaining the occurrence of organic pollutants in environments across different climatic regions.

**KEYWORDS:** wet deposition, snow, scavenging, amplification, Antarctica, PFASs, OPEs, PAHs, PCBs, PBDEs



## INTRODUCTION

Persistent organic pollutants (POPs) have the potential for long-range atmospheric transport (LRAT) from source to remote regions due to their persistence, semivolatility, or transference to the atmosphere with sea-spray aerosols.<sup>1–3</sup> The study of LRAT and atmospheric deposition has been central in previous assessments of the occurrence of POPs at regional and global scales. Some of these mechanisms were previously described, such as cold trapping,<sup>4</sup> enhanced deposition due to the biological pump,<sup>5,6</sup> degradation pump enhanced deposition,<sup>7,8</sup> temperature and biological pump-driven grasshopping over the oceans,<sup>4,9</sup> and retardation of grasshopping due to sorption to soil and vegetation organic matter.<sup>10,11</sup> Many of these transport mechanisms are fugacity-driven diffusive fluxes between the air and the receiving surface (water, soils, and vegetation). Conversely, wet deposition by rain and snow is independent of the fugacity gradient between the air and surface but is dependent on the capacity of rain and snow to scavenge atmospheric pollutants.<sup>12–14</sup> These deposition processes can lead to higher concentrations of organic pollutants in water and soils than those derived from air–surface partitioning only, a process named “amplification.” However, the amplification of organic pollutants has received little attention. Previous works focused on the role of snow deposition amplifying the concentrations of POPs in soils and

seawater.<sup>15–19</sup> Comparatively, the role of rain deposition has received less attention.

Wet deposition by rain or snow<sup>12</sup> scavenges POPs found in the gas and aerosol phases very efficiently and can amplify concentrations of POPs in other environmental compartments, such as in seawater<sup>18–21</sup> or soils.<sup>22–25</sup> This process has been proven to be especially effective for snow scavenging and is one of the main entries of POPs in cold regions, such as high-mountain and polar regions.<sup>18,26–28</sup> Partition toward the snowflakes is favored at low temperatures. This together with the high specific surface area of snow, makes this deposition process crucial to understand the occurrence of a large variety of POPs in polar regions.<sup>17–19,26,27,29</sup> A meta-analysis of the snow–air partition constants ( $K_{SA}$ ), estimated as the ratio of POP concentrations in snow and air, from previously reported simultaneous field measurements, showed that snow amplification was relevant for diverse families of POPs, independent of their volatility.<sup>18</sup> The same work showed that seawater–air fugacity ratios of polychlorinated biphenyls (PCBs) were

**Received:** May 20, 2021

**Revised:** August 2, 2021

**Accepted:** August 17, 2021

**Published:** September 23, 2021



highly correlated with the product of  $K_{SA}$  and the dimensionless Henry's law constant ( $H'$ ), a measure of snow amplification of fugacity. Therefore, coastal seawater mirrored the PCBs in snow due to snowmelt releasing POPs. This and other observations of the occurrence of POPs provide the field evidence that pollution in the environment is underpredicted when considering air-surface diffusive partitioning only.

Rain also scavenges atmospheric pollutants found in the gas and aerosol phases, but the role of rain deposition, occurring under various climatic regimes, remains unclear, especially in terms of amplification of concentrations and fugacities. Snow and rain deposition have been compared previously, suggesting that snow is significantly more efficient than rain when collecting POPs from the atmosphere.<sup>26</sup> Nowadays, the occurrence of legacy and emerging POPs have been reported in rain from diverse climatic regions, with what has become a remarkable body of field evidence of the role of rain as a deposition process.<sup>30–74</sup>

The wet deposition flux ( $F_{WD}$ ,  $\text{pg m}^{-2} \text{d}^{-1}$ ) is given by

$$F_{WD} = p_0 C_{\text{rain}} \quad (1)$$

where  $p_0$  ( $\text{m d}^{-1}$ ) is the precipitation depth per day and  $C_{\text{rain}}$  ( $\text{pg m}^{-3}$ ) is the POP concentration in rain. It is useful to refer to the wet deposition flux for the raining period and for the atmospheric concentrations, which are often measured instead of concentrations in the rain. Then, the estimation of the wet deposition flux is given by

$$F_{WD\_rain} = K_{RA}(C_{\text{air},g} + C_{\text{air},p}) \frac{p_0}{f} \quad (2)$$

where  $C_{\text{air},p}$  is the concentration in the particulate phase of air,  $C_{\text{air},g}$  is the concentration in the gas phase, and  $f$  is the fractional occurrence of rain. The rain–air partition constant ( $K_{RA}$ ), also called the water/air scavenging ratio or washout ratio, is given by<sup>33</sup>

$$K_{RA} = \frac{C_{\text{rain}}}{C_{\text{air},p} + C_{\text{air},g}} = K_{RP} \theta + K_{RG}(1 - \theta) \quad (3)$$

where  $K_{RG}$  and  $K_{RP}$  are the rain–air partition constants for the gas phase and particulate phase, respectively.  $\theta$  (dimensionless) is the fraction of aerosol-bound POPs to the total atmospheric POP concentration ( $C_{\text{air},p}/C_{\text{air},p} + C_{\text{air},g}$ ). These dimensionless partition constants can be estimated by

$$K_{RP} = \frac{C_{\text{rain}}}{C_{\text{air},p}} \quad (4)$$

$$K_{RG} = \frac{C_{\text{rain}}}{C_{\text{air},g}} \quad (5)$$

$K_{RA}$  is thus the scavenging ratio including both the gas- and particulate-phase concentrations in air, while  $K_{RP}$  and  $K_{RG}$  only consider the scavenging of aerosol- or gas-phase POPs, respectively.

The values of  $K_{RP}$  depend, in a complex manner, on meteorology, aerosol characteristics, and chemical properties and have generally been determined empirically. Field derived values are highly variable with a mean value<sup>13</sup> of  $2 \times 10^5$ . Conversely,  $K_{RG}$  depends on the raindrop-air diffusive partitioning and the POP adsorption on the raindrop surface from the gas phase. Thus,

$$K_{RG} = K_{RG,dissolved} + K_{RG,adsorbed} \quad (6)$$

Therefore,  $K_{RG}$  is the sum of the absorption and adsorption terms ( $K_{RG,dissolved}$  and  $K_{RG,adsorbed}$ ).  $K_{RG,dissolved}$  is given by the inverse of the dimensionless Henry's law constant ( $1/H'$ ). On the other hand,  $K_{RG,adsorbed}$  depends on the water interface-air partition constant and the size distribution of rain drops.<sup>13,75</sup> The second term in eq 6 has been suggested to be important for polycyclic aromatic hydrocarbons (PAHs), dioxins, and furans (PCDD/Fs) but not for PCBs.<sup>13,75</sup> When the adsorption to raindrops is important, then it is possible that there is an amplification of gas-phase POPs in rain.


The objectives of this work are (i) to report a meta-analysis of the rain–air partitioning of POPs previously reported in the literature, with a contribution of a new data set of perfluoroalkyl substances (PFAS), organophosphate ester (OPEs), and PAHs in the Antarctic rain, (ii) to assess the potential for rain amplification for different legacies and emerging POPs, and (iii) to compare rain and snow amplification of POPs.

## MATERIALS AND METHODS

**Site Description and Sampling.** The samples for this study were collected during two sampling campaigns. The first campaign was carried out in Deception Island ( $62^\circ 58' \text{S}$   $60^\circ 39' \text{W}$ ) during the Antarctic summer 2016–2017, while the other one was in Livingston Island ( $62^\circ 36' \text{S}$   $60^\circ 30' \text{O}$ ) during the Antarctic summer 2017–2018. These two islands are located in the South Shetland Archipelago (Figure S1) in the Antarctic Peninsula. In this area, periods with temperatures above freezing during summer allow rain to fall.<sup>76</sup> The samples for PFAS analysis were collected in both islands; however, the samples for OPEs and PAHs were collected only in Livingston Island.

The rain samples were collected with a stainless steel tray, from which 2 L was poured into Teflon bottles for the analysis of OPEs and PAHs ( $n = 10$ ), and 2 L was poured into polypropylene bottles for the analysis of PFAS ( $n = 21$ ). After 30–60 min of sample collection, the samples were filtered through precombusted GF/F glass fiber filters (47 mm diameter, Whatman 0.7  $\mu\text{m}$  mesh size) before passing through their specific cartridges at the Antarctic research station's laboratory. The PFAS extraction was based on a solid-phase extraction (SPE) procedure using Oasis WAX cartridges (6  $\text{cm}^3$ , 150 mg; Waters).<sup>77</sup> On the other hand, OPEs and PAHs were extracted using an SPE procedure with Oasis hydrophilic–lipophilic balance cartridges (6  $\text{cm}^3$ , 200 mg; Waters) by following an established methodology<sup>78,79</sup> with some modifications. Samples were spiked with recovery standards before the SPE (Table S1). Aerosol samples were collected (Figure S1) using a high volume air sampler (MCV, Collbató, Spain) operating at  $40 \text{ m}^3 \text{h}^{-1}$ . The air was drawn through a precombusted and preweighed quartz microfiber filter (QM-A; Whatman, 8–10 inches) to collect aerosol-bound compounds (total suspended particle). Twenty aerosol samples were analyzed for PFAS and 6 aerosol samples were analyzed for OPEs and PAHs. All the samples/cartridges were stored at  $-20^\circ \text{C}$ , after the sampling, until analysis in an ultraclean laboratory in Barcelona.


The procedures followed for the extraction, identification, and quantification of PFAS, OPEs, and PAHs are described in Annex S1 in Supporting Information. Quality assurance and quality control are reported in Annex S2 in Supporting Information. Recoveries and limits of detection are summarized in Tables S2 and S3 in Supporting Information.



**Figure 1.** Meta-analysis of rain–air particulate partition constants ( $K_{RP}$ ) for various families of organic pollutants. The results shown are the mean and the standard deviation of  $\log K_{RP}$ .

**Criteria for Meta-Analysis of Rain–Air Partition Constants.** We reviewed the previous reports of organic pollutants in rain and air (particulate and gas phase), and a total of 45 publications were found and used for this study.<sup>30–74</sup> There are additional reports of field concentrations of organic pollutants in rain,<sup>80–101</sup> but they lack concurrently measured atmospheric concentrations, or data were not given in the manuscript, and thus rain–air partition constants could not be estimated. Table S4 summarizes the data set contained in the 45 publications used for the partition constants. With the new data reported for PFAS, OPEs, and PAHs in this study, a total of 46 data sets were used in this meta-analysis. The meta-analysis was made for the three rain–air partition constants (eqs 3–5). In some of the previous studies, one, two, or three of the coefficients,  $K_{RP}$ ,  $K_{RG}$ , and  $K_{RA}$  were already

provided. For the other works reporting concentrations, we calculated the respective partition constants by following eqs 3–5. We were not able to perform the meta-analysis with all data sets for the three different coefficients as some works only reported concentrations for the particulate, or gas phase, or total atmospheric concentration. In the case of PCBs and hexachlorocyclohexane (HCH), if the authors provided only the concentrations in the gas phase, both  $K_{RG}$  and  $K_{RA}$  were estimated, as it is well known that these POPs are mostly found in the gas phase (low  $\theta$  value in eq 3). These data were used to derive Figures 1, 2, 3, 4 and S5–S9 and are reported in Tables S4–S11. The results shown in these figures are the mean and the standard deviation of the partition constants for each compound and each data set. The standard deviation is not given when the original work provided only one value of the



**Figure 2.** Meta-analysis of rain–gas partition constants ( $K_{RG}$ ) for various families of organic pollutants. The results shown are the mean and the standard deviation of  $\log K_{RG}$ .


coefficients for each compound or reported only the mean (details in Table S4). We focused on rain–air partition coefficients as reported directly from chemical measurements. We did not consider estimates of these partition constants from models, even if these were partly based on measurements. For the estimation of  $K_{RG}H'$  (amplification potential), we used  $H'$  values at 298.15 K (Table S8). All the calculations, statistics, and plots for the meta-analysis were performed using R Studio 1.4 (Figures 1, 2, 3, 4, 5, S1–S9, Tables S12).

## RESULTS AND DISCUSSION

**Concentrations of PFAS, OPEs, and PAHs in Rain and Aerosols from Maritime Antarctica.** The concentrations of the individual PFAS, OPEs, and PAHs measured in the rain and aerosol samples from Deception and Livingston Islands

(the Antarctic Peninsula) are summarized in Tables S9–S11 (Supporting Information) and shown in Figures S2–S4. The average and range concentrations of  $\Sigma$ PFASs in rain and aerosol samples from Deception Island were 3,600 (660–7,600)  $\text{pg L}^{-1}$  and 0.13 (0.056–0.32)  $\text{pg m}^{-3}$ , respectively. The average and range concentrations of  $\Sigma$ PFASs in rain and aerosol samples from Livingston Island were 3,600 (400–8,400)  $\text{pg L}^{-1}$  and 0.19 (0.0067–0.71)  $\text{pg m}^{-3}$ , respectively. To the best of our knowledge, this is the first report of POP concentrations in rain from Antarctica. These concentrations in rain were 1 order of magnitude lower than those reported in Northern Germany.<sup>65</sup> On the other hand, these PFAS concentrations in rain were similar to those measured in snow from Livingston Island.<sup>19</sup> PFAS concentrations in aerosol





**Figure 3.** Meta-analysis of rain–air (particulate + gas phase) partition constants ( $K_{RA}$ ) for various families of organic pollutants. The results shown are the mean and the standard deviation of  $\log K_{RA}$ .

samples were comparable to those reported previously at Livingston Island.<sup>2</sup>

The average and range concentrations of  $\Sigma$ OPEs in rain and aerosol samples from Livingston Island were 35,800 (9730–93,600)  $\text{pg L}^{-1}$  and 41.7 (13.3–28.3)  $\text{pg m}^{-3}$ , respectively. The concentrations reported here for rain were 1 order of magnitude lower than those reported in Germany.<sup>87,102</sup> The aerosol-phase concentrations of OPEs were comparable with those reported in the Western Antarctic Peninsula for 2014–2018,<sup>103</sup> which ranged from 5.75 to 238  $\text{pg m}^{-3}$ .

The average concentrations of  $\Sigma$ PAHs in rain samples from Livingston Island averaged 9.05 (2.36–17.8)  $\text{ng L}^{-1}$  and concentrations of  $\Sigma$ PAHs in aerosol samples averaged 0.02

(0.005–0.03)  $\text{ng m}^{-3}$ . These aerosol-phase concentrations from Livingston Island were comparable to the  $\Sigma$ PAH concentrations previously reported from the Antarctic and Southern Ocean atmosphere,<sup>104</sup> where  $\Sigma$ PAH average concentrations from Livingston Island ranged from 0.03 to 0.09  $\text{ng m}^{-3}$ . The  $\Sigma$ PAH concentrations in rain for this study were significantly lower than those reported in rain during the Malaspina 2010 circumnavigation in the tropical and subtropical oceans.<sup>71</sup>

**Rain Scavenging of Aerosol-Bound POPs.** Particle scavenging by rain is characterized by  $K_{RP}$ . The efficiency of the washout of aerosols by rain depends on a number of factors, such as the aerosol and raindrop size distributions,




**Figure 4.** Predicted rain amplification of the fugacity ratio between rain and air given by the product of the rain–air (gas phase) partition constant and the dimensionless Henry's law constant ( $f_w/f_G = K_{RG}H'$ ) for various families of organic pollutants. The results shown are the mean and the standard deviation of  $\log K_{RG}H'$ .

among other factors.<sup>13,75</sup> There are a number of reported measurements of  $K_{RP}$  for PCBs and PAHs, some of which have been used in modeling exercises for wet deposition.<sup>13,36,44,45,50,58,62,66,105</sup> We compiled all the previous reports of simultaneous occurrence of organic pollutants in rain and aerosols, with 24 studies reporting concentrations of PCBs, PCDD/Fs, PAHs, organochlorine pesticides (OCPs), PBDEs, and PFAS, together with the new data set from this work for PAHs, OPEs, and PFAS from Antarctica. Figure 1 shows the comparison of  $K_{RP}$  from this meta-analysis. The compound-specific average  $\log K_{RP}$  value ranged between 2.6 and 11.5; the mean value was 5.5.

The  $\log K_{RP}$  value of PAHs ranged between 2.6 and 11.5, representing the highest variability. Among these, the values

derived from studies by Ligocki et al. (1985)<sup>33</sup> showed the most different values of  $K_{RP}$  compared with all the other studies, especially for the high molecular weight (MW) PAHs. Naphthalene and their methylated compounds showed the highest  $\log K_{RP}$  average value, ranging from 4.9 to 11.5. With the exception of PAHs, for which there was a high variability of  $K_{RP}$ , the other pollutant families showed similar  $K_{RP}$  values among the different studies. The  $\log K_{RP}$  values of PCBs and OCPs presented similar ranges between studies, within the range 4.0–7.0 and 2.7–6.6, respectively. The  $\log K_{RP}$  value of PCDD/Fs ranged between 4.0 and 5.0.<sup>37,38</sup>

As far as we know, the field-derived  $K_{RP}$  value of OPEs is reported here for the first time, with the  $\log K_{RP}$  value ranging from 4.1 to 7.3. The  $\log K_{RP}$  value of PBDEs ranged between



**Figure 5.** Comparison of amplification of organic pollutants by rain and snow. The results show the mean of  $K_{RG}/K_{SA}$  and its error estimated using the uncertainty propagation approach (Annex S3).

4.01 and 7.0.<sup>49,50,59,60,62,63,68</sup> The  $K_{RP}$  value ranged between 3.6 and 8.8 for PFAS.<sup>57,65</sup> We could only compare  $K_{RP}$  for various PFAS from two different studies that measured concentrations simultaneously in rain and aerosols. In addition, Barton et al. (2007)<sup>57</sup> reported  $K_{RP}$  values for PFOA only. The  $K_{RP}$  values from previously reported concentrations<sup>65,106</sup> were comparable with those measured here (Figure 1).

The aerosol type and the physical and chemical properties of the compound could influence the values of the scavenging ratios. OPEs and PFAS showed the highest average values for  $\log K_{RP}$ , together with naphthalene and some methylnaphthalenes. Overall, the  $K_{RP}$  values of the different chemicals showed high variability. Nevertheless, these were significantly correlated with  $K_{OA}$  and  $H'$  ( $K_{AW}$ ), but correlations explained a

small percentage of the variability ( $r = 0.39$  or  $r = 0.2$ , respectively) (Figure S9 and Table S12). In addition, we compared  $K_{RP}$  for aerosols having different origins, by classifying the field studies between those performed in urban and continental areas and coastal and open oceans (Figure S5). A Tukey HSD test was carried out for performing pairwise comparison between the means of  $K_{RP}$  for different aerosol types, which showed significant differences between continental/urban areas and coastal/open ocean aerosols. Conversely, there were no significant differences between continental and urban areas and between coastal and open oceans. For modeling purposes, and for chemicals for which field-derived  $K_{RP}$  values (Figure 1) are not available, this meta-

analysis shows that a value for  $\log K_{RP}$  between 5 and 6 would be reasonable (Figure 1).

**Rain Scavenging of Gas-Phase POPs.** Scavenging or washout of gas-phase organic pollutants by rain is characterized by  $K_{RG}$ . There are also a number of reported  $K_{RG}$  field measurements for PCBs, OCPs, PAHs, PBDEs, and PCDD/Fs (Figure 2). Conversely, emerging pollutants have barely been measured concurrently in the gas phase and rain, with only one study reporting these for neutral PFAS (MeFOSE, EtFOSE, and MeFBSE).<sup>65,106</sup> Figure 2 shows the meta-analysis of  $K_{RG}$  with data from 35 studies. The compound-specific average  $\log K_{RG}$  value ranged between 1.1 and 9.6; the mean value was 4.1.

PAHs showed a high variability of  $\log K_{RG}$  (ranging between 1.1 and 7.3), while PCBs and OCPs showed similar  $K_{RG}$  values among the different studies. In the case of PAHs, the  $K_{RG}$  value derived from studies by Poster and Baker (1996)<sup>40</sup> presented the lowest values in comparison with the rest of the studies. Generally,  $K_{RG}$  increases as the number of aromatic rings increases, with the exception of naphthalene and alkyl-naphthalene (3.4–9.6), which have similar  $K_{RG}$  to high molecular weight (MW) PAHs, such as indeno[1,2,3-*cd*]-pyrene, dibenzo[*a,h*]anthracene, and benzo[*g,h,i*]perylene, ranging from 3.8 to 7.9. There is the possibility that there is some redissolution from aerosols to the rainwater dissolved phase, which would cause a sampling artifact explaining the high  $K_{RG}$  values observed for some chemicals, but this would be not consistent with the strong association of high MW PAHs with aerosol soot carbon. PCBs and OCPs presented similar compound-specific  $\log K_{RG}$  values among the different studies, ranging between 1.6–7.1 and 1.4–6.2, respectively. The  $\log K_{RG}$  value of PCDD/Fs ranged between 3.8 and 6.5. The  $\log K_{RG}$  values of PBDEs and neutral PFAS ranged from 3.2 to 8.1 and 5.3 to 5.4, respectively.

$K_{RG}$  depends on the raindrop-air diffusive partitioning and the POP adsorption on the raindrop surface from the gas phase.  $K_{RG,dissolved}$  equals the inverse of  $H'$ , and thus, knowing the field-derived  $K_{RG}$ , we could estimate  $K_{RG,adsorbed}$  (eq 6). Figure S6 shows that the  $K_{RG,adsorbed}$  value ranged between 1.1 and 9.6, with a mean of 4.1. Only for  $K_{RG}$  for PAHs reported by Poster and Baker,  $K_{RG,adsorbed}$  was negligible. For all other data sets and chemicals, adsorption on the raindrop is predicted to be not only important but also to dominate as a scavenging mechanism of gas-phase POPs from the atmosphere. This suggests that the common modeling practice of estimating  $K_{RG}$  as  $1/H'$  induces an underestimation by several orders of magnitude of the importance of rain deposition of POPs. In fact,  $K_{RG}$  showed a weak correlation with  $H'$  (spearman  $r = -0.156$ ,  $n = 498$ ,  $p < 0.001$ ) (Figure S8 and Table S12). For modeling  $K_{RG}$  of chemicals other than those shown in Figure 2, a mean value of  $10^{4.5}$  can be used but with high uncertainty.

**Amplification of POPs by Wet Deposition.** The overall importance of rain scavenging of both gas- and aerosol-phase POPs is characterized by  $K_{RA}$  (Figure 3), which can only be reported for those studies providing the concentration in both the aerosol and gas phases separately (39 studies, Table S4). The  $\log K_{RA}$  value ranged between 1.2 and 10.1, with such large variability observed mainly for PAHs. Such large variability of  $K_{RP}$ ,  $K_{RG}$ , and  $K_{RA}$  for PAHs is surprising. This is not due to limitations or difficulties in their chemical analysis as PAHs are at atmospheric concentrations several orders of magnitude higher than other POPs, such as PCBs, PCDDs/Fs, or OPEs. A characteristic of low MW PAHs is that they

degrade in the air and water by photodegradation and biodegradation.<sup>8,107</sup> High MW PAHs are protected by association, adsorption, or incorporation into the particles or in the black carbon.<sup>112,113</sup> Such degradation in rain could be a reason explaining such large variability of the rain–air partition constants for these chemicals. Confirmation of this hypothesis would require further work. Previously, it was shown that snow amplification of PAHs was reduced due to degradation,<sup>17</sup> and it could occur similarly for rain amplification.  $\log K_{RA}$  correlations with the chemical properties explained a small fraction of the variability for most chemical classes (Figure S7 and Table S12).

The amplification potential of POP fugacity by rain is given by

$$\frac{f_w}{f_g} = K_{RG}H'$$

where  $f_w$  and  $f_g$  are the POP fugacity in the deposited water and gas phases, respectively. Figure 4 shows the  $\log K_{RG}H'$  values of PCBs, OCPs, PAHs, PCDD/Fs, PBDEs, and three neutral PFAS. With the exception of some studies, there is an amplification of concentration in rain for all compounds, which is maximum for naphthalene, alkyl-naphthalenes, and neutral PFASs.

The amplification of fugacities by rain can occur when adsorption to raindrops is a significant process. This is the case for most POPs (Figure S6). Thus, there is a generalized amplification of the fugacities for all POPs (Figure 4), which surprisingly is especially more relevant for the more volatile chemicals, which can be as high as 6 orders of magnitude. For other POPs, the amplification potential is still important but generally below 3 orders of magnitude.

**Rain and Snow Amplification of POPs.** The investigation on which of the two wet deposition processes (snow or rain) is the most effective in scavenging organic compounds has been a recurrent topic in the “fate and transport” field,<sup>12,16,26,28,108</sup> but this comparison was often focused on predictions from models rather than field-derived assessments. Furthermore, snow has received more attention as an amplification mechanism for POPs than rain. This is especially relevant in polar environments, even though rainfall occurrence is predicted to increase in the coming decades.<sup>76</sup> The meta-analysis performed here allows for calculating the ratio between  $K_{RG}$  versus  $K_{SA}$  (the snow-air partition coefficient) from the field-derived data. For such a comparison, we use  $K_{SA}$  as estimated in a companion meta-analysis reported elsewhere.<sup>18</sup>

Figure 5 shows  $\log K_{RG}/K_{SA}$  for comparative purposes. Such comparisons could be done for PCBs, PAHs, neutral PFAS, and PBDEs. In addition to the mean, we calculated the error using the uncertainty propagation approach (Annex S3). The  $\log K_{RG}/K_{SA}$  value ranged between −3.16 and 1.9, presenting negative values of  $\log K_{RG}/K_{SA}$  for most PAHs and some PCB congeners, while positive values of  $\log K_{RG}/K_{SA}$  for PBDEs, low MW PAHs, and some PCB congeners. Therefore, the fieldwork carried out during the last four decades shows that snow and rain amplification of POPs are of comparable magnitude, with differences that are compound specific.

Snow deposition is limited to cold regions and/or cold periods of time, while rain precipitation occurs widely for different seasons and across climatic regions. Furthermore, there are observations that concentrations in rivers increase



after strong rain events.<sup>60,82,83,109–111</sup> Such large concentrations would be driven by the amplification of POPs by rain and the focusing of water in rivers from the watershed. Future work should be focused on studying the role of wet deposition on the cycle and occurrence of organic pollutants, especially in terms of its spatial and temporal dynamics, and extending this assessment to chemicals of emerging concern. Climate change induces a perturbation of the magnitude and frequency of precipitation events, which should be considered as a potential factor influencing the POP dynamics and amplification under a scenario of global environmental change.

## ■ ASSOCIATED CONTENT

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.est.1c03295>.

Analytical procedures, quality assurance/quality control, sampling location for the rain and aerosol samples, PFAS concentrations in aerosols and rain samples from Deception and Livingston Islands, OPEs and PAHs concentrations in aerosols and rain samples from Livingston Island, Meta-analysis of rain-air particulate partition constants, Pearson's correlations, details of target, recovery, and internal standards for PFAS, OPEs, and PAHs, PFAS, OPE, and PAH sample recoveries of recovery standards, details of limits of detection for PFAS and OPEs for rain and aerosol samples from Deception and Livingston Islands, data used in the meta-analysis of rain-air partition constants,  $K_{RP}$  mean,  $K_{RG}$  mean, and  $K_{RA}$  mean for each compound and each data set, dimensionless Henry's law constant values, and Pearson's correlations between  $\log K_{RA}$ ,  $\log K_{RG}$ , and  $\log K_{RP}$  with  $\log K_{aw}$ ,  $\log K_{oa}$ , and  $\log K_{ow}$  (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

Jordi Dachs – Institute of Environmental Assessment and Water Research, Spanish National Research Council (IDAEA-CSIC), Barcelona, Catalonia 08034, Spain; [orcid.org/0000-0002-4237-169X](https://orcid.org/0000-0002-4237-169X); Email: [jordi.dachs@idaea.csic.es](mailto:jordi.dachs@idaea.csic.es)

### Authors

Gemma Casas – Institute of Environmental Assessment and Water Research, Spanish National Research Council (IDAEA-CSIC), Barcelona, Catalonia 08034, Spain; Department of Instrumental Analysis and Environmental Chemistry, Institute of Organic Chemistry, Spanish National Research Council (IQOG-CSIC), Madrid 28006, Spain

Alicia Martínez-Varela – Institute of Environmental Assessment and Water Research, Spanish National Research Council (IDAEA-CSIC), Barcelona, Catalonia 08034, Spain

Maria Vila-Costa – Institute of Environmental Assessment and Water Research, Spanish National Research Council (IDAEA-CSIC), Barcelona, Catalonia 08034, Spain

Begoña Jiménez – Department of Instrumental Analysis and Environmental Chemistry, Institute of Organic Chemistry, Spanish National Research Council (IQOG-CSIC), Madrid 28006, Spain; [orcid.org/0000-0002-2401-4648](https://orcid.org/0000-0002-2401-4648)

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.est.1c03295>

## Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the staff of the Marine Technology Unit (UTM-CSIC) for their logistical support during the sampling campaign at Livingston Island, M. Pizarro for technical assistance, and The State Meteorological Agency (AEMET) for the meteorological assistance. This work was supported by Spanish Ministry of science to G.C. and A.M.-V. through predoctoral fellowships and projects SENTINEL (CTM2015-70535-P) and ANTOM (PGC2018-096612-B-I00). This research is part of POLARCSIC activities. The research group of Global Change and Genomic Biogeochemistry receives support from the Catalan Government (2017SGR800). Special thanks to TERNUA for sponsoring technical eco-friendly clothing and gear equipment for Antarctic campaigns.

## ■ REFERENCES

- (1) Lohmann, R.; Breivik, K.; Dachs, J.; Muir, D. Global Fate of POPs: Current and Future Research Directions. *Environ. Pollut.* **2007**, *150*, 150–165.
- (2) Casas, G.; Martínez-Varela, A.; Roscales, J. L.; Vila-Costa, M.; Dachs, J.; Jiménez, B. Enrichment of Perfluoroalkyl Substances in the Sea-Surface Microlayer and Sea-Spray Aerosols in the Southern Ocean. *Environ. Pollut.* **2020**, *267*, 115512.
- (3) Johansson, J. H.; Salter, M. E.; Acosta Navarro, J. C.; Leck, C.; Nilsson, E. D.; Cousins, I. T. Global Transport of Perfluoroalkyl Acids via Sea Spray Aerosol. *Environ. Sci.: Processes Impacts* **2019**, *21*, 635–649.
- (4) Wania, F.; Mackay, D. A Global Distribution Model for Persistent Organic Chemicals. *Sci. Total Environ.* **1995**, *160–161*, 211–232.
- (5) Dachs, J.; Lohmann, R.; Ockenden, W. A.; Méjanelle, L.; Eisenreich, S. J.; Jones, K. C. Oceanic Biogeochemical Controls on Global Dynamics of Persistent Organic Pollutants. *Environ. Sci. Technol.* **2002**, *36*, 4229–4237.
- (6) Galbán-Malagón, C.; Berrojalbiz, N.; Ojeda, M. J.; Dachs, J. The Oceanic Biological Pump Modulates the Atmospheric Transport of Persistent Organic Pollutants to the Arctic. *Nat. Commun.* **2012**, *3*, 862.
- (7) Galbán-Malagón, C. J.; Berrojalbiz, N.; Gioia, R.; Dachs, J. The “Degradative” and “Biological” Pumps Controls on the Atmospheric Deposition and Sequestration of Hexachlorocyclohexanes and Hexachlorobenzene in the North Atlantic and Arctic Oceans. *Environ. Sci. Technol.* **2013**, *47*, 7195–7203.
- (8) González-Gaya, B.; Martínez-Varela, A.; Vila-Costa, M.; Casal, P.; Cerro-Gálvez, E.; Berrojalbiz, N.; Lundin, D.; Vidal, M.; Mompeán, C.; Bode, A.; Jiménez, B.; Dachs, J. Biodegradation as an Important Sink of Aromatic Hydrocarbons in the Oceans. *Nat. Geosci.* **2019**, *12*, 119–125.
- (9) Jurado, E.; Dachs, J. Seasonality in the “Grasshopping” and Atmospheric Residence Times of Persistent Organic Pollutants over the Oceans. *Geophys. Res. Lett.* **2008**, *35*, L17805.
- (10) Dalla Valle, M.; Jurado, E.; Dachs, J.; Sweetman, A. J.; Jones, K. C. The Maximum Reservoir Capacity of Soils for Persistent Organic Pollutants: Implications for Global Cycling. *Environ. Pollut.* **2005**, *134*, 153–164.
- (11) Ockenden, W. A.; Breivik, K.; Meijer, S. N.; Steinnes, E.; Sweetman, A. J.; Jones, K. C. The Global Re-Cycling of Persistent Organic Pollutants Is Strongly Retarded by Soils. *Environ. Pollut.* **2003**, *121*, 75–80.
- (12) Bidelman, F. Wet and Dry Deposition of Organic Compounds Are Controlled by Their Vapor-Particle Partitioning. *Environ. Sci. Technol.* **1988**, *22*, 361–367.

- (13) Jurado, E.; Jaward, F.; Lohmann, R.; Jones, K. C.; Simó, R.; Dachs, J. Wet Deposition of Persistent Organic Pollutants to the Global Oceans. *Environ. Sci. Technol.* **2005**, *39*, 2426–2435.
- (14) Jurado, E.; Jaward, F. M.; Lohmann, R.; Jones, K. C.; Simó, R.; Dachs, J. Atmospheric Dry Deposition of Persistent Organic Pollutants to the Atlantic and Inferences for the Global Oceans. *Environ. Sci. Technol.* **2004**, *38*, 5505–5513.
- (15) Macdonald, R.; Mackay, D. Contaminant Amplification in the Environment. *Environ. Sci. Technol.* **2002**, *36*, 456A.
- (16) Meyer, T.; Wania, F. Organic Contaminant Amplification during Snowmelt. *Water Res.* **2008**, *42*, 1847–1865.
- (17) Casal, P.; Cabrerizo, A.; Vila-Costa, M.; Pizarro, M.; Jiménez, B.; Dachs, J. Pivotal Role of Snow Deposition and Melting Driving Fluxes of Polycyclic Aromatic Hydrocarbons at Coastal Livingston Island (Antarctica). *Environ. Sci. Technol.* **2018**, *52*, 12327–12337.
- (18) Casal, P.; Casas, G.; Vila-Costa, M.; Cabrerizo, A.; Pizarro, M.; Jiménez, B.; Dachs, J. Snow Amplification of Persistent Organic Pollutants at Coastal Antarctica. *Environ. Sci. Technol.* **2019**, *53*, 8872–8882.
- (19) Casal, P.; Zhang, Y.; Martin, J. W.; Pizarro, M.; Jiménez, B.; Dachs, J. Role of Snow Deposition of Perfluoroalkylated Substances at Coastal Livingston Island (Maritime Antarctica). *Environ. Sci. Technol.* **2017**, *51*, 8460–8470.
- (20) Bigot, M.; Hawker, D. W.; Cropp, R.; Muir, D. C.; Jensen, B.; Bossi, R.; Bengtson Nash, S. M. Spring Melt and the Redistribution of Organochlorine Pesticides in the Sea-Ice Environment: A Comparative Study between Arctic and Antarctic Regions. *Environ. Sci. Technol.* **2017**, *51*, 8944–8952.
- (21) Geisz, H. N.; Cochran, M. A.; Fraser, W. R.; Ducklow, H. W.; Ducklow, H. W. Melting Glaciers: A Probable Source of DDT to the Antarctic Marine Ecosystem. *Environ. Sci. Technol.* **2008**, *42*, 3958–3962.
- (22) Cabrerizo, A.; Dachs, J.; Moeckel, C.; Ojeda, M.-J.; Caballero, G.; Barceló, D.; Jones, K. C. Ubiquitous Net Volatilization of Polycyclic Aromatic Hydrocarbons from Soils and Parameters Influencing Their Soil-Air Partitioning. *Environ. Sci. Technol.* **2011**, *45*, 4740–4747.
- (23) Cabrerizo, A.; Dachs, J.; Barceló, D. Soil-Air Exchange Controls on Background Atmospheric Concentrations of Polychlorinated Biphenyls (PCBs); Organochlorine Pesticides (OCPs); and Polycyclic Aromatic Hydrocarbons (PAHs): A Case Study from Temperate Regions. *ACS Symposium Series, Occurrence, Fate and Impact of Atmospheric Pollutants on Environmental and Human Health*; American Chemical Society, 2013; Vol. 19–38, p 1149.
- (24) Khairy, M. A.; Dickhut, R.; Lohmann, R.; Lohmann, R. Levels; Sources and Chemical Fate of Persistent Organic Pollutants in the Atmosphere and Snow along the Western Antarctic Peninsula. *Environ. Pollut.* **2016**, *216*, 304–313.
- (25) Ma, J.; Hung, H.; Tian, C.; Kallenborn, R. Revolatilization of Persistent Organic Pollutants in the Arctic Induced by Climate Change. *Nat. Clim. Change* **2011**, *1*, 255–260.
- (26) Lei, Y. D.; Wania, F. Is Rain or Snow a More Efficient Scavenger of Organic Chemicals? *Atmos. Environ.* **2004**, *38*, 3557–3571.
- (27) Herbert, B. M. J.; Villa, S.; Halsall, C. J. Chemical Interactions with Snow: Understanding the Behavior and Fate of Semi-Volatile Organic Compounds in Snow. *Ecotoxicol. Environ. Saf.* **2006**, *63*, 3–16.
- (28) Wang, X.; Halsall, C.; Codling, G.; Xie, Z.; Xu, B.; Zhao, Z.; Xue, Y.; Ebinghaus, R.; Jones, K. C. Accumulation of Perfluoroalkyl Compounds in Tibetan Mountain Snow: Temporal Patterns from 1980 to 2010. *Environ. Sci. Technol.* **2014**, *48*, 173–181.
- (29) Sanchis, J.; Cabrerizo, A.; Galbán-Malagón, C.; Barceló, D.; Farré, M.; Dachs, J. Unexpected Occurrence of Volatile Dimethylsiloxanes in Antarctic Soils; Vegetation; Phytoplankton; and Krill. *Environ. Sci. Technol.* **2015**, *49*, 4415–4424.
- (30) Bidleman, T. F.; Christensen, E. J. Atmospheric Removal Processes for High Molecular Weight Organochlorines. *J. Geophys. Res., C: Oceans Atmos.* **1979**, *84*, 7857–7862.
- (31) Atlas, E.; Giam, C. S. Global Transport of Organic Pollutants: Ambient Concentrations in the Remote Marine Atmosphere. *Science* **1981**, *211*, 163–165.
- (32) Pankow, J. F.; Isabelle, L. M.; Asher, W. E. Trace Organic Compounds in rain. 1. Sampler Design and Analysis by Adsorption/Thermal Desorption (ATD). *Environ. Sci. Technol.* **1984**, *18*, 310–318.
- (33) Ligocki, M. P.; Leuenberger, C.; Pankow, J. F. Trace Organic Compounds in Rain-II. Gas Scavenging of Neutral Organic Compounds. *Atmos. Environ.* **1985**, *19*, 1609–1617.
- (34) Atlas, E.; Giam, C. S. Ambient Concentration and Precipitation Scavenging of Atmospheric Organic Pollutants. *Water, Air, Soil Pollut.* **1988**, *38*, 19–36.
- (35) McVeety, B. D.; Hites, R. A. Atmospheric Deposition of Polycyclic Aromatic Hydrocarbons to Water Surfaces: A Mass Balance Approach. *Atmos. Environ.* **1988**, *22*, 511–536.
- (36) Duinker, J. C.; Bouchertall, F. On the Distribution of Atmospheric Polychlorinated Biphenyl Congeners between Vapor Phase, Aerosols and Rain. *Environ. Sci. Technol.* **1989**, *23*, 57–62.
- (37) Eltzt, B. D.; Hites, R. A. Atmospheric Transport and Deposition of Polychlorinated Dibenzo-p-Dioxins and Dibenzofurans. *Environ. Sci. Technol.* **1989**, *23*, 1396–1401.
- (38) Koester, C. J.; Hites, R. A. Wet and Dry Deposition of Chlorinated Dioxins and Furans. *Environ. Sci. Technol.* **1992**, *26*, 1375.
- (39) Dickhut, R. M.; Gustafson, K. E. Atmospheric Washout of Polycyclic Aromatic Hydrocarbons in the Southern Chesapeake Bay Region. *Environ. Sci. Technol.* **1995**, *29*, 1518–1525.
- (40) Poster, D. L.; Baker, J. E. Influence of Submicron Particles on Hydrophobic Organic Contaminants in Precipitation. 1. Concentrations and Distributions of Polycyclic Aromatic Hydrocarbons and Polychlorinated Biphenyls in Rainwater. *Environ. Sci. Technol.* **1996**, *30*, 341–348.
- (41) Khalil Granier, L.; Chevreuil, M. Behaviour and Spatial and Temporal Variations of Polychlorinated Biphenyls and Lindane in the Urban Atmosphere of the Paris Area, France. *Atmos. Environ.* **1997**, *31*, 3787–3802.
- (42) Franz, T. P.; Eisenreich, S. J. Snow Scavenging of Polychlorinated Biphenyls and Polycyclic Aromatic Hydrocarbons in Minnesota. *Environ. Sci. Technol.* **1998**, *48*, 1771.
- (43) Wania, F.; Haugen, J.-E. Long Term Measurements of Wet Deposition and Precipitation Scavenging of Hexachlorocyclohexanes in Southern Norway. *Environ. Pollut.* **1999**, *105*, 381–386.
- (44) Park, J.-S.; Wade, T. L.; Sweet, S. T. Atmospheric Deposition of PAHs; PCBs; and Organochlorine Pesticides to Corpus Christi Bay; Texas. *Atmos. Environ.* **2002**, *36*, 1707–1720.
- (45) Van Ry, D. A.; Glenn, C. L.; Nelson, E. D.; Eisenreich, S. J.; Totten, L. A.; Eisenreich, S. J. Wet Deposition of Polychlorinated Biphenyls in Urban and Background Areas of the Mid-Atlantic States. *Environ. Sci. Technol.* **2002**, *36*, 3201–3209.
- (46) Offenberg, J. H.; Baker, J. E. Precipitation Scavenging of Polychlorinated Biphenyls and Polycyclic Aromatic Hydrocarbons along an Urban to Over-Water Transect. *Environ. Sci. Technol.* **2002**, *36*, 3763–3771.
- (47) Backe, C.; Larsson, P.; Agrell, C. Spatial and Temporal Variation of Polychlorinated Biphenyl (PCB) in Precipitation in Southern Sweden. *Sci. Total Environ.* **2002**, *285*, 117–132.
- (48) Agrell, C.; Larsson, P.; Okla, L.; Agrell, J. PCB Congeners in Precipitation; Wash out Ratios and Depositional Fluxes within the Baltic Sea Region; Europe. *Atmos. Environ.* **2002**, *36*, 371–383.
- (49) Takase, Y.; Murayama, H.; Mitobe, H.; Aoki, T.; Yagoh, H.; Shibuya, N.; Shimizu, K.-i.; Kitayama, Y. Persistent Organic Pollutants in Rain at Niigata; Japan. *Atmos. Environ.* **2003**, *37*, 4077–4085.
- (50) Mandalakis, M.; Stephanou, E. G. Wet Deposition of Polychlorinated Biphenyls in the Eastern Mediterranean. *Environ. Sci. Technol.* **2004**, *38*, 3011–3018.
- (51) Sahu, S. K.; Pandit, G. G.; Sadasivan, S. Precipitation Scavenging of Polycyclic Aromatic Hydrocarbons in Mumbai; India. *Sci. Total Environ.* **2004**, *318*, 245–249.

- (52) ter Schure, A. F. H.; Larsson, P.; Agrell, C.; Boon, J. P. Atmospheric Transport of Polybrominated Diphenyl Ethers and Polychlorinated Biphenyls to the Baltic Sea. *Environ. Sci. Technol.* **2004**, *38*, 1282–1287.
- (53) ter Schure, A. F. H.; Agrell, C.; Bokenstrand, A.; Sveder, J.; Larsson, P.; Zegers, B. N. Polybrominated Diphenyl Ethers at a Solid Waste Incineration Plant II: Atmospheric Deposition. *Atmos. Environ.* **2004**, *38*, 5149–5155.
- (54) Gioia, R.; Offenber, J. H.; Gigliotti, C. L.; Totten, L. A.; Du, S.; Eisenreich, S. J. Atmospheric Concentrations and Deposition of Organochlorine Pesticides in the US Mid-Atlantic Region. *Atmos. Environ.* **2005**, *39*, 2309–2322.
- (55) Tspakis, M.; Apostolaki, M.; Eisenreich, S.; Stephanou, E. G. Atmospheric Deposition and Marine Sedimentation Fluxes of Polycyclic Aromatic Hydrocarbons in the Eastern Mediterranean Basin. *Environ. Sci. Technol.* **2006**, *40*, 4922–4927.
- (56) Blanchard, M.; Teil, M. J.; Chevreuil, M. The Seasonal Fate of PCBs in Ambient Air and Atmospheric Deposition in Northern France. *J. Atmos. Chem.* **2006**, *53*, 123–144.
- (57) Barton, C. A.; Kaiser, M. A.; Russell, M. H. Partitioning and Removal of Perfluorooctanoate during Rain Events: The Importance of Physical-Chemical Properties. *J. Environ. Monit.* **2007**, *9*, 839–846.
- (58) Holoubek, I.; Klánová, J.; Jarkovský, J.; Kohoutek, J. Trends in Background Levels of Persistent Organic Pollutants at Kosetice Observatory, Czech Republic. Part I. Ambient Air and Wet Deposition 1996–2005. *J. Environ. Monit.* **2007**, *9*, 557–563.
- (59) Venier, M.; Hites, R. A. Atmospheric Deposition of PBDEs to the Great Lakes Featuring a Monte Carlo Analysis of Errors. *Environ. Sci. Technol.* **2008**, *42*, 9058–9064.
- (60) Mariani, G.; Canuti, E.; Castro-Jiménez, J.; Christoph, E. H.; Eisenreich, S. J.; Hanke, G.; Skejo, H.; Umlauf, G. Atmospheric Input of POPs into Lake Maggiore (Northern Italy): PBDE Concentrations and Profile in Air; Precipitation; Settling Material and Sediments. *Chemosphere* **2008**, *73*, S114–S121.
- (61) He, J.; Balasubramanian, R. A Study of Precipitation Scavenging of Semivolatile Organic Compounds in a Tropical Area. *J. Geophys. Res. Atmos.* **2009**, *114*, 12201.
- (62) Noël, M.; Dangerfield, N.; Hourston, R. A. S.; Belzer, W.; Shaw, P.; Yunker, M. B.; Ross, P. S. Do Trans-Pacific Air Masses Deliver PBDEs to Coastal British Columbia; Canada? *Environ. Pollut.* **2009**, *157*, 3404–3412.
- (63) Zhang, B.-Z.; Guan, Y.-F.; Zeng, E. Y.; Zeng, E. Y. Occurrence of Polybrominated Diphenyl Ethers in Air and Precipitation of the Pearl River Delta; South China: Annual Washout Ratios and Depositional Rates. *Environ. Sci. Technol.* **2009**, *43*, 9142–9147.
- (64) Birgül, A.; Tasdemir, Y.; Cindoruk, S. S. Atmospheric Wet and Dry Deposition of Polycyclic Aromatic Hydrocarbons (PAHs) Determined Using a Modified Sampler. *Atmos. Res.* **2011**, *101*, 341–353.
- (65) Dreyer, A.; Matthias, V.; Weinberg, I.; Ebinghaus, R. Wet Deposition of Poly- and Perfluorinated Compounds in Northern Germany. *Environ. Pollut.* **2010**, *158*, 1221–1227.
- (66) Günindi, M.; Tasdemir, Y. Wet and Dry Deposition Fluxes of Polychlorinated Biphenyls (PCBs) in an Urban Area of Turkey. *Water, Air, Soil Pollut.* **2011**, *215*, 427–439.
- (67) Liu, F.; Xu, Y.; Liu, J.; Liu, D.; Li, J.; Zhang, G.; Li, X.; Zou, S.; Lai, S. Atmospheric Deposition of Polycyclic Aromatic Hydrocarbons (PAHs) to a Coastal Site of Hong Kong; South China. *Atmos. Environ.* **2013**, *69*, 265–272.
- (68) Guo, L.-C.; Bao, L.-J.; Wu, F.-C.; Yue, Q.; Zeng, E. Y. Seasonal Deposition Fluxes and Removal Efficiency of Atmospheric Polybrominated Diphenyl Ethers in a Large Urban Center: Importance of Natural and Anthropogenic Factors. *Environ. Sci. Technol.* **2014**, *48*, 11196–11203.
- (69) Shahpoury, P.; Lammel, G.; Holubová Šmejkalová, A.; Klánová, J.; Přibylková, P.; Váňa, M. Polycyclic Aromatic Hydrocarbons; Polychlorinated Biphenyls; and Chlorinated Pesticides in Background Air in Central Europe - Investigating Parameters Affecting Wet Scavenging of Polycyclic Aromatic Hydrocarbons. *Atmos. Chem. Phys.* **2015**, *15*, 1795–1805.
- (70) Zhang, L.; Cheng, I.; Muir, D.; Charland, J.-P. Scavenging Ratios of Polycyclic Aromatic Compounds in Rain and Snow in the Athabasca Oil Sands Region. *Atmos. Chem. Phys.* **2015**, *15*, 1421–1434.
- (71) González-Gaya, B.; Fernández-Pinos, M.-C.; Méjanelle, L.; Abad, E.; Piña, B.; Duarte, C. M.; Jiménez, B.; Dachs, J. High Atmosphere–Ocean Exchange of Semivolatile Aromatic Hydrocarbons. *Nat. Geosci.* **2016**, *9*, 438–442.
- (72) Li, P.-h.; Wang, Y.; Li, Y.-h.; Wai, K.-m.; Li, H.-l.; Tong, L. Gas-Particle Partitioning and Precipitation Scavenging of Polycyclic Aromatic Hydrocarbons (PAHs) in the Free Troposphere in Southern China. *Atmos. Environ.* **2016**, *128*, 165–174.
- (73) IADN Data Visualization, <https://iadviz.iu.edu/datasets/index.html> (accessed on July 8th, 2021).
- (74) Zhang, Z.; Lin, G.; Lin, T.; Zhang, R.; Jin, L.; Di, Y. Occurrence; Behavior; and Fate of Organophosphate Esters (OPEs) in Subtropical Paddy Field Environment: A Case Study in Nanning City of South China. *Environ. Pollut.* **2020**, *267*, 115675.
- (75) Simcik, M. The Importance of Surface Adsorption on the Washout of Semivolatile Organic Compounds by Rain. *Atmos. Environ.* **2004**, *38*, 491–501.
- (76) Vignon, É.; Roussel, M.; Gorodetskaya, I. V.; Genthon, C.; Berne, A. Present and Future of Rainfall in Antarctica. *Geophys. Res. Lett.* **2021**, *48*, No. e2020GL092281.
- (77) Yamashita, N.; Kannan, K.; Taniyasu, S.; Horii, Y.; Okazawa, T.; Petrick, G.; Gamo, T. Analysis of Perfluorinated Acids at Parts-per-Quadrillion Levels in Seawater Using Liquid Chromatography-Tandem Mass Spectrometry. *Environ. Sci. Technol.* **2004**, *38*, 5522–5528.
- (78) Sun, Y.; De Silva, A. O.; Muir, D. C. G.; Spencer, C.; Lehnher, I.; MacInnis, J. J.; MacInnis, J. J. Glacial Melt Inputs of Organophosphate Ester Flame Retardants to the Largest High Arctic Lake. *Environ. Sci. Technol.* **2020**, *54*, 2734–2743.
- (79) Zhong, M.; Tang, J.; Guo, X.; Guo, C.; Li, F.; Wu, H. Occurrence and Spatial Distribution of Organophosphorus Flame Retardants and Plasticizers in the Bohai; Yellow and East China Seas. *Sci. Total Environ.* **2020**, *741*, 140434.
- (80) Van Noort, P. C. M.; Wondergem, E. Scavenging of Airborne Polycyclic Aromatic Hydrocarbons by Rain. *Environ. Sci. Technol.* **1985**, *19*, 1044–1048.
- (81) Leuenberger, C.; Czuczwa, J.; Heyerdahl, E.; Giger, W. Aliphatic and Polycyclic Aromatic Hydrocarbons in Urban Rain; Snow and Fog. *Atmos. Environ.* **1988**, *22*, 695–705.
- (82) Totten, L. A.; Panangadan, M.; Eisenreich, S. J.; Cavallo, G. J.; Fikslin, T. J. Direct and Indirect Atmospheric Deposition of PCBs to the Delaware River Watershed. *Environ. Sci. Technol.* **2006**, *40*, 2171–2176.
- (83) Kim, S.-K.; Kannan, K. Perfluorinated Acids in Air; Rain; Snow; Surface Runoff; and Lakes: Relative Importance of Pathways to Contamination of Urban Lakes. *Environ. Sci. Technol.* **2007**, *41*, 8328–8334.
- (84) Motelay-Massei, A.; Ollivon, D.; Garban, B.; Tiphagne-Larcher, K.; Zimmerlin, I.; Chevreuil, M. PAHs in the Bulk Atmospheric Deposition of the Seine River Basin: Source Identification and Apportionment by Ratios; Multivariate Statistical Techniques and Scanning Electron Microscopy. *Chemosphere* **2007**, *67*, 312–321.
- (85) Taniyasu, S.; Kannan, K.; Yeung, L. W. Y.; Kwok, K. Y.; Lam, P. K. S.; Yamashita, N. Analysis of Trifluoroacetic Acid and Other Short-Chain Perfluorinated Acids (C2–C4) in Precipitation by Liquid Chromatography-Tandem Mass Spectrometry: Comparison to Patterns of Long-Chain Perfluorinated Acids (C5–C18). *Anal. Chim. Acta* **2008**, *619*, 221–230.
- (86) Delhomme, O.; Rieb, E.; Millet, M. Polycyclic Aromatic Hydrocarbons Analyzed in Rainwater Collected on Two Sites in East of France (Strasbourg and Erstein). *Polycyclic Aromatic Compounds*; Taylor & Francis Group, 2008; Vol. 28, pp 472–485.



- (87) Regnery, J.; Püttmann, W. Organophosphorus Flame Retardants and Plasticizers in Rain and Snow from Middle Germany. *Clean: Soil, Air, Water* **2009**, *37*, 334–342.
- (88) Liu, W.; Jin, Y.; Sasaki, K.; Saito, N.; Nakayama, S. F.; Sato, I.; Tsuda, S. Perfluorosulfonates and Perfluorocarboxylates in Snow and Rain in Dalian, China. *Environ. Int.* **2009**, *35*, 737–742.
- (89) Gaga, E. O.; Tuncel, S. G. Sources and Wet Deposition Fluxes of Polycyclic Aromatic Hydrocarbons (PAHs) in an Urban Site 1000 Meters High in Central Anatolia (Turkey). *Environ. Forensics* **2009**, *10*, 286–298.
- (90) Wang, Y.; Li, P.-h.; Li, H.-l.; Liu, X.-h.; Wang, W.-x. PAHs Distribution in Precipitation at Mount Taishan, China. Identification of Sources and Meteorological Influences. *Atmos. Res.* **2010**, *95*, 1–7.
- (91) Cavalcante, R. M.; Sousa, F. W.; Nascimento, R. F.; Silveira, E. R.; Viana, R. B. Influence of Urban Activities on Polycyclic Aromatic Hydrocarbons in Precipitation: Distribution; Sources and Depositional Flux in a Developing Metropolis; Fortaleza; Brazil. *Sci. Total Environ.* **2012**, *414*, 287–292.
- (92) Murray, M. W.; Andren, A. W. Precipitation Scavenging of Polychlorinated Biphenyl Congeners in the Great Lakes Region. *Atmos. Environ., Part A* **1992**, *26*, 883–897.
- (93) Cetin, B.; Odabasi, M.; Bayram, A. Wet Deposition of Persistent Organic Pollutants (POPs) in Izmir; Turkey. *Environ. Sci. Pollut. Res.* **2016**, *23*, 9227–9236.
- (94) Shimizu, M. S.; Mott, R.; Potter, A.; Zhou, J.; Baumann, K.; Surratt, J. D.; Turpin, B.; Avery, G. B.; Harfmann, J.; Kieber, R. J.; Mead, R. N.; Skrabal, S. A.; Willey, J. D. Atmospheric Deposition and Annual Flux of Legacy Perfluoroalkyl Substances and Replacement Perfluoroalkyl Ether Carboxylic Acids in Wilmington; NC; USA. *Environ. Sci. Technol. Lett.* **2021**, *8*, 366–372.
- (95) Škrdlíková, L.; Landlová, L.; Klánová, J.; Lammel, G. Wet Deposition and Scavenging Efficiency of Gaseous and Particulate Phase Polycyclic Aromatic Compounds at a Central European Suburban Site. *Atmos. Environ.* **2011**, *45*, 4305–4312.
- (96) Van Drooge, B. L.; Torres-García, C. J.; Cuevas, E. Deposition of Semi-Volatile Organochlorine Compounds in the Free Troposphere of the Eastern North Atlantic Ocean. *Mar. Pollut. Bull.* **2001**, *42*, 628–634.
- (97) Ollivon, D.; Blanchoud, H.; Motelay-Massei, A.; Garban, B. Atmospheric Deposition of PAHs to an Urban Site, Paris, France. *Atmos. Environ.* **2002**, *36*, 2891–2900.
- (98) De Rossi, C.; Bierl, R.; Riefstahl, J. Organic Pollutants in Precipitation: Monitoring of Pesticides and Polycyclic Aromatic Hydrocarbons in the Region of Trier (Germany). *Phys. Chem. Earth* **2003**, *28*, 307–314.
- (99) Berger, U.; Jarnberg, U.; Kallenborn, R. *Perfluorinated Alkylated Substances (PFAS) in the European Nordic Environment*, 2004.
- (100) Loewen, M.; Halldorson, T.; Wang, F.; Tomy, G. Fluorotelomer Carboxylic Acids and PFOS in Rainwater from an Urban Center in Canada. *Environ. Sci. Technol.* **2005**, *39*, 2944–2951.
- (101) Scott, B. F.; Spencer, C.; Mabury, S. A.; Muir, D. C. G. Poly and Perfluorinated Carboxylates in North American Precipitation. *Environ. Sci. Technol.* **2006**, *40*, 7167–7174.
- (102) Mihajlović, I.; Fries, E. Atmospheric Deposition of Chlorinated Organophosphate Flame Retardants (OFR) onto Soils. *Atmos. Environ.* **2012**, *56*, 177–183.
- (103) Wang, C.; Wang, P.; Zhao, J.; Fu, M.; Zhang, L.; Li, Y.; Yang, R.; Zhu, Y.; Fu, J.; Zhang, Q.; Jiang, G. Atmospheric Organophosphate Esters in the Western Antarctic Peninsula over 2014–2018: Occurrence; Temporal Trend and Source Implication. *Environ. Pollut.* **2020**, *267*, 115428.
- (104) Cabrerizo, A.; Galbán-Malagón, C.; Del Vento, S.; Dachs, J. Sources and Fate of Polycyclic Aromatic Hydrocarbons in the Antarctic and Southern Ocean Atmosphere. *Global Biogeochem. Cycles* **2014**, *28*, 1424–1436.
- (105) Mackay, D.; Paterson, S.; Schroeder, W. H. Model Describing the Rates of Transfer Processes of Organic Chemicals Between Atmosphere and Water. *Environ. Sci. Technol.* **1986**, *20*, 810–816.
- (106) Dreyer, A.; Weinberg, I.; Temme, C.; Ebinghaus, R. Polyfluorinated Compounds in the Atmosphere of the Atlantic and Southern Oceans: Evidence for a Global Distribution. *Environ. Sci. Technol.* **2009**, *43*, 6507–6514.
- (107) Keyte, I. J.; Harrison, R. M.; Lammel, G. Chemical Reactivity and Long-Range Transport Potential of Polycyclic Aromatic Hydrocarbons—a Review. *Chem. Soc. Rev.* **2013**, *42*, 9333–9391.
- (108) Wania, F.; Hoff, J. T.; Jia, C. Q.; Mackay, D. The Effects of Snow and Ice on the Environmental Behaviour of Hydrophobic Organic Chemicals. *Environ. Pollut.* **1998**, *102*, 25–41.
- (109) Capel, P. D.; Larson, S. J.; Winterstein, T. A. The Behaviour of 39 Pesticides in Surface Waters as a Function of Scale. *Hydrol. Process.* **2001**, *15*, 1251–1269.
- (110) Olivella, M. À. Polycyclic Aromatic Hydrocarbons in Rainwater and Surface Waters of Lake Maggiore; a Subalpine Lake in Northern Italy. *Chemosphere* **2006**, *63*, 116–131.
- (111) Chen, M.; Wang, C.; Wang, X.; Fu, J.; Gong, P.; Yan, J.; Yu, Z.; Yan, F.; Nawab, J. Release of Perfluoroalkyl Substances From Melting Glacier of the Tibetan Plateau: Insights Into the Impact of Global Warming on the Cycling of Emerging Pollutants. *J. Geophys. Res. Atmos.* **2019**, *124*, 7442–7456.
- (112) Pankow, J. F.; Bidleman, T. F. Effects of Temperature, TSP and per Cent Non-Exchangeable Material in Determining the Gas-Particle Partitioning of Organic Compounds. *Atmos. Environ., Part A* **1991**, *25*, 2241–2249.
- (113) Dachs, J.; Eisenreich, S. J. Adsorption onto aerosol soot carbon dominates gas-particle partitioning of polycyclic aromatic hydrocarbons. *Environ. Sci. Technol.* **2000**, *34*, 3690–3697.