



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

Correspondence

Re: “Endothelitis in COVID-19-Positive Patients after Extremity Amputation for Acute Thrombotic Events”



To Editor,

Ilonzo et al.¹ reported, in their recent and interesting article, their clinical experience in 4 patients affected by COVID-19 and undergoing major limb amputation secondary to acute irreversible ischemia. On histological examination with hematoxylin/eosin, they found inflammatory cells associated with endothelium/apoptotic bodies, mononuclear cells, small vessel congestion, and lymphocytic endotheliitis and concluded that the findings in these patients are more likely an infectious angiitis due to COVID-19.

COVID-19 may predispose to both venous and arterial thromboembolic disease due to high-grade inflammation, hypoxia, immobilization, and diffuse intravascular coagulation, but direct damage to the endothelium by the virus is not yet demonstrated. Conversely, the microorganisms can generally induce the expression of numerous factors, including tissue factor on monocytes and macrophages, by binding to pattern-recognizing receptors on immune cells. The triggering of host inflammatory reaction also results in increased production of pro-inflammatory cytokines that have pleiotropic effects, including activation of coagulation.

In our experience, we have observed numerous cases of venous and arterial thromboembolism not only in the acute phase of COVID-19, but (even more interestingly) even after recovery.²⁻⁶

Despite successful treatment of initial COVID19, there is the possibility of acute thrombosis during follow-up, despite normalization of hemostatic and inflammatory parameters. The real prevalence of this event is difficult to determine. We think that our patients support the hypothesis that the integrity and functional characteristics of the endothelial cells, initially deranged during the viral infection may persist for a longer period, despite apparent normalization of hemostatic parameters.⁷⁻⁹

Whether SARS-CoV-2 is able to directly attack vascular endothelial cells expressing high levels of ACE2, and then lead to abnormal coagulation and sepsis, still needs to be explored. The question of how the SARS-CoV-2 spreads to extrapulmonary organs and the mechanism of endothelial damage remains an enigma.

DECLARATION OF COMPETING INTEREST

None.

*Antonio Bozzani**, *Vittorio Arici*
Vascular and Endovascular
Surgery, Foundation I.R.C.C.S.
Policlinico San Matteo, Pavia,
Italy

Guido Tavazzi
Anesthesiology and Intensive
Care Unit, Fondazione IRCCS
Policlinico San Matteo, Pavia,
Italy
Department of Medical, Surgical,
Diagnostic and Pediatric Science,
University of Pavia, Pavia, Italy

Stefano Boschini
Vascular and Endovascular
Surgery, Foundation I.R.C.C.S.
Policlinico San Matteo, Pavia,
Italy

Francesco Mojoli
Anesthesiology and Intensive
Care Unit, Fondazione IRCCS
Policlinico San Matteo, Pavia,
Italy
Department of Medical, Surgical,
Diagnostic and Pediatric Science,
University of Pavia, Pavia, Italy

Raffaele Bruno
Infectious Diseases Unit,
Fondazione IRCCS Policlinico
San Matteo, Pavia, Italy
Department of Medical, Surgical,
Diagnostic and Pediatric Science,
University of Pavia, Pavia, Italy

Antonio V. Sterpetti
The Sapienza University of Rome

Franco Ragni
Vascular and Endovascular
Surgery, Foundation I.R.C.C.S.
Policlinico San Matteo, Pavia,
Italy

Correspondence to: Antonio Bozzani, MD, Vascular and Endovascular Surgery, Foundation I.R.C.C.S. Policlinico San Matteo, P.le Golgi 19, 27100 Pavia, Italy
E-mail: a.bozzani@smatteo.pv.it

REFERENCES

1. Ilonzo N, Kumar S, Borazan N, et al. Endothelitis in COVID-19-positive patients after extremity amputation for acute thrombotic events. *Ann Vasc Surg* 2020; S0890-5096(20)31101-8. doi:[10.1016/j.avsg.2020.12.004](https://doi.org/10.1016/j.avsg.2020.12.004).
2. Bozzani A, Arici V, Tavazzi G, et al. Acute arterial and deep venous thromboembolism in COVID-19 patients: risk factors and personalized therapy. *Surgery* 2020;168:987–92. doi:[10.1016/j.surg.2020.09.009](https://doi.org/10.1016/j.surg.2020.09.009).
3. Bozzani A, Tavazzi G, Arici V, et al. Acute deep vein thrombosis in COVID 19 hospitalized patients. Risk factors and clinical outcomes. *Phlebology* 2020 Sep 14;268355520958598. Online ahead of print. doi:[10.1177/0268355520958598](https://doi.org/10.1177/0268355520958598).
4. Bozzani A, Arici V, Franciscone MM, et al. Severe acute respiratory syndrome coronavirus 2 infection and the upper limb deep vein thrombosis risk. *Ann Vasc Surg* 2020;66:11–13 Epub 2020 Apr 23. doi:[10.1016/j.avsg.2020.04.037](https://doi.org/10.1016/j.avsg.2020.04.037).
5. Perotti C, Arici V, Cervio M, et al. Allogeneic lethally irradiated cord blood mononuclear cells in no-option critical limb ischemia: a "box of rain". *Stem Cells Dev* 2013;22:2806–12 Epub 2013 Jul 13. doi:[10.1089/scd.2013.0172](https://doi.org/10.1089/scd.2013.0172).
6. Arici V, Bozzani A, Odero A. Successful endovascular treatment of a bronchial artery aneurysm refractory to transcatheter embolization. *Ann Vasc Surg* 2013;27 802.e13-5. doi:[10.1016/j.avsg.2012.07.029](https://doi.org/10.1016/j.avsg.2012.07.029).
7. Arici V, Perotti C, Fabrizio C, et al. Autologous immuno magnetically selected CD133+ stem cells in the treatment of no-option critical limb ischemia: clinical and contrast enhanced ultrasound assessed results in eight patients. *J Transl Med* 2015;13:342. doi:[10.1186/s12967-015-0697-4](https://doi.org/10.1186/s12967-015-0697-4).
8. Bozzani A, Arici V, Tavazzi G, et al. Acute thrombosis of lower limbs arteries in the acute phase and after recovery from COVID19. *Ann Surg* 2021 Online ahead of print. doi:[10.1097/SLA.0000000000004700](https://doi.org/10.1097/SLA.0000000000004700).
9. Bozzani A, Pallini M, Arici V, et al. Re-organization of the vascular surgery department during the acute phase of the COVID19 outbreak: lessons learned and future perspectives. *Ann Vasc Surg* 2021 Online ahead of print. doi:[10.1016/j.avsg.2020.12.001](https://doi.org/10.1016/j.avsg.2020.12.001).