


Radial incision and cutting under gel immersion as a useful method for refractory anastomotic stricture




► **Fig. 1** The stricture was refractory and difficult for the scope to pass through despite endoscopic balloon dilation.

The radial incision and cutting method, in which the stricture is incised and the scar tissue excised, has shown promising results in the treatment of refractory anastomotic strictures [1,2]. It is useful for refractory strictures that are not adequately dilated after repeated endoscopic balloon dilation.

Gel immersion endoscopy with clear gel (VISCOCLEAR; Otsuka Pharmaceuticals Factory, Chiyoda, Japan) is used to displace blood and clots to ensure a good field of view in situations with heavy bleeding or residue [3,4].

The patient was a 47-year-old man who underwent low anterior resection with creation of a temporary colostomy for rectal cancer. However, a postoperative anastomotic stricture was observed, and endoscopic balloon dilation was performed twice on an outpatient basis after discharge. Because the stricture was refractory and resulted in difficulty passing the scope despite the two dilations (► **Fig. 1**), radial incision and cutting was performed (► **Video 1**).

An ITknife nano (KD-612; Olympus Corp., Tokyo, Japan) was inserted into the stricture, and the scar was incised in the endo cut mode. The incision with the endo cut caused a multifocal hemorrhage resulting in difficulty maintaining the field of




► **Video 1** Radial incision and cutting under gel immersion for the treatment of refractory anastomotic stenosis is safe and feasible.


view. When the gel was used, the bleeding point became clear, and it was possible to make an incision with a good field of view while recognizing the line of the muscle layer even with heavy bleeding (► **Fig. 2**). After the procedure, passage of the scope was possible, and no complications were observed (► **Fig. 3**). Endoscopy showed improvement of the stricture 2 weeks later (► **Fig. 4**).

Gel immersion endoscopy ensured a good field of view, even in situations with heavy bleeding. Radial incision and cutting under gel immersion is a safe and efficient method for the treatment of refractory colorectal anastomotic strictures.


Endoscopy_UCTN_Code_CCL_1AB_2AD_3AF



► **Fig. 2** It was possible to make an incision with a good field of view under gel immersion endoscopy even in a situation with heavy bleeding.



► **Fig. 3** It was possible for the scope to pass through after radial incision and cutting.




► **Fig. 4** Endoscopy showed improvement of the stricture 2 weeks after the procedure.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Keisaku Yamada, Masahiro Tajika  **Tsutomu Tanaka, Sachiyo Oonishi, Tomoyasu Kamiya, Yasumasa Niwa**

Department of Endoscopy, Aichi cancer center Hospital, Nagoya, Japan

Corresponding author

Masahiro Tajika, MD
Department of Endoscopy, Aichi Cancer Center Hospital, 1-1 Kanokoden, Chikusa-ku, Nagoya 464-8681, Japan
Fax: +81-52-763-5233
mtajika@aichi-cc.jp

References

- [1] Inoue T, Ibusuki M, Kitano R et al. Radial incision and cutting combined with balloon dilation for refractory pancreaticojejunostomy anastomotic strictures. *Endosc Int Open* 2021; 9: E578–E582
- [2] Osera S, Ikematsu H, Odagaki T et al. Efficacy and safety of endoscopic radial incision and cutting for benign severe anastomotic stricture after surgery for lower rectal cancer (with video). *Gastrointest Endosc* 2015; 81: 770–773
- [3] Yano T, Nemoto D, Ono K et al. Gel immersion endoscopy: a novel method to secure the visual field during endoscopy in bleeding patients (with videos). *Gastrointest Endosc* 2016; 83: 809–811
- [4] Miura Y, Yano T, Takezawa T et al. Gel immersion endoscopy simplifies hemostasis during endoscopic submucosal dissection using the pocket-creation method. *Endoscopy* 2018; 50: E294–E295

Bibliography

Endoscopy 2022; 54: E832–E833

DOI 10.1055/a-1824-4843

ISSN 0013-726X

published online 13.5.2022

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>