

SHORT PAPER

Open Access

Preliminary evidence about the effects of meditation on interoceptive sensitivity and social cognition

Margherita Melloni^{1,2,5}, Lucas Sedeño^{1,2,5}, Blas Couto^{1,2,5}, Martín Reynoso¹, Carlos Gelormini^{1,2,5}, Roberto Favaloro¹, Andrés Canales-Johnson^{2,3}, Mariano Sigman^{4,5,6}, Facundo Manes^{1,2,5,7} and Agustín Ibáñez^{1,2,5*}

Abstract

Background: Interoception refers to the conscious perception of body signals. Mindfulness is a meditation practice that encourages individuals to focus on their internal experiences such as bodily sensations, thoughts, and emotions. In this study, we selected a behavioral measure of interoceptive sensitivity (heartbeat detection task, HBD) to compare the effect of meditation practice on interoceptive sensitivity among long term practitioners (LTP), short term meditators (STM, subjects that completed a Mindfulness-Based Stress Reduction (MBSR) program) and controls (non-meditators). All participants were examined with a battery of different tasks including mood state, executive function and social cognition tests (emotion recognition, empathy and theory of mind).

Findings: Compared to controls, both meditators' groups showed lower levels of anxiety and depression, but no improvement in executive function or social cognition performance was observed (except for lower scores compared to controls only in the personal distress dimension of empathy). More importantly, meditators' performance did not differ from that of nonmeditators regarding cardiac interoceptive sensitivity.

Conclusion: Results suggest no influence of meditation practice in cardiac interoception and in most related social cognition measures. These negative results could be partially due to the fact that awareness of heartbeat sensations is not emphasized during mindfulness/vipassana meditation and may not be the best index of the awareness supported by the practice of meditation.

Keywords: Interoception, Meditation, Mindfulness, Social cognition, Heartbeat detection task

Background

Interoception involves the conscious perception of feelings from inside the body [1-3]. Interoception has been proposed to modulate social cognition processes such as motivational behavior [2], empathy [4], and theory of mind (ToM), which have been suggested to be supported by emotional and body feedback information [4].

Meditation is a form of mental training [5] encouraging individuals to focus on their internal experiences, such as bodily sensations, thoughts, and emotions [6].

One component of meditation involves the development of interoceptive attention to visceral sensations [7]. Additionally, meditation practice promotes the development of prosocial behavior [8].

Previous findings reported no difference in interoception accuracy between meditators and nonmeditators [5,9]. In these studies, a heartbeat discrimination paradigm was selected: participants had to discriminate whether their heartbeats synchronized with either auditory or visual cues [10]. Consequently, subjects had to attend at the same time to their cardiac sensation and to external stimuli which have been shown to affect interoceptive performance [11]. We selected a different heartbeat detection paradigm [12] to avoid the possible interference of external stimuli. Moreover, given the relationship between interoception and social cognition [2,4,13,14], we included

* Correspondence: aibanez@ineco.org.ar

¹Laboratory of Experimental Psychology and Neuroscience (LPEN), INECO (Institute of Cognitive Neurology) and Institute of Neuroscience, Favaloro, Favaloro University, C1078AAI, Buenos Aires, Argentina

²UDP-INECO Foundation Core on Neuroscience (UICoN), Diego Portales University, Santiago, Chile

Full list of author information is available at the end of the article

tasks of emotion recognition, empathy and ToM to test the association among bodily perception, social cognition and meditation practice. Moreover, considering the interaction between executive functions (EF) and social cognition domains (emotional processing [15], ToM [16] and empathy [17]), EF abilities were also evaluated.

Our aim was to compare the effect of meditation practice on interoceptive sensitivity and related measures among long term practitioners (LTP), subjects that completed a Mindfulness-Based Stress Reduction (MBSR) program (short term meditators, STM) and controls (nonmeditators). We predicted that long term practitioners would show enhanced interoceptive sensitivity, reflected in a better performance in heartbeat detection and related domains of social cognition.

Methods

Subjects

Ten nonmeditators, 9 short-term meditators and 10 long-term practitioners participated. The LTP group's mean was 4.35 (SD = 2.17) years of continued practice and the STM completed an 8-week Mindfulness-Based Stress Reduction (MBSR) program (see criteria in the Additional file 1: Table S1). Controls had never attended a yoga or meditation course. Groups were age, gender and education matched. We controlled body mass index because it influences the interoceptive performance [18]. Participants had no history of drug abuse, neurological or psychiatric conditions. Participants provided an informed consent in accordance with the Declaration of Helsinki and the study was approved by the institutional ethics committee.

Neuropsychological and clinical evaluation

Participants completed the Beck's Depression Inventory (BDI) and the State Trait Anxiety Inventory (STAI) to evaluate mood and affective state, respectively. EF were assessed with the INECO Frontal Screening (IFS) [19] indexing 8 EF (see Additional file 1: Table S1) and the Stroop test.

Social cognition tasks

A description of social cognition tasks (empathy, theory of mind and emotion recognition) is provided in Table 1 (see also Additional file 1: Table S1 for a detailed explanation of the materials and methods).

Interoception

Heartbeat detection task (HBD)

The HBD is a motor tracking test that assesses interoception sensitivity [12]. Participants had to tap a key on a keyboard along with their heartbeat in different conditions (see Table 1 and the Additional file 1: Table S1 data for a more detailed explanation).

Data analysis

Demographic, neuropsychological, and experimental data were compared among groups using ANOVA and Tukey's HSD *post-hoc* tests. For categorical variables (e.g., gender), Kruskal-Wallis tests were applied. Mixed repeated measured ANOVA was performed for HBD, with a within-subject factor (the four conditions) and a between-subject factor (group).

Table 1 Interoception and social cognition domain assessed and tasks employed

Interoception	Task	Description
Interoception sensitivity	HBD	The HBD is a motor tracking test that assesses interoception sensitivity. Participants had to tap a key on a keyboard along with their heartbeat in different conditions. First, as a motor-control condition, participants had to follow an audio-recording of a heartbeat. Next, they had to follow their heartbeat without external feedback (intero-pre condition). Then they had to do the same while receiving simultaneous auditory feedback of their own heart provided through online EKG register (feedback condition). Finally, they had to follow their own heartbeat without feedback (intero-post condition). These conditions offer a measure of audio-motoric performance (first condition), and a cardiac interoceptive measure (second and fourth conditions), prior to and after the feedback condition, respectively. During this task we also measured heart rate and heart rate variability to control their possible influence on interoception sensitivity; results showed no differences among groups (see Additional file 2).
Social cognition domain	Task	Description
Emotional recognition	Emotional morphing	This task assesses recognition of six basic emotion expressions and consists of 48 morphing faces randomly presented on a screen (see Additional file 1).
Theory of mind	ToM	This test assesses the emotional inference aspect of the ToM. Consist of 36 pictures of the eye region of a face. Participants chose which of four words best described the person's thoughts or feelings in each picture.
Empathy	IRI	The IRI is a 28-item self-report questionnaire that separately measures both the cognitive and affective components of empathy.

HBD=Heartbeat Detection task; ToM=Theory of Mind; IRI=Index of Interpersonal Reactivity.

Results

Demographic and neuropsychological results

No differences were found in gender [$H = 4.90$, $p = 0.86$], age [$F (2, 25) = 0.95$, $p = 0.39$, $\eta_p^2 = 0.07$], formal education [$F (2, 25) = 2.13$, $p = 0.13$, $\eta_p^2 = 0.14$] or body mass index [$F (2, 21) = 1.47$, $p = 0.25$, $\eta_p^2 = 0.12$] among groups.

Groups showed similar EF performance measured by the IFS [$F (2, 25) = 1.50$, $p = 0.24$, $\eta_p^2 = 0.10$]. There were no differences in the three condition of the Stroop task, word [$F (2, 23) = 0.20$, $p = 0.81$, $\eta_p^2 = 0.01$], color [$F (2, 23) = 1.40$, $p = 0.26$, $\eta_p^2 = 0.10$] and incongruent color-word [$F (2, 23) = 0.35$, $p = 0.70$, $\eta_p^2 = 0.03$]. No interference effect was found [$F (2, 23) = 1.88$, $p = 0.17$, $\eta_p^2 = 0.14$] (See Table 2).

Clinical evaluation

We observed a significant difference for BDI score among groups [$F (2, 25) = 4.12$, $p < 0.05$, $\eta_p^2 = 0.24$]. Post-hoc comparisons (Tukey HSD test, MS = 34.97;

$df = 25.00$) revealed higher scores of depressive symptoms in controls compared to STM ($p < 0.05$). We did not observe between group differences for STAI-State subscale [$F (2, 25) = 1.87$, $p = 0.17$, $\eta_p^2 = 0.13$]. However, significant differences for STAI-Trait subscale [$F (2, 25) = 3.74$, $p < 0.05$, $\eta_p^2 = 0.23$] were observed; post hoc comparisons (Tukey test, HSD, MS = 69.98; $df = 25.00$) showed controls had significantly higher anxiety scores ($p < 0.05$) than LTM.

Social cognition measures

Emotion recognition: No differences were observed regarding total accuracy [$F (2, 25) = 2.49$, $p = 0.10$, $\eta_p^2 = 0.16$]. However, per category analysis showed significant differences in disgust recognition among groups [$F (2, 25) = 4.1$, $p < 0.05$, $\eta_p^2 = 0.24$]. A post-hoc comparison (Tukey HSD test, MS = 0.01; $df = 25.00$) revealed lower accuracy performance in LTM group ($p < 0.05$) than controls (see Figure 1a). Groups did not differ regarding RTs of average


Table 2 Demographic, neuropsychological and clinical results

	F	p	Controls	STM	LTM
Gender	4.90 (H)	.86	2 M : 8 F M= 37.30; SD= 9.12 (22 – 49) M=16.10;	4 M : 4 F M= 41.12; SD= 12.15 (25 – 55) M= 16.13;	7 M : 3 F M= 43.80; SD=10.55 M= 17.90;
Age (years)	.95	.40			
Formal education (years)	2.13	.13	SD=.74 (15 – 17) M= 22.94;	SD= 1.73 (12 – 17) M= 22.88;	SD= 3.25 (12 – 25) M= 24.87;
Body mass index	1.47	.25	SD= 2.75 (19.43 – 26.67) M= 25.95;	SD= 3.15 (17.63 – 26.51) M= 27.44;	SD= 2.49 (21.60 – 28.57) M= 26.25;
IFS (Ineco Frontal Score)	1.50	.24	SD= 1.50 (23 – 28) M= 5.02;	SD= 2.47 (23 – 30) M= 11.33;	SD= 1.69 (24 – 30) M= 5.36;
Stroop Interference score	1.88	.17	SD= 8.06 (-6.59 – 15.95) M=9.90;	SD= 7.20 (5.54 – 27.70) M=2.88;	SD= 7.03 (-2.64 – 15.85) M=3.50;
BDI-II	4.12	.03	SD=6.94 (2 – 22) M=40.30;	SD=2.30 (0 – 7) M=33.25;	SD=6.70 (0 – 22) M=30.30;
STAI Trait	3.74	.03	SD=9.75 (25 – 61) M=34.20;	SD=6.54 (27 – 43) M=30.88;	SD=8.12 (23 – 51) M=27.90;
STAI State	1.87	.17	SD=10.87 (20 – 54)	SD=2.70 (28 – 34)	SD=4.86 (23 – 37)

M= mean.

SD= standard deviation.

Numbers within () are min-max.



emotions recognition [$F(2, 25) = 1.84, p = 0.17, \eta_p^2 = 0.12$]. Conversely, significant differences among groups were observed for disgust recognition [$F(2, 25) = 3.97, p < 0.05, \eta_p^2 = 0.24$]. Post-hoc comparisons showed significantly slower RT for controls than STM group ($p < 0.5$). No other differences were observed (see Figure 1b).

Empathy: Group differences were found in Personal distress subscale [$F(2, 25) = 7.88, p < 0.01, \eta_p^2 = 0.38$]. A post-hoc comparison (Tuckey HSD, MS = 13.53; df = 25.00) showed that both LTM and STM groups scored lower than controls ($p < 0.01$, for both). No other difference was observed (see Figure 1c).

Theory of mind (ToM): No group differences were observed [$F(2, 25) = 1.10, p = 0.34, \eta_p^2 = 0.08$] (see Figure 1d).

Interoception

No group effects [$F(2, 25) = 0.57, p = 0.57, \eta_p^2 = 0.04$] or condition \times group interaction [$F(6, 75) = 0.59, p = 0.72, \eta_p^2 = 0.04$] were observed. Thus, there were no significant differences in the ability to track their heartbeats (interoceptive conditions) or an external cued heartbeat (motor and feedback conditions), in any of the four conditions (See Figure 2). Only an expected [12] and irrelevant effect of condition was observed (see Additional file 2: Table S2).

Discussion

This is the first study assessing the influence of meditation practice both in cardiac interoception and in social cognition using a range of tasks. We selected a HBD

task that avoids the possible interference of external stimuli [11] previously reported [5,9].

No differences in EF or demographic variables were observed. Related to mood and affective scales, controls showed higher STM (depression) and LTM (anxiety-trait) scores. These results might reflect the possible influence of skills acquired during meditation practice (without considering its length), such as stress coping and emotional regulation abilities, which could help to deal with anxiety and depression situations. These skills might modulate mood perception as more euthymic and positive [20].

Regarding interoception, we replicated negative results previously reported [5,9]. Body awareness includes one internal (viscera and blood composition) and one external stream (taste, smell, pressure sensations and pain [21]). Consequently, cardiac sensations might be considered as a basic modality of visceral perception that relies mostly on internal drive (the heart being an internal organ), which is why it would be more difficult to gain conscious inspection. Respiration is unique among interoceptive signals as it involves external pressure information from the nose and chest, and it is susceptible of voluntary control and straightforward conscious perception. During meditation, attention is commonly directed towards breathing [5], where more consistent results have been shown [2,7]. These findings suggest that cardiac perception might not be the most suitable index to reflect meditation influence on interoception.

Few group differences were observed in social cognition domains. The lower accuracy in disgust recognition found




Figure 2 Heartbeat detection task (HBD). The Accuracy Index can vary between 0 and 1, with higher scores indicating better accuracy. No differences were found among groups in any condition. Vertical bars indicate standard deviation.

in LTM compared to controls might be related to their lower cardiac interoceptive sensitivity (given the common insular involvement for interoception and disgust recognition [22]). However, this is speculative because interoceptive differences were not significant among groups.

Both meditators' groups showed significantly lower empathy scores compared to controls only in the personal distress subscale, an index of emotional contagion by others' distress [23]. This is unsurprising since one of the aims of meditation is the regulation of responsiveness to stressors [24]. Finally, no difference in ToM was observed. Overall, despite the few differences reported, groups have similar social cognition performance suggesting that meditation practice in this study may not impact on these abilities.

Our study suffers from important limitations. First, the sample size should be increased to allow more informative analysis (i.e. correlations, multiple regressions) about the association among meditation, interoception and social cognition. However, it is worth highlighting that we reported preliminary data about interoception sensitivity measure with a novel method, and that previous research has employed similar sample size [9]. Second, further studies should cover a multidimensional interoceptive assessment (not only cardiac but also breathing, cardiac, visceral, etc.) and including both awareness and sensibility dimensions. Finally, groups' homogeneity should be guaranteed by measuring variables that might bias visceral perception such as physical state, volume stroke, blood pressure and contractility (Additional file 2: Table S2).

Conclusion

In conclusion, no influence of meditation practice in cardiac interoception and related social cognition measures

was observed. Based on the existence of diverse interoceptive signals, a more extensive assessment of each visceral source (other than cardiac one) may be necessary to disentangle the influence of meditation on interoceptive sensitivity.

Additional files

Additional file 1: Methods. In this Additional file 1 we provide a supplementary and detailed description of the materials and methods used in the study

Additional file 2: HBD additional results. In this Additional file 2 we provide a supplementary description of others interoceptive results.

Abbreviations

HBD: Heartbeat detection task; LTP: Long term practitioners; STM: Short term meditators; MBSR: Mindfulness-based stress reduction; ToM: Theory of mind; EF: Executive functions; BDI: Beck's depression inventory; STAI: State trait anxiety inventory; IFS: INECO frontal screening; IRI: Interpersonal reactivity index.

Competing interests

All the authors declare that they have no competing interests with respect to this study or its publication.

Authors' contributions

MM and LS collected the data, statistically analyzed the data and wrote the first draft of the manuscript. BC was involved in the study conception and design, writing the protocol and contributed to the drafting of the manuscript. MR contributed in collecting the data and revising the final version of the manuscript. CG contributed to writing the final version of the manuscript. ACJ and MS contributed to revising the final version of the manuscript. FM contributed to revising the final version of the manuscript. AI is the head of our laboratory, was involved in the study conception and design and contributed to writing the final version of the manuscript. All authors read and approved the final manuscript.

Authors' information

Margherita Melloni, Lucas Sedeño as the first author.

Acknowledgements

We thank the "Asociación Mindfulness Argentina" for providing the experimental subjects and the place for examination made on this work. Specifically, we are grateful for Mrs Clara Badino and Mr Julio Laurindo predisposition to participate and advice on mediator's sample selection. We also thank all participants of this study. This research was partially supported by CONICET, INECO Foundation, CONICYT/FONDECYT Regular (1130920), FONCYT-PICT 2012-0412, FONCYT-PICT 2012-1309, and James McDonnell Foundation Grants. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of those grants.

Author details

¹Laboratory of Experimental Psychology and Neuroscience (LPEN), INECO (Institute of Cognitive Neurology) and Institute of Neuroscience, Favaloro, Favaloro University, C1078AAI, Buenos Aires, Argentina. ²UDP-INECO Foundation Core on Neuroscience (UIFCoN), Diego Portales University, Santiago, Chile. ³Medical Research Council Cognition and Brain Sciences Unit, Cambridge, CB2 7EF, UK. ⁴Physics Department, Laboratory of Integrative Neuroscience, FCEyN UBA and IFIBA, Conicet, Pabellón 1, Ciudad Universitaria, 1428 Buenos Aires, Argentina. ⁵National Scientific and Technical Research Council (CONICET), Buenos Aires, Argentina. ⁶Universidad Torcuato Di Tella, Almirante Juan Saenz Valiente 1010, Buenos Aires, C1428BJJ, Argentina. ⁷Australian Research Council (ACR) Centre of Excellence in Cognition and its Disorders, New South Wales, Australia.

Received: 2 August 2013 Accepted: 17 December 2013

Published: 23 December 2013

References

1. Cameron OG: **Interoception: the inside story—a model for psychosomatic processes.** *Psychosom Med* 2001, **63**:697–710.
2. Craig A: **How do you feel? Interoception: the sense of the physiological condition of the body.** *Nat Rev Neurosci* 2002, **3**:655–666.
3. Barrett LF, Quigley KS, Bliss-Moreau E, Aronson KR: **Interoceptive sensitivity and self-reports of emotional experience.** *Journal Personality Social Psychology* 2004, **87**:684–697.
4. Lamm C, Singer T: **The role of anterior insular cortex in social emotions.** *Brain Struct Funct* 2010, **214**:579–591.
5. Khalsa SS, Rudrauf D, Damasio AR, Davidson RJ, Lutz A, Tranel D: **Interoceptive awareness in experienced meditators.** *Psychophysiology* 2008, **45**:671–677.
6. Hölzel B, Lazar S, Gard T, Schuman-Olivier Z, Vago D, Ott U: **How Does Mindfulness Meditation Work? Proposing Mechanisms of Action From a Conceptual and Neural Perspective.** *Perspect Psychol Sci* 2011, **6**:537–559.
7. Farb NA, Segal ZV, Anderson AK: **Mindfulness meditation training alters cortical representations of interoceptive attention.** *Social Cognitive and Affective Neurosci* 2013, **8**:15–26.
8. Vago DR, Silbersweig DA: **Self-awareness, self-regulation, and self-transcendence (S-ART): a framework for understanding the neurobiological mechanisms of mindfulness.** *Front Hum Neurosci* 2012, **6**:296.
9. Nielsen L, Kaszniak AW: **Awareness of subtle emotional feelings: a comparison of long-term meditators and nonmeditators.** *Emotion* 2006, **6**:392–405.
10. Whitehead WE, Drescher V, Heiman P: **Relation of heart rate control to heartbeat perception.** *Biofeedback Self Regul* 1977, **2**:371–392.
11. Pennebaker JW: *The psychology of physical symptoms.* New York: Springer; 1982.
12. Couto B, Salles A, Sedeño L, Peradejordi M, Bartfeld P, Canales-Johnson A, Dos Santos YV, Huepe D, Bekinschtein T, Sigman M, et al: **The man who feels two hearts: the different pathways of interoception.** *Social Cognitive and Affective Neurosci* 2013. published on line.
13. Brass M, Haggard P: **The hidden side of intentional action: the role of the anterior insular cortex.** *Brain Struct Funct* 2010, **214**:603–610.
14. Dunn BD, Galton HC, Morgan R, Evans D, Oliver C, Meyer M, Cusack R, Lawrence AD, Dalgleish T: **Listening to your heart. How interoception shapes emotion experience and intuitive decision making.** *Psychol Sci* 2010, **21**:1835–1844.
15. Pessoa L: **On the relationship between emotion and cognition.** *Nat Rev Neurosci* 2008, **9**:148–158.
16. Ahmed FS, Stephen Miller L: **Executive function mechanisms of theory of mind.** *J Autism and Developmental Disorders* 2011, **41**:667–678.
17. Singer T, Lamm C: **The social neuroscience of empathy.** *Ann N Y Acad Sci* 2009, **1156**:81–96.
18. Rouse CH, Jones GE, Jones KR: **The effect of body composition and gender on cardiac awareness.** *Psychophysiology* 1988, **25**:400–407.
19. Torralva T, Roca M, Gleichgercht E, Lopez P, Manes F: **INECO Frontal Screening (IFS): a brief, sensitive, and specific tool to assess executive functions in dementia.** *J The International Neuropsychological Society: JINS* 2009, **15**:777–786.
20. Britton WB, Shahar B, Szepesnwi O, Jacobs WJ: **Mindfulness-based cognitive therapy improves emotional reactivity to social stress: results from a randomized controlled trial.** *Behav Ther* 2012, **43**:365–380.
21. Cameron OG: *Visceral Sensory Neuroscience: Interoception.* New York: Oxford University Press; 2002.
22. Wicker B, Keysers C, Plailly J, Royet JP, Gallese V, Rizzolatti G: **Both of us disgusted in My insula: the common neural basis of seeing and feeling disgust.** *Neuron* 2003, **40**:655–664.
23. Davis M: **Measuring individual differences in empathy: evidence for a multidimensional approach.** *J Personality and Social Psychology* 1983, **44**:113–126.
24. Kabat-Zinn J, Nhat Hanh T: *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness.* New York: Bantam Dell; 1990.

doi:10.1186/1744-9081-9-47

Cite this article as: Melloni et al.: Preliminary evidence about the effects of meditation on interoceptive sensitivity and social cognition. *Behavioral and Brain Functions* 2013 9:47.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

