



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



SARS-CoV-2 attachment to host cells is possibly mediated via RGD-integrin interaction in a calcium-dependent manner and suggests pulmonary EDTA chelation therapy as a novel treatment for COVID 19

Tikam Chand Dakal

Genome and Computational Biology Lab, Department of Biotechnology, Mohanlal Sukhadia University, Udaipur 313001, Rajasthan, India

ABSTRACT

SARS-CoV-2 is a highly contagious virus that has caused serious health crisis world-wide resulting into a pandemic situation. As per the literature, the SARS-CoV-2 is known to exploit human ACE2 receptors (similar to previous SARS-CoV-1) for gaining entry into the host cell for invasion, infection, multiplication and pathogenesis. However, considering the higher infectivity of SARS-CoV-2 along with the complex etiology and pathophysiological outcomes seen in COVID-19 patients, it seems that there may be an alternate receptor for SARS-CoV-2. I performed comparative protein sequence analysis, database based gene expression profiling, bioinformatics based molecular docking using authentic tools and techniques for unveiling the molecular basis of high infectivity of SARS-CoV-2 as compared to previous known coronaviruses. My study revealed that SARS-CoV-2 (previously known as 2019-nCoV) harbors a RGD motif in its receptor binding domain (RBD) and the motif is absent in all other previously known SARS-CoVs. The RGD motif is well known for its role in cell-attachment and cell-adhesion. My hypothesis is that the SARS-CoV-2 may be (via RGD) exploiting integrins, that have high expression in lungs and all other vital organs, for invading host cells. However, an experimental verification is required. The expression of ACE2, which is a known receptor for SARS-CoV-2, was found to be negligible in lungs. I assume that higher infectivity of SARS-CoV-2 could be due to this RGD-integrin mediated acquired cell-adhesive property. Gene expression profiling revealed that expression of integrins is significantly high in lung cells, in particular $\alpha v \beta 6$, $\alpha 5 \beta 1$, $\alpha v \beta 8$ and an ECM protein, ICAM1. The molecular docking experiment showed the RBD of spike protein binds with integrins precisely at RGD motif in a similar manner as a synthetic RGD peptide binds to integrins as found by other researchers. SARS-CoV-2 spike protein has a number of phosphorylation sites that can induce cAMP, PKC, Tyr signaling pathways. These pathways either activate calcium ion channels or get activated by calcium. In fact, integrins have calcium & metal binding sites that were predicted around and in vicinity of RGD-integrin docking site in our analysis which suggests that RGD-integrins interaction possibly occurs in calcium-dependent manner. The higher expression of integrins in lungs along with their previously known high binding affinity ($\sim K_D = 4.0$ nM) for virus RGD motif could serve as a possible explanation for high infectivity of SARS-CoV-2. On the contrary, human ACE2 has lower expression in lungs and its high binding affinity ($\sim K_D = 15$ nM) for spike RBD alone could not manifest significant virus-host attachment. This suggests that besides human ACE2, an additional or alternate receptor for SARS-CoV-2 is likely to exist. A highly relevant evidence never reported earlier which corroborate in favor of RGD-integrins mediated virus-host attachment is an unleashed cytokine storm which causes due to activation of TNF- α and IL-6 activation; and integrins role in their activation is also well established. Altogether, the current study has highlighted possible role of calcium and other divalent ions in RGD-integrins interaction for virus invasion into host cells and suggested that lowering divalent ion in lungs could avert virus-host cells attachment.

1. Introduction

Ever since the recent emergence of novel coronavirus (SARS-CoV-2, earlier known as 2019-nCoV) in the Wuhan city of China and its subsequent transmission in other countries has resulted into serious health crisis as well as breakdown of socio-economic development. Scientific community from all over the world is industriously engaged in and committed to find a potent therapeutic solution for the treatment of COVID-19. As of 10th April, 1,521,252 confirmed cases and 92,798 deaths were recorded as a cumulative data from different parts of the world (WHO Situation Report no. 81 available at who.int accessed on 12-04-2020). At the onset of the disease, the infected symptomatic

patients experience hyperthermia, pharyngeal congestion, cough, and anosmia (in some cases); however, as disease progress more than fifty percent patients develop severe labored breathing, clinically referred to as dyspnoea or tachypnoea (Huang et al., 2020; Qiu et al., 2020; Wang et al., 2020a, 2020b, 2020c). Advance stages are characterized by severe pulmonary inflammation, fibrosis and obstructions of the bronchioles resulting in a pneumonia-like pathophysiological condition (Huang et al., 2020; Qiu et al., 2020; Wang et al., 2020a, 2020b, 2020c). In both symptomatic and asymptomatic COVID-19 patients, SARS-CoV-2 manages to cause significant damages to multiple organs before any patients could realize they are infected with SARS-CoV-2. This is because neither any neurological indications nor any signs of heart, kidney and liver

E-mail address: tc.dakal@mlsu.ac.in.

<https://doi.org/10.1016/j.imbio.2020.152021>

Received 21 April 2020; Accepted 18 October 2020

Available online 5 November 2020

0171-2985/© 2020 Elsevier GmbH. All rights reserved.

failure are seen at the onset of disease in these patients (Qiu et al., 2020).

As of today, there is no specific antiviral therapy available in any form, either vaccine or drug or others, to combat COVID-19 and its infection. The neutralizing antibodies that were previously tested and found successful against SARS-CoV-1 have displayed inappreciable cross-reactivity against SARS-CoV-2. For some antibodies (such as CR3022) that could bind to SARS-CoV-2, their neutralizing efficacy has not been tested and appreciated yet (Tian et al., 2020; Yuan et al., 2020; Wrapp et al., 2020). It has been asserted that the binding sites for these monoclonal antibodies on SARS-CoV-2 are vulnerable and it is only an assumption that antibodies that could bind strongly would possibly neutralize the SARS-CoV-2 also. Moreover, cross-reacting neutralizing antibodies are also doubted for their ability to confer prolonged protection against SARS-CoV-2. So far, we have no other choice than to use previously tested drugs or to solely oblige known tactics as precautionary measures against COVID-19. Recently, some researchers suggested the use of combination of remdesivir (a broad-range antiviral drug) and chloroquine for effective control of SARS-CoV-2 infection under *in vitro* condition (Devaux et al., 2020; Wang et al., 2020a, 2020b, 2020c). Similarly, hydroxychloroquine and azithromycin has also been referred to as a potent therapeutic weapon against the SARS-CoV-2 virus (Colson et al., 2020). Drug repurposing approach, which involved a screening of a thousand of molecules showed that HIV protease inhibitors, RNA-dependent RNA polymerase (RdRp) inhibitors and some other inhibitors and agonists such as methisazone (an inhibitor of protein synthesis), CGP42112A (an angiotensin AT2 receptor agonist) and ABT450 (an inhibitor of the non-structural protein 3-4A) could become promising treatment options for COVID-19 (Gordon et al., 2020; Li et al., 2020; Shah et al., 2020). Lu (2020) suggested some treatment options for COVID-19 that include use of nucleoside analogues, neuraminidase inhibitors, lopinavir or ritonavir, remdesivir, 3TC/TDF/EFV monotherapy or combination therapy (DNA polymerase inhibitors), anti-inflammatory or immune-suppressive drugs, just to name a few. Besides this, some traditional Chinese medicine, for instance, Shu-FengJieDu and Lianhuaqingwen capsules could also be useful (Lu, 2020). Using virtual screening (Kandeel and Al-Nazawi, 2020), epigenetic dysregulation (Sawalha et al., 2020), protein-protein interaction mapping (Cava et al., 2020), integrated network pharmacological approach (Wang et al., 2020a, 2020b, 2020c), and similar such approaches, a number of other drugs have also been repurposed for COVID-19 treatment. However, most of the drugs are either in developmental stages or under clinical trials (<https://clinicaltrials.gov/>). In the midst of this pandemic situation, it is obvious that there exists no preventive therapy for highly contagious SARS-CoV-2 and this is strikingly alarming to all of us.

A number of studies demonstrated the key role of human ACE2 in virus attachment to host cells (Wan et al., 2020; Zhao et al., 2020). The three-dimensional crystal structure of SARS-CoV-2 spike receptor binding domain complexed with its receptor, human ACE2 (Angiotensin converting enzyme 2), has been already solved (Chen et al., 2020). All other structural, functional and antigenicity related information related to SARS-CoV-2 are also available (Walls et al., 2020). Based on biophysical data, it has been demonstrated that the human ACE2 binds to SARS-CoV-2 with greater affinity than SARS-CoV-1 (Wrapp et al., 2020). However, owing to their low copy number (protein expression) (Chen et al., 2020), their high affinity (for SARS-CoV-2 than SARS-CoV-1) alone could not manifest reasonable virus-host cell attachment, at least in lung cells. When this manuscript study was under progress, Sigrist and coauthors (2020) showed that SARS-CoV-2 harbors a RGD motif and integrins (that display high affinity for RGD motifs) may be involved in facilitating virus entry into host cells. However, the study did not explain the full mechanistic state of affairs involved in RGD-integrins interaction and virus entry into the host cells. I also found RGD motif in the spike receptor binding domain of SARS-CoV-2 and studied mechanistic basis of RGD-integrin (and other ECM protein such as ICAM1) mediated virus invasion into host cells. This study is the first

study to present striking evidence (substantiated by existing facts in literature) favoring the role of calcium and other divalent ions (magnesium, manganese etc.) in RGD-integrins mediated virus attachment with the host cells for and that lowering the concentration of calcium and other divalent ions in lungs could be a possible mechanism to avert SARS-CoV-2 infection and invasion. Herein, I did comparative protein sequence analysis, motif scanning, gene expression data analysis and bioinformatics based molecular docking using most trustworthy tools and techniques. The results generated in this study indicated and underscored the key role of calcium and other divalent ions in mediating virus-host cell attachment for invasion into human lung cells possibly via integrins and other cell adhesion proteins present on the host cell surface. I designed and suggested novel pulmonary EDTA chelation therapy as a technical simple (requires nebulizer/inhaler/nasal sprays that are available in all hospitals and clinics), quick (1-3hr a day, 2-3 times a week), safe (EDTA/EGTA are FDA approved for calcium chelation), affordable (estimated cost approx. 50-100 USD) and efficient (based of efficacy data from literature) treatment therapy for COVID-19 patients.

2. Materials & methods

2.1. Sequence retrieval

The coronavirus related nucleotide and protein sequences were retrieved from GenBank (<https://www.ncbi.nlm.nih.gov/>), UniProtKB (<https://www.uniprot.org>) and SARS Coronavirus 2 data hub of the NCBI (<https://www.nih.gov/coronavirus>).

2.2. Sequence alignment

The protein sequence of SARS-CoV-1 and SARS-CoV-2 coronaviruses were subjected to pair-wise sequence alignment using Clustal Omega using default setting (<https://www.ebi.ac.uk/Tools/msa/clustalo/>) (Madeira et al., 2019).

2.3. Motif scanning and prediction

The Pfam, Prosite and HAMAP profiles of the S protein from the coronavirus was ascertained using MyHits Motif Scan (SIB, Switzerland) (https://myhits.isb-sib.ch/cgi-bin/motif_scan). The putative N-linked glycosylation and other post-translation modification sites were predicted as Frequent Pattern in MyHits outputs (Pagni et al., 2007).

2.4. Gene expression profiling

The expression profile of ACE2 receptor protein and integrins in lungs was ascertained using Gene Expression Database at EBI (<https://www.ebi.ac.uk/gxa/home>) and the Human Protein Atlas (<https://www.proteinatlas.org/>) (Uhlén et al., 2015, 2019).

2.5. Calcium and magnesium binding sites in integrins

In order to understand the role of divalent ions in RGD-integrins mediated virus-host cells attachment. A composite calcium and magnesium binding-site in integrins were predicted using IonCom (<http://zhanglab.ccmb.med.umich.edu/IonCom/>), which used an integrated approach based on *ab initio* training and template-based transfers (Hu et al., 2016) and predicts calcium binding sites in a given protein by searching four or more oxygen atoms on protein surface arranged in a spherical manner. The input PDB file of integrins such as $\alpha 5\beta 1$ (PDB ID: 3VI3) and $\alpha v\beta 6$ (5FFG) were subjected for ion binding sites' prediction to specifically predict calcium and magnesium binds sites.

2.6. Spike RBD docking with integrins

The PDB file of the spike receptor binding domain (PDB ID: 6LZG) and integrins such as $\alpha 5\beta 1$ (PDB ID: 3VI3 and 3VI4) and $\alpha v\beta 6$ (PDB ID: 5FFG) were retrieved from Protein Data Bank (<http://www.rcsb.org/structure/>). The protein–protein docking was performed using HDock server which rely on template-based modeling and *ab initio* free docking using default parameters (hdock.phys.hust.edu.cn/). The docked structures were visualized and high resolution photographs were generated in Chimera 1.10.

2.7. Spike RBD docking with EDTA

The PDB file of the spike receptor binding domain was retrieved from Protein Data Bank (as mentioned above). The PDB file of the EDTA was online obtained and converted in Mol2 file in Chimera 1.10.1 version. The docking was done using Swissdock server of SIB (<http://www.swissdock.ch/docking>). Protein–ligand binding modes were clustered according to their rank based on average FullFitness output data (Grosdidier et al., 2007).

3. Results

3.1. N-terminus of spike protein is variable

Pairwise sequence alignment of SARS-CoV-1 and SARS-CoV-2 spike protein revealed that the N-terminus (especially upto 250 aa) of SARS-CoV-2 is highly divergent than that of the SARS-CoV-1 (Fig. 1). The S1 Glycoprotein (from 268 to 304 aa in SARS-CoV-1 and from 282 to 317 aa in SARS-CoV-2) domain which plays an important role in recognition of host cell receptor were found divergent at 10 amino acid positions. The spike receptor binding domain is of 253 amino acid residues in SARS-

CoV-1 and ranges from 317 to 569 aa; whereas, it is of 254 amino acid residues in SARS-CoV-2 and stretch from 330 to 583 aa. There were 49 mismatches and 1 insertion/deletion that render SARS-CoV-1 and SARS-CoV-2 approximately 80% sequence similarity. The S2 glycoprotein domain is a large 544 amino acid residues long region at the C-terminus of the spike protein which plays crucial role in virus fusion with host cells. This region has been predicted to be less divergent than S1 glycoprotein and RBD. There were 49 amino acids substitution which amount for approximately 9% sequence divergence between SARS-CoV-1 and SARS-CoV-2. The cysteine rich region at the C-terminus part of the SARS-CoV-2 spike protein has an additional Cys residue (actually Ala > Cys) at 1247 position (corresponds to Ala residue at 1229 position in SARS-CoV-1 spike protein in alignment). This region is also involved in virus-induced membrane fusion (Chang et al., 2000). The presence extra Cys residue results in the formation of a C_{X6}CC motif, which is usually found in envelop proteins of some viruses, for instance Ebola virus and murine leukemia virus, and is required for virus fusion with host cell (Johnston and Radke, 2000; Lee and Saphire, 2009).

I also found two insertion sequences with the stretch HVSGTNGTKRFD⁶⁹⁻⁸⁰ and RFQTLALHRSYLTPGDSSSGWTAG²³⁶⁻²⁶¹ at the N-terminus region of spike protein and these were seen as discontinuous in the amino acid sequence as predicted by the pair-wise sequence alignment. However, considering that these inserts are very short and appeared in the hypervariable region of viral spike protein, the most likely assumption for their origin is that they might have arisen naturally. Besides this, the receptor binding domain of the spike protein in SARS-CoV-2 has a stretch of sequence (TEIYQAGSTPCNGVEGF⁴⁷⁰⁻⁴⁸⁶) showing mismatch with SARS-CoV-1.



Fig. 1. Pair-wise sequence alignment of SARS-CoV-1 and SARS-CoV-2 spike protein RBD sequence using Clustal Omega.

3.2. Increased number of phosphorylation sites in SARS-CoV-2 led to increased viral infection and pathogenesis

The number and position of putative N-linked glycosylation sites, post-translation modification sites that can induce cAMP, CK2, Tyr, PKC signaling pathways and Myristyl sites were predicted to be varying in SARS-CoV-1 and SARS-CoV-2 (Table 1). Although both coronaviruses have equal number of N-linked glycosylation sites but the position of sites differs (Table 1). Furthermore, phosphorylation sites of all types (except PKC phosphorylation) were predicted to be in more number in SARS-CoV-2 spike protein than in SARS-CoV-1. The phosphorylation of viral proteins is actuated by host cells' kinases has a significant impact on viral infection, replication, multiplication and cytotoxicity (Bretaña et al., 2012; Keating and Striker, 2012; Keck et al., 2015), I assume that the increased number of phosphorylation sites in SARS-CoV-2 spike protein could be one of the major reasons for its higher infectivity as observed in case of COVID-19. This can be ascertained using *in-vitro* experiments. These pathways either activate calcium ion channels or get activated by calcium. This suggested that virus might be directly involved in regulating these calcium ion channels or might be dependent upon presence of Ca²⁺ ion or other divalent ions for attachment with host cells.

3.3. SARS-CoV-2 spike receptor binding domain harbors a RGD motif which is absent in SARS-CoV-1

The spike protein sequence of SARS-CoV-1 and SARS-CoV-2 were subjected to motif scanning and prediction using MyHits Motif Scan at SIB server. A number of motifs were predicted in the spike protein sequence such as RGD (from 403 to 405 aa in receptor binding domain of SARS-CoV-2) (Table 2). The RGD motif (K403R substitution) was predicted in spike receptor binding domain of SARS-CoV-2 and the same was not predicted in the SARS-CoV-1 spike protein and other previous known coronaviruses. This motif was originally found in fibronectin, which is an extracellular matrix protein and this motif plays a major role in cell adhesion and attachment. A number of proteins are known to interact with these RGD motifs. One such group of proteins are integrins that have strong affinity for RGD motif ($-K_D = 4.0$ nM) and they employ this motif to efficiently mediate cell adhesion. Eight out of approximately twenty known integrins recognize and exploit RGD motifs for cell adhesion (Ruoslahti, 1996; Teoh et al., 2016). Presence of motif RGD in

Table 1

Prediction of putative N-linked glycosylation and other PTM sites using MyHits Motif Scan of SIB. Top scored N-linked Glycosylation sites are represented in bold.

S. No.	Pattern	Position	
		SARS-CoV-1	SARS-CoV-2
1.	ASN_Glycosylation	29, 65, 73, 109, 118, 119, 158, 227, 269, 318, 330, 357, 589, 602, 691, 699, 783, 1056, 1080, 1116, 1140, 1155, 1176	17, 61, 74, 122, 149, 165, 234, 282, 331, 343, 603, 616, 657, 709, 717, 801, 1074, 1098, 1134, 1158, 1173, 1194
2.	CAMP_Phospho_Site	343	356, 528
3.	CK2_Phospho_Site	12, 20, 271, 561, 644, 716, 798, 809, 964, 1118, 1129, 1178	50, 108, 151, 221, 250, 284, 734, 816, 827, 982, 1136, 1147, 1160, 1196
4.	Myristyl	86, 104, 225, 246, 298, 368, 418, 531, 587, 634, 653, 682, 739, 862, 871, 890, 953, 1113, 1153, 1228	72, 89, 184, 232, 311, 381, 431, 446, 476, 545, 601, 648, 667, 700, 757, 880, 889, 908, 971, 1093, 1131, 1171, 1246
5.	PKC_Phospho_Site	36, 92, 173, 215, 289, 363, 541, 561, 670, 775, 795, 980, 1019, 1087, 1129	19, 76, 95, 302, 376, 415, 555, 632, 680, 813, 998, 1037, 1105, 1147
6.	Tyr_Phospho_Site	188, 715	417, 733

Table 2

Artificial neural network based predicted motifs in the protein sequence of SARS-CoV-1 and SARS-CoV-2.

S. No.	Motifs	Position		Motif's Role/Function
		HCoV-1	HCoV-2	
1	S1 Glycoprotein	268–304	519–592	Viral attachment
2	Potato Inhibitor 1	309–320	Not predicted	NA
3	Spike Receptor Binding domain	317–569	330–583	Attachment with host cells
4	RGD	Not Predicted	403–405	Possibly attachment with host cells
5	FMRP	437–447	Not predicted	NA
6	S2 Glycoprotein	641–1247; 694–1237	684–1265; 712–1255	Fusion of viral and cell membrane as well as fusion of infected cells with adjoining cells.
7	Borrelia Repeat	662–679	Not Predicted	NA
8	DUF16	892–971	Not Predicted	Protein of unknown function
9	Fusion Glycoprotein F0	910–935	Not Predicted	Induces fusion of viral and cellular membranes leading to delivery of the nucleocapsid into the cytoplasm
10	BIG1	Not predicted	1–3	NA
11	Cysteine-rich Domain	1217–1236	1235–1254	Membrane fusion

SARS-CoV-2 spike receptor binding domain is a strikingly alarming as this motif is expected to facilitate even stronger attachment between virus and human target cells, and thus, rendering human cells more susceptible and vulnerable to infection. Motif scan also predicted fusion glycoprotein F0 (from 910 to 935 aa in S2 glycoprotein of SARS-CoV-1) and FMRFamide (from 437 to 447 aa in receptor binding domain of SARS-CoV-1). The absence of fusion glycoprotein F0 in SARS-CoV-2 may also have potential deleterious effect because this motif acts as a potent antigen that are targeted by neutralizing antibodies induced by virus infection as a part of humoral immune response and adaptive immunity (Prabakaran et al., 2006) and its absence in SARS-CoV-2 spike protein explains why monoclonal antibodies raised against SARS-CoV-1 could not show cross-reactivity with SARS-CoV-2 spike protein. This also accounts for the reason why an appropriate and adequate immune response could not mount in COVID-19 patients, especially in aged and immunocompromised COVID-19 patients, which eventually led to patients' death. The FMRFamide (Phe-Met-Arg-Phe-amide) like peptides (also known as FLPs) (predicted to be absent in SARS-CoV-2), are one particular group of Arg-Phe-amides (RFamides), are known to play a central role in parasite neuromuscular biology (Peymen et al., 2014); however, their role in pathogenesis of SARS-CoV-2 cannot be ascertained yet and same for other predicted motifs (Table 2).

3.4. ACE2 expression is high in digestive route while integrins expression is high in lung cells

Presence of RGD motif in the spike receptor binding domain of the SARS-CoV-2 and the fact that integrins display a strong affinity for proteins harboring RGD motifs (Liu, 2009) suggested and indicated that virus possibly enters into host cells via integrins receptor (Ruoslahti, 1988; Van Agthoven et al., 2014). In order to assess this, I analyzed the gene expression data for all eight integrins at Human Protein Atlas (<https://www.proteinatlas.org/>) (Uhlén et al., 2015, 2019) and found that expression of three integrins (α v β 6, α 5 β 1, α 8 β 1) is significantly high in lung cells and other vital organs of human body that have been found to be affected by SARS-CoV-2 infection (Fig. 2A–C). These three




Fig. 2. Gene expression profile of RGD binding integrins $\alpha\beta6$ (A), $\alpha5\beta1$ (B), $\alpha8\beta1$ (C) and human ICAM1 (D) in lungs as obtained and plotted at the Human Protein Atlas. (<https://www.proteinatlas.org/>). The bar shows expression level of respective protein in different human tissues and organs. The expression peaks that correspond to lungs have been shown using red arrow to ease visualization and verification of data. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

integrins were used for molecular docking experiments. I also found that expression of an ECM protein ICAM1 is significantly higher (even higher than integrins) in lungs (Fig. 2D) suggesting that SARS-CoV-2 spike protein might be using these integrins and/or ECM proteins such as ICAM1 (or others) for invading host cells. On the contrary, the ACE2 RNA is detected in very low quantity (almost unidentifiable) in lungs and detectable expression peaks were detected in other organs such as duodenum, small intestine, colon, gallbladder, kidney, testis, and heart muscles (Fig. 3). In congruence with this, Xu and coauthors (2020a, 2020b) found high expression of ACE2 in oral mucosa and Zhang and coauthors (2020) demonstrated digestive route as a potential mechanism of ACE2 mediated virus infection based on the single-cell transcriptomic analysis. On the contrary, Chen and coauthors (2020) observed relatively low levels of ACE2 mRNA expression in lungs which further supports my claims.

3.5. SARS-CoV-2 spike RBD interacts with integrins and ICAM1 precisely at RGD motif

The RGD motif is the minimal indispensable requirement for integrins to bind with any viral protein which virus can use for attachment with host cells (Hussein et al., 2015). Nagae and coauthors (2012) demonstrated that synthetic RGD tripeptide binds to the integrin $\alpha5\beta1$ (PDB ID: 3V14). In order to obtain molecular evidences for integrins mediated SARS-CoV-2 entry into lung cells, I docked the viral spike RBD domain (PDB ID: 6LZG) onto the structure of integrin $\alpha5\beta1$ (PDB ID:

3V13 without synthetic RGD tripeptide) (Nagae et al., 2012) and integrin $\alpha\beta6$ (PDB ID: 5FFG) (Dong et al., 2017) and found that spike RBD binds with integrin precisely via the RGD motif as expected (Fig. 4A). The predicted docking energy score was -302.33 (ΔG) and the ligand RMSD score was 71.80 Å for spike RBD-integrin $\alpha\beta6$ docking; and for the spike RBD-integrin $\alpha5\beta1$, the scores were respectively -315.45 and 37.4 Å. I focused on only $\alpha\beta6$, $\alpha5\beta1$, and $\alpha8\beta1$ as they have been found to be expressing abundantly in lungs tissues/cells. All the other integrins are also known to bind to RGD peptides with different degree of affinity and I did in-silico work on those that show higher affinity for RGD. I checked the binding of other integrins with RGD and I found that the stability of integrin-RGD is low (data not shown). I found that the spike RGD interacts with loop turns AAG¹⁹⁹⁻²⁰¹, GSYFWQ²²⁵⁻²³⁰ and RQASSIYDDA²⁶⁰⁻²⁷⁰ of the $\alpha5$ chain of the integrin $\alpha5\beta1$ (Nagae et al., 2012) (Fig. 4B). The latter (RQASSIYDDA²⁶⁰⁻²⁷⁰) has also been predicted as one of the calcium binding sites as per the UniProt KD database (ITGA5; UniProt ID: P06756) and our IonCon analysis. Presence of Ca^{+2} ion binding sites around and in vicinity of RGD-integrins docking site suggest that RGD-integrin interaction is indeed calcium-dependent (Fig. 4B). ICAM1 is also known to bind integrins and this mechanism is exploited by some viruses (such as Rhinoviruses) to gain entry into respiratory system for pathogenesis (Bella et al., 1998).




Fig. 3. Gene expression profile of ACE2 in lungs as obtained and plotted at Gene Expression Database at EBI (<https://www.ebi.ac.uk/gxa/home>) (A) and the Human Protein Atlas (<https://www.proteinatlas.org/>) (B). The expression peak that corresponds to lungs have been shown using red arrow to ease visualization and verification of data. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 4. Molecular docking of viral spike RBD domain onto the structure of $\alpha 5\beta 1$ (PDB ID: 3VI3) and $\alpha v\beta 6$ (PDB ID: 5FFG) showing interaction via the RGD motif present in the spike RBD domain (A). The molecular docking of viral spike RBD domain onto the structure of $\alpha 5\beta 1$ (PDB ID: 3VI3) showed that the spike RBD is interacting with the $\alpha 5\beta 1$ through its RGD motif in the same way (B, left panel) as synthetic RGD peptide binds and interact to $\alpha 5\beta 1$ (PDB ID: 3VI4) (B, right panel) as already known in literature (Nagae et al., 2012). The docking was done using HDock server (hdock.phys.hust.edu.cn/). Spike protein is shown in red and synthetic RGD peptide is shown in green. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

3.6. Molecular docking suggests EDTA chelation therapy as a novel treatment for COVID-19

The RGD motifs are well known in the field of cell biology, cell therapy and tissue engineering because of their remarkable cell-adhesive property. Extracellular matrix proteins, such as fibronectin and laminin, possess RGD motif and are frequently coated onto the surface of biomaterials (petri-dishes, T-flasks) for facilitating human cells adhesion onto the surface of the biomaterials. Bivalent ions such as Ca^{+2} & Mg^{+2} play indispensable role in promoting such cell adhesion and EDTA is used as a chelating agent for disaggregating Ca^{+2} mediated cell bondage.

Two features in SARS-CoV-2 spike protein suggest virus attachment with host cells via integrins in a calcium-dependent manner: 1) presence of RGD motif in receptor binding domain of SARS-CoV-2 and 2) presence of a number of phosphorylation sites for key pathways that regulates calcium ion channels, and secondly, Ca^{+2} signaling. Similarly, integrins possess calcium and magnesium binding sites (as predicted by IonCon) for interaction with RGD motif containing proteins, and the same in current context are spike protein of SARS-CoV-2. Besides this, integrins are also known to regulate cell adhesion by altering intracellular calcium dynamics and have involvement in regulation of ion channels and calcium signaling to exert pathophysiological impact in other diseases (Sjaastad and Nelson, 1997; Arcangeli and Becchetti, 2010). Altogether, it appears that SARS-CoV-2 enters into human cells via interaction of their spike receptor RGD motif with integrins in a Ca^{+2} dependent manner.

Altogether, the current study highlighted the possible role of calcium and other divalent ions in RGD-integrins binding for virus invasion and suggested that lowering divalent ion in lungs could avert virus-host cells binding. This can be accomplished by EDTA or EGTA, both are well known chelating agents I strongly believe that phenotypic remodeling and regulation of Ca^{+2} ions (Berridge, 2012) by pulmonary EDTA chelation can discourage the SARS-CoV-2 virus attachment with human lung cells which will led to attainment of a state of protection against this virus. Finally, EDTA chelation therapy targeting lungs (as in discussion part) holds immense potential in successful management and treatment of COVID-19 patients.

3.7. RGD-Integrins binding cause activation of downstream integrin-mediated cell signaling (GO:0007229)

Further, it seems that RGD-integrins binding would relay or activate downstream integrins mediated cell signaling pathway (GO:0007229) which comprise of 106 human genes related to different biological processes and cellular pathways (<http://amigo.geneontology.org/>; Data not shown). The experimental evidences collected from UniProt KB databases show that integrins play key role in viral attachment and TGF- β and IL-1 mediated signaling. Other signaling pathways that are known to be regulated by integrins upon interaction with different viruses are FGF1 signaling, FGF2 signaling, NRG1-ERBB signaling, CX3CL1 signaling and CD40-CD40LG signaling (Table 3).

4. Discussion

In any disease or pathological condition where integrin expression would be high, I can expect patient to be highly vulnerable to COVID-19. Integrin expression has been found high in case of cancer (Bianconi et al., 2016) and diabetes (Roth et al., 1993; Miller et al., 2014) and I assume the susceptibility for SARS-CoV-2 infection and COVID-19 would be high in patients suffering from any diseases in which integrin expression is high. The immune-diagnosis and profiling of COVID-19 patients showed elevated levels of pro-inflammatory cytokines including IL-6 and IL-1 β in serum (Huang et al., 2020; Tan et al., 2020; Qin et al., 2020; Xu et al., 2020a, 2020b). They also exhibited an increased serum levels of other cytokine and chemokine molecules such

Table 3

RGD binding Integrins, their receptors and their role in virus attachment and cell signaling as sourced from UniProt KB database.

S. N.	Integrin	UniProt ID	Receptor	Virus Attachment	Function
1	α v β 3 ITGAV- ITGB3 CD61	P06756 P05106	CD40LG	Herpes virus 8	FGF1 signaling
			FGF1/FGF2	HHV-8	FGF2 signaling
			CX3CL1	Coxsackievirus A9	NRG1-ERBB signaling
			PLA2G2A	Hantaan virus	CX3CL1 signaling
			NRG1	Cytomegalovirus/	IL-1B signaling
			IGF2	HHV-5	Human metapneumovirus
			IL-1B	Human	IL-1B signaling
			FBN1	Human	IL-1B signaling
			CD40-LG	Human	CD40-CD40L signaling
			IGF2	parechovirus 1	IGF2 signaling
2	α v β 5 ITGAV- ITGB5	P06756 P18084	Fibronectin	Adenovirus type c	TGF- β 1 signaling
				Coxsackievirus A9	
				Coxsackievirus B1	
				Coxsackievirus A9	
				Coxsackievirus B1	
				Herpes simplex virus-1	
				HHV-1	
				Data not found	
3	α v β 6 ITGAV- ITGB6	P06756 P18564	Fibronectin	Coxsackievirus A9	TGF- β 1 activation
			Cytotactin	Coxsackievirus B1	Clathrin-mediated endocytosis
			TGF- β 1	Herpes simplex virus-1	
			FBN1	HHV-1	
				Data not found	
4	α v β 8 ITGAV- ITGB8	P06756 P26012	Fibronectin		TGF- β 1 activation
			TGF- β 1		
5	α 5 β 1 ITGA5- ITGB1 CD29	P08648 P05556	Fibrinogen	Cytomegalovirus/	CD40-CD40L signaling
			Fibronectin	HHV-5	IL1B signaling
			PLA2G2A	Epstein-Barr virus/HHV-4	IL1B signaling
			CD40LG	Human parvovirus B19	
			IL1B	Mammalian reovirus	
			FBN1	HHV-1	
				Human metapneumovirus	
6	α 8 β 1 ITGA8- ITGB1	P53708 P05556	TNC	Data not found	TGF- β 1 signaling
			FN1		
			SPP1		
			TGFB1/		
			TGFB3		
			VTN		
			NPNT		

as IL-2, IL-8, IL-17, G-CSF, GM-CSF, IP10, MCP1, MIP1 α (also known as CCL3) and TNF (Huang et al., 2020; Shi et al., 2020; Qin et al., 2020; Xu et al., 2020a, 2020b). Also, C-reactive protein and D-dimer are found to be abnormally high (Cao, 2020). High levels of pro-inflammatory cytokines may lead to tissue shedding and damage in lungs, heart, liver and kidney, i.e., multiple organ failure (Cao, 2020). The most evident pathophysiological outcome of SARS-CoV-2 infection is activation of immune cells and massive infiltration of neutrophils and macrophages that eventually led to “cytokine storm” resulting in severe lung inflammations (such as pulmonary edema and alveolar damages) and can be characterized by pneumonia-like symptoms and respiratory failure (Mehta et al., 2020). Herein, I underline another strong and convincing

evidence in favor of virus-host attachment through RGD-integrins, i.e., the cytokine storm, caused due to activation of pro-inflammatory cytokines such as TNF- α and IL-6 activation; and integrins role in their activation is also well established (Blobe et al., 2000; Annes et al., 2002; Lu et al., 2002; Sheppard, 2008; Hung et al., 2018). Integrin $\alpha\beta1$ regulates Ca^{+2} ion transient responses to IL-1 and TGF- β (Parekh et al., 2014). Integrins also regulates calcium mobilization in associated pathways such as Go-mediated cell signaling (Erb et al., 2001). Besides this, envelop proteins of virus, including previous coronaviruses, are Ca^{+2} channeling viroporins that exploit host cell signaling and are involved in virus entry, replication, immune evasion leading to activation of autophagy and inflammasome responses such as that of NLRP3 (Nieto-Torres et al., 2015; Hyser and Estes, 2015). In context of SARS-CoV2 and integrins attachment, what exact role does the dynamics (in terms of mobilization, concentration, frequency, timing) of intracellular and extracellular calcium play is still not clear; however, understanding this dynamics would greatly assist us in appropriately and adequately controlling the host cells' response to SARS-CoV-2 so as to counteract virus infection and invasion. However, it is apparently clear that high Ca^{+2} ion concentration in lungs is detrimental in case of SARS-CoV-2 infection.

A number of studies demonstrated the key role of human ACE2 in virus attachment to host cells (Wan et al., 2020; Zhao et al., 2020). Previous coronaviruses have also been found to enter into host cells using clathrin and caveolae-independent pathway (Inoue et al., 2007; Wang et al., 2008). Integrin $\alpha\beta6$ have also been implicated in mediating clathrin-dependent endocytosis (Table 3). Based on the biophysical data, it has been demonstrated that the human ACE2 binds to SARS-CoV-2 with greater affinity than SARS-CoV-1 (Wrapp et al., 2020). However, owing to their low copy number (gene expression), their high affinity (for SARS-CoV-2 than SARS-CoV-1) alone could not manifest reasonable virus-host cell attachment, at least in lung cells. On the other hand, the affinity of integrins for RGD motif of the spike RBD is considerably high. The binding affinity of ACE2 for whole spike RBD is $K_D = 15.2$ nM ($K_D =$ dissociation constant) (Wrapp et al., 2020) and on the other hand the binding affinity of integrins for RGD is close to $K_D = 4.0$ nM (Bernhagen et al., 2017, 2019; Ma et al., 2017). Since, the binding affinity has inverse relationship with K_D , I can rule out that virus spike protein would prefer binding to integrins rather than ACE2, at least in lung cells.

However, ACE2 expression has been found to be considerably high in the digestive tract from duodenum to rectum as well as in gall bladder and kidney. Therefore, ACE2 could be responsible for virus infection in these organs. This study presents striking evidence favoring the role of calcium and other divalent ions (magnesium, manganese etc.) in RGD-integrins mediated virus attachment with the host cells in lung cells for infection and invasion.

Chelating agents are long known for treating metal toxicity in humans in the context to several disease conditions such as coronary heart disease, neurotoxicity and arthritis etc. (Goyer et al., 1995; Bamonti et al., 2011; Ferrero, 2016; Fulgenzi and Ferrero, 2019). The EDTA is a well-known chelating agent and is a FDA approved drug for treatment of hypercalcemia conditions. In EDTA chelation therapy disodium EDTA (or similar sodium-EDTA derivative) is used as a chelating agent and is administered intravenously for the chelating divalent ions and the resulting chelates then come out in urine after passing through glomerulus filtration (Bamonti et al., 2011; Ferrero, 2016; Fulgenzi and Ferrero, 2019).

I have proposed EDTA chelation therapy as a potent anti-viral therapy against SARS-CoV-2. Pulmonary EDTA chelation therapy has already been executed clinically for asthmatic patients and is also known to increase the efficacy of nebulized bronchodilators (Asmus et al., 2001). Sodium salts of EDTA can be used for chelating Ca^{+2} ions as calcium-EDTA has higher stability than sodium-EDTA (Blaurock-Busch and Busch, 2014) and the resulting Ca-EDTA chelate would be soluble that can easily excrete out of body through glomerular filtration (Fig. 5). I have designed EDTA chelation therapy as a novel therapeutic strategy for safe, fast, cost-effective and efficient treatment of COVID-19 based on genomic understanding of SARS-CoV-2 virus spike protein. In EDTA chelation therapy, disodium EDTA will be administered in the body (preferably through nebulizer/inhaler/nasal spray or intravenously) so as to reach to its target organ, i.e., lungs as shown in Fig. 5. In normal disease cells, virus could make entry into host cells *via* binding of RGD motif (in their spike receptor binding domain) to the integrins receptor proteins present on the surface of human lung cells. Ca^{+2} ions (or other bivalent ions) would augment the binding of viral spike protein with host cells and subsequently facilitate viral invasion into cells for infection, replication and pathogenesis. On the contrary, using EDTA chelation therapy, Ca^{+2} or other bivalent ions can be chelated and can be




Fig. 5. Pulmonary EDTA chelation therapy which can be clinically executed through a nebulizer or inhaler to allow sodium-EDTA to pass into the lungs. The sodium-EDTA will chelates Ca^{+2} ions and other divalent ions making them unavailable for RGD-integrin mediated virus attachment to the host cells. A novel strategy for safe, technically simple, quick, cost-effective and efficient treatment of COVID 19 patients.

made unavailable to viral spike protein for attaching to host cells' integrins. In parallel, strict control over diet (or medication) for ensuring proper availability of Ca^{+2} in the body may also be required for successful phenotypic remodeling for EDTA chelation therapy. In this way, virus could not enter cells and a state of protection against SARS-CoV-2 and COVID-19 can be achieved (Fig. 5).

Besides ceasing the RGD-integrin mediated virus attachment to host cells, treatment with EDTA or EGTA is expected to cease SARS-CoV-2 viral infection in a multifaceted way. Firstly, I expect that chelating agents would disintegrate the viral membrane as Ca^{+2} forms major component of retrovirus membrane and are known to stabilize envelop proteins and others (Wunderlich and Sydow, 1982). Second, I also expect chelation of the metal cofactors within the enzyme active site of the SARS-CoV-2, as seen in HIV-1 IN enzyme (Hicks and Gulick, 2009). Third, EDTA is expected to lower metal-ion dependent reactive oxygen species (ROS) production and would also lipid peroxidation (Roussel et al., 2009). EDTA has also been shown to induce protective antioxidant and anti-inflammatory response in case of liver fibrosis (González-Cuevas et al., 2011). Finally, as COVID-19 patients exhibit excessive cytokine secretion (called as cytokine storm) (Mehta et al., 2020), use of EDTA is expected to render an anti-inflammatory and immunosuppressive effect which would be highly beneficial. As such there are plenty of studies that confirm EDTA safe use in chelation therapy with no potent adversity and fatality (Roussel et al., 2009). The EDTA concentrations greater than 0.6 mM may lead to reduced energy metabolism in lung cells (Asmus et al., 2001). Besides this, some complications such as hypocalcemia, proteinuria and renal insufficiency are known to associate with EDTA chelation therapy and I suggest that a strict monitoring protocol, comprising routine cytokine profiling, blood examination, urine test and CT scanning etc., must be practiced while treating patients with pulmonary EDTA chelation therapy.

5. Conclusion

In combination with the EDTA/EGTAbased pulmonary chelation therapy, modulation of integrins expression using integrins inhibitors or anti-integrin antibodies could also serve as a mechanism to treat COVID-19 patients (Henderson et al., 2013; Hatley et al., 2018; Sigrist et al., 2020). The expression of such integrins can be down regulated by modulation of intracellular redox ions such as superoxide anion ($\text{O}_2^{\cdot-}$) and H_2O_2 (Puchsaka et al., 2016). Besides this, some bioactive compounds/molecules such as Ouabain are also known to affect integrins expression in lung cells (Ninsontia and Chanvorachote, 2014). Application of synthetic RGD tripeptide or its derivatives (linear, cyclic etc.) can also specifically block integrins and could also serve as a possible mechanism to discourage virus-host attachment (Bernhagen et al., 2017, 2019; Ma et al., 2017). Besides this, PGNase treatment is also expected to be equally effective as the mass spectrometric studies revealed that all major glycan classes of human lung glycome can be released effectively by PNGase F (Jia et al., 2020). Since, spike protein of SARS-CoV-2 has a number of sites for cAMP, CK2, PKC and Tyr phosphorylation, I believe synergistic clinical application of other potential drugs such as kinase inhibitors can also be beneficial in COVID-19 treatment. Therapies specifically targeting TNF- α and IL-6 mediated signaling to down-regulate cytokines secretion seems to be a promising approach to deal with the situation of cytokine storm. In this regard, tocilizumab (anti-IL6 monoclonal antibody) alone or in combination with infliximab and Emodin are also worth testing for COVID-19 treatment (Cao, 2020). Besides this, I also believe that blocking the calcium ion channels (or others) and downregulation of integrin-mediated cell signaling (GO:0007229) pathways such as FGF1 signaling, FGF2 signaling, NRG1-ERBB signaling, CX3CL1 signaling and CD40-CD40LG signaling are also available as a potent immunotherapeutic solutions against COVID-19. Finally, in the era of multi-omics technologies, medicines/therapies based on comparative coronavirus genome analysis and understanding of individual pathophysiology could pave way towards successful

development of effective therapeutic solution for combating COVID-19.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

I acknowledge Mohanlal Sukhadia University, Udaipur for the infrastructure facility.

Funding statement

The work has been conducted in absence of any funding.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.imbio.2020.152021>.

References

- Annes, J.P., Rifkin, D.B., Munger, J.S., 2002. The integrin $\alpha\text{V}\beta\text{6}$ binds and activates latent TGF β3 . *FEBS Lett.* 511, 65–68. PMID:11821050.
- Arcangeli, A., Becchetti, A., 2010. Integrin structure and functional relation with ion channels. *Adv. Exp. Med. Biol.* 674, 1–7.
- Asmus, M.J., Barros, M.D., Liang, J., Chesrown, S.E., Hendeles, L., 2001. Pulmonary function response to EDTA, an additive in nebulized bronchodilators. *J. Allergy Clin. Immunol.* 107, 68–72.
- Bamonti, F., Fulgenzi, A., Novembrino, C., Ferrero, M.E., 2011. Metal chelation therapy in rheumatoid arthritis: a case report. Successful management of rheumatoid arthritis by metal chelation therapy. *Biometals* 24, 1093–1098. <https://doi.org/10.1007/s10534-011-9467-9>.
- Bella, J., Kolatkar, P.R., Marlor, C.W., Greve, J.M., Rossmann, M.G., 1998. The structure of the two amino-terminal domains of human ICAM-1 suggests how it functions as a rhinovirus receptor and as an LFA-1 integrin ligand. *Proc. Natl. Acad. Sci. U S A.* 95, 4140–4145.
- Bernhagen, D., Jungbluth, V., Quilis, N.G., Dostalek, J., White, P.B., Jalink, K., Timmerman, P., 2019. Bicyclic RGD peptides with exquisite selectivity for the integrin $\alpha\text{v}\beta\text{3}$ receptor using a "Random Design" approach. *ACS Comb. Sci.* 21, 198–206. <https://doi.org/10.1021/acscmb.8b00144>.
- Bernhagen, D., De Laporte, L., Timmerman, P., 2017. High-affinity RGD-knottin peptide as a new tool for rapid evaluation of the binding strength of unlabeled RGD-peptides to $\alpha\text{v}\beta\text{3}$, $\alpha\text{v}\beta\text{5}$, and $\alpha\text{5}\beta\text{1}$ integrin receptors. *Anal. Chem.* 89, 5991–5997.
- Berridge, M.J., 2012. Calcium signalling remodelling and disease. *Biochem. Soc. Trans.* 40, 297–309. <https://doi.org/10.1042/BST20110766>.
- Bianconi, D., Unseld, M., Prager, G.W., 2016. Integrins in the spotlight of cancer. *Int. J. Mol. Sci.* 17 (12), 2037. <https://doi.org/10.3390/ijms17122037>.
- Bretaña, N.A., Lu, C.T., Chiang, C.Y., Su, M.G., Huang, K.Y., Lee, T.Y., Weng, S.L., 2012. Identifying protein phosphorylation sites with kinase substrate specificity on human viruses. *PLOS ONE* 7, e40694 <https://doi.org/10.1371/journal.pone.0040694>.
- Blaurock-Busch, E., Busch, Y.M., 2014. Comparison of chelating agents DMPS, DMSA and EDTA for the diagnosis and treatment of chronic metal exposure. *Br. J. Med. Med. Res.* 4, 1821–1835.
- Blobe, G.C., Schiemann, W.P., Lodish, H.F., 2000. Role of transforming growth factor- β in human disease. *N. Engl. J. Med.* 342, 1350–1358.
- Cao, X., 2020. COVID-19: immunopathology and its implications for therapy. *Nat. Rev. Immunol.* <https://doi.org/10.1038/s41577-020-0308-3>.
- Cava, C., Bertoli, C., Castiglioni, I., 2020. In-silico discovery of candidate drugs against Covid-19. *Viruses* 12, 404. <https://doi.org/10.3390/v12040404>.
- Chang, K.W., Sheng, Y., Gombold, J.L., 2000. Coronavirus-induced membrane fusion requires the cysteine-rich domain in the spike protein. *Virology* 269, 212–224.
- Chen, Y., Guo, Y., Pan, Y., Zhuang, J.Z., 2020. Structure analysis of the receptor binding of 2019-nCoV. *Biochem. Biophys. Res. Commun.* 525, 135–140.
- Colson, P., Rolain, J.M., Lagier, J.C., Brouqui, P., Raoult, D., 2020. Chloroquine and hydroxychloroquine as available weapons to fight COVID-2019. *Int. J. Antimicrob. Agents* 55, 105932.
- Devaux, C.A., Rolain, J.M., Colson, P., Raoult, D., 2020. New insights on the antiviral effects of chloroquine against coronavirus: what to expect for COVID-19? *Int. J. Antimicrob. Agents* 55 (5), 105938 <https://doi.org/10.1016/j.ijantimicag.2020.105938>.
- Dong, X., Zhao, B., Iacob, R., et al., 2017. Force interacts with macromolecular structure in activation of TGF- β . *Nature* 542, 55–59.
- Erb, L., Liu, J., Ockerhausen, J., Kong, Q., Garrad, R.C., Griffin, K., Neal, C., Krugh, B., Santiago-Pérez, L.I., González, F.A., Gresham, H.D., Turner, J.T., Weisman, G.A., 2001. An RGD sequence in the P2y2 receptor interacts with $\alpha\text{v}\beta\text{3}$ integrins and is

- required for Go-mediated signal transduction. *J. Cell Biol.* 53, 491–502. <https://doi.org/10.1083/jcb.153.3.491>.
- Ferrero, M.E., 2016. Rationale for the successful management of EDTA chelation therapy in human burden by toxic metals. *BioMed Res. Int.* 2016, 8274504.
- Fulgenzi, A., Ferrero, M.E., 2019. EDTA chelation therapy for the treatment of neurotoxicity. *Int. J. Mol. Sci.* 20, 1019. <https://doi.org/10.3390/ijms20051019>.
- González-Cuevas, J., Navarro-Partida, J., Marquez-Aguirre, A.L., Bueno-Topete, M.R., Beas-Zarate, C., Armendáriz-Borunda, J., 2011. Ethylenediaminetetraacetic acid induces antioxidant and anti-inflammatory activities in experimental liver fibrosis. *Redox Rep.* 16, 62–70. <https://doi.org/10.1179/174329211X13002357050851>.
- Gordon, C.J., Tchesnokov, E.P., Woolner, E., Perry, J.K., Feng, J.Y., Porter, D.P., Gotte, M., 2020. Remdesivir is a direct-acting antiviral that inhibits RNA-dependent RNA polymerase from severe acute respiratory syndrome coronavirus 2 with high potency. *J. Biol. Chem.* <https://doi.org/10.1074/jbc.RA120.013679>.
- Goyer, R.A., Cherian, M.G., Jones, M.M., Reigart, J.R., 1995. Role of chelating agents for prevention, intervention and treatment of exposures to toxic metals. *Environ. Health Perspect.* 103, 1048–1052.
- Grosdidier, A., Zoete, V., Michielin, O., 2007. EADock: Docking of small molecules into protein active sites with a multiobjective evolutionary optimization. *Proteins* 67, 1010–1025.
- Hatley, R.J.D., Macdonald, S.J.F., Slack, R.J., Le, J., Ludbrook, S.B., Lukey, P.T., 2018. Anov-RGD integrin inhibitor toolbox: drug discovery insight, challenges and opportunities. *Angew. Chem. Int. Ed. Engl.* 57, 3298–3321. <https://doi.org/10.1002/anie.201707948>.
- Hicks, C., Gulick, R.M., 2009. Raltegravir: the first HIV type 1 integrase inhibitor. *Clin. Infect. Dis.* 48, 931–939. <https://doi.org/10.1093/cid/cin929>.
- Henderson, N.C., Arnold, T.D., Katamura, Y., Giacomini, M.M., Rodriguez, J.D., McCarty, J.H., Pellicoro, A., Raschperger, E., Betsholtz, C., Ruminski, P.G., Griggs, D.W., Prinsen, M.J., Maher, J.J., Iredale, J.P., Lacy-Hulbert, A., Adams, R.H., Sheppard, D., 2013. Targeting of alphaV integrin identifies a core molecular pathway that regulates fibrosis in several organs. *Nat. Med.* 19, 1617–1624. <https://doi.org/10.1038/nm.3282>.
- Hu, X., Dong, Q., Yang, J., Zhang, Y., 2016. Recognizing metal and acid radical ion binding sites by integrating ab initio modeling with template-based transfersals. *Bioinformatics* 32, 3260–3269.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., Xiao, Y., Gao, H., Guo, L., Xie, J., Wang, G., Jiang, R., Gao, Z., Jin, Q., Wang, J., Cao, B., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497–506.
- Hung, C.F., Wilson, C.L., Chow, Y.H., Schnapp, L.M., 2018. Role of integrin alpha8 in murine model of lung fibrosis. *PLOS ONE* 13, e0197937. <https://doi.org/10.1371/journal.pone.0197937>.
- Hussein, H.A.M., Walker, L.R., Abdel-Raouf, U.M., Desouky, S.A., Montasser, A.K.M., Akula, S.M., 2015. Beyond RGD: virus interactions with integrins. *Arch. Virol.* 160, 2669–2681. <https://doi.org/10.1007/s00705-015-2579-8>.
- Hysler, J.M., Estes, M.K., 2015. Pathophysiological consequences of calcium-conducting viroporins. *Annu. Rev. Virol.* 2, 473–496. <https://doi.org/10.1146/annurev-virology-100114-054846>.
- Inoue, Y., Tanaka, N., Tanaka, Y., Inoue, S., Morita, K., Zhuang, M., Hattori, T., Sugamura, K., 2007. Clathrin-dependent entry of severe acute respiratory syndrome coronavirus into target cells expressing ACE2 with the cytoplasmic tail deleted. *J. Virol.* 81, 8722–8729.
- Johnston, E.R., Radke, K., 2000. The SU and TM envelope protein subunits of bovine leukemia virus are linked by disulfide bonds, both in cells and in virions. *J. Virol.* 74, 2930–2935.
- Jia, N., Byrd-Leotis, L., Matsumoto, Y., Gao, C., Wein, A.N., Lobby, J.L., Kohlmeier, J.E., Steinhauer, D.A., Cummings, R.D., 2020. The human lung glycome reveals novel glycan ligands for influenza A virus. *Sci Rep.* 10, 5320. <https://doi.org/10.1038/s41598-020-62074-z>.
- Kandeel, M., Al-Nazawi, M., 2020. Virtual screening and repurposing of FDA approved drugs against COVID-19 main protease. *Life Sci.* 117627 <https://doi.org/10.1016/j.lfs.2020.117627>.
- Keating, J.A., Striker, R., 2012. Phosphorylation events during viral infections provide potential therapeutic targets. *Rev. Med. Virol.* 22, 166–181. <https://doi.org/10.1002/rmv.722>.
- Keck, F., Ataey, P., Amaya, M., Bailey, C., Narayanan, A., 2015. Phosphorylation of single stranded RNA virus proteins and potential for novel therapeutic strategies. *Viruses* 7, 5257–5273. <https://doi.org/10.3390/v7102872>.
- Lee, J.E., Saphire, E.O., 2009. Ebolavirus glycoprotein structure and mechanism of entry. *Future Virol.* 4, 621–635.
- Li, H., Zhou, Y., Zhang, M., Wang, H., Zhao, Q., Liu, J., 2020. Updated approaches against SARS-CoV-2. *Antimicrob. Agents Chemother.* <https://doi.org/10.1128/AAC.00483-20>.
- Liu, S., 2009. Radiolabeled cyclic RGD peptides as integrin $\alpha v \beta 3$ -targeted radiotracers: maximizing binding affinity via bivalency. *Bioconjug. Chem.* 20, 2199–2213.
- Lu, H., 2020. Drug treatment options for the 2019-new coronavirus (2019-nCoV). *Biosci. Trends* 14, 69–71. <https://doi.org/10.5582/bst.2020.01020>.
- Lu, M., Munger, J.S., Steadele, M., Busald, C., Tellier, M., Schnapp, L.M., 2002. Integrin $\alpha 8 \beta 1$ mediates adhesion to LAG-TGF $\beta 1$. *J. Cell Sci.* 115, 4641–4648.
- Ma, Y., Ai, G., Zhang, C., Zhao, M., Dong, X., Han, Z., Wang, Z., Zhang, M., Liu, Y., Gao, W., Li, S., Gu, Y., 2017. Novel linear peptides with high affinity to $\alpha v \beta 3$ integrin for precise tumor identification. *Theranostics* 7, 1511–1523. <https://doi.org/10.7150/thno.18401>.
- Madeira, F., Park, Y.M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey, A.R.N., Potter, S.C., Finn, R.D., Lopez, R., 2019. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res.* 47, W636–W641. <https://doi.org/10.1093/nar/gkz268>.
- Mehta, P., McAuley, D.F., Brown, M., Sanchez, E., Tattersall, R.S., Manson, J.J., 2020. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 395, P1033–1034. [https://doi.org/10.1016/S0140-6736\(20\)30628-0](https://doi.org/10.1016/S0140-6736(20)30628-0).
- Miller, C.G., Pozzi, A., Zent, R., Schwarzbauer, J.E., 2014. Effects of high glucose on integrin activity and fibronectin matrix assembly by mesangial cells. *Mol. Biol. Cell* 25, 2315–2317.
- Nagae, M., Re, S., Mihara, E., Nogi, T., Sugita, Y., Takagi, J., 2012. Crystal structure of $\alpha 5 \beta 1$ integrin ectodomain: atomic details of the fibronectin receptor. *J. Struct. Biol.* 197, 131–140. <https://doi.org/10.1083/jcb.201111077>.
- Nieto-Torres, J.L., Verdía-Báguena, C., Jimenez-Guardeno, J.M., Regla-Nava, J.A., Castaño-Rodríguez, C., Fernandez-Delgado, R., Torres, J., Aguilera, V.M., Enjuanes, L., 2015. Severe acute respiratory syndrome coronavirus E protein transports calcium ions and activates the NLRP3 inflammasome. *Virology* 485, 330–339. <https://doi.org/10.1016/j.virol.2015.08.010>.
- Ninsontia, C., Chanvorachote, P., 2014. Ouabain mediates integrin switch in human lung cancer cells. *Anticancer Res.* 34, 5495–5502.
- Pagni, M., Ioannidis, V., Cerutti, L., Zahn-Zabal, M., Jongeneel, C.V., Hau, J., Martin, O., Kuznetsov, D., Falquet, L., 2007. Improvements to an interactive resource for analyzing protein sequences. *Nucleic Acids Res.* 35, W433–W437. <https://doi.org/10.1093/nar/gkm352>.
- Parekh, R., Lorenzo, M.K., Shin, S.Y., Pozzi, A., Clark, A.L., 2014. Integrin $\alpha 1 \beta 1$ differentially regulates cytokine-mediated responses in chondrocytes. *Osteoarthr. Cartil.* 22, 499–508.
- Peymen, K., Watteyne, J., Froninckx, L., Schoofs, L., Beets, I., 2014. The FMRFamide-like peptide family in nematodes. *Front. Endocrinol.* 5, 1–21. <https://doi.org/10.3389/fendo.2014.00090>.
- Prabakaran, P., Gan, J., Feng, Y., Zhu, Z., Choudhary, V., Xiao, X., Ji, X., Dimitrov, D.S., 2006. Structure of severe acute respiratory syndrome coronavirus receptor-binding domain complexed with neutralizing antibody. *J. Biol. Chem.* 281, 15829–15836.
- Puchaka, P., Chaotham, C., Chanvorachote, P., 2016. α -Lipoic acid sensitizes lung cancer cells to chemotherapeutic agents and anoikis via integrin $\beta 1/\beta 3$ downregulation. *Int. J. Oncol.* 49, 1445–1456. <https://doi.org/10.3892/ijo.2016.3624>.
- Qin, C., Zhou, L., Hu, Z., Zhang, S., Yang, S., Tao, Y., Xie, C., Ma, K., Shang, K., Wang, W., Tian, D.-S., 2020. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clin. Infect. Dis.* <https://doi.org/10.1093/cid/ciaa248>.
- Qiu, H., Wu, J., Hong, L., Luo, Y., Song, Q., Chen, D., 2020. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet.* [https://doi.org/10.1016/S1473-3099\(20\)30198-5](https://doi.org/10.1016/S1473-3099(20)30198-5).
- Roth, T., Podesta, F., Stepp, M.A., Boeri, D., Lorenzi, M., 1993. Integrin overexpression induced by high glucose and by human diabetes: potential pathway to cell dysfunction in diabetic microangiopathy. *Proc. Natl. Acad. Sci. USA* 90 (20), 9640–9644. <https://doi.org/10.1073/pnas.90.20.9640>.
- Roussel, A.M., Hininger-Favier, I., Waters, R.S., Osman, M., Fernholz, K., Anderson, R.A., 2009. EDTA Chelation therapy, without added vitamin C, decreases oxidative DNA damage and lipid peroxidation. *Altern. Med. Rev.* 14, 56–62.
- Ruoslahti, E., 1988. Fibronectin and its receptors. *Ann. Rev. Biochem.* 57, 375–413.
- Ruoslahti, E., 1996. RGD and other recognition sequences for integrins. *Annu. Rev. Cell Dev. Biol.* 12, 697–715.
- Sawalha, A.H., Zhao, M., Coit, P., Lu, Q., 2020. Epigenetic dysregulation of ACE2 and interferon-regulated genes might suggest increased COVID-19 susceptibility and severity in lupus patients. *ClinImmunol* 108010. <https://doi.org/10.1016/j.clim.2020.108410>.
- Shah, B., Modi, P., Sagar, S.R., 2020. In silico studies on therapeutic agents for COVID-19: drug repurposing approach. *Life Sci.* <https://doi.org/10.1016/j.lfs.2020.117652>.
- Sheppard, D., 2008. The role of integrins in pulmonary fibrosis. *Eur. Respir. Rev.* 17, 157–162. <https://doi.org/10.1183/09059180.00010909>.
- Shi, Y. et al., 2020. Immunopathological characteristics of coronavirus disease 2019 cases in Guangzhou, China. Preprint at medRxiv. <https://doi.org/10.1101/2020.03.12.20034736>.
- Sigrist, C.J.A., Bridge, A., Mercier, P.L., 2020. A potential role for integrins in host cell entry by SARS-CoV-2. *Antiviral Res.* 177, 104759 <https://doi.org/10.1016/j.antiviral.2020.104759>.
- Sjaastad, M.D., Nelson, W.J., 1997. Integrin-mediated calcium signaling and regulation of cell adhesion by intracellular calcium. *Bioessays* 19, 47–55.
- Teoh, C.M., Tan, S.S.L., Tran, T., 2016. Integrins as therapeutic targets for respiratory diseases. *Curr. Mol. Med.* 15, 714–734.
- Tan, M., Liu, Y., Zhou, R., Deng, X., Li, F., Liang, K., Shi, Y., 2020. Immunopathological characteristics of coronavirus disease 2019 cases in Guangzhou, China. *Immunology* 160, 261–268. <https://doi.org/10.1111/imm.13223>.
- Tian, X., Li, C., Huang, A., Xia, S., Lu, S., Shi, Z., Lu, L., Jiang, S., Yang, Z., Wu, Y., Ying, T., 2020. Potent binding of 2019 novel coronavirus spike protein by a SARS coronavirus-specific human monoclonal antibody. *Emerg. Microbes Infect.* 9, 382–385. <https://doi.org/10.1080/22221751.2020.1729069>.
- Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., Olsson, I., Edlund, K., Lundberg, E., Navani, S., Szizyarto, C.A., Odeberg, J., Djureinovic, D., Takanen, J.O., Hober, S., Alm, T., Edqvist, P.H., Berling, H., Tegel, H., Mulder, J., Rockberg, J., Nilsson, P., Schwenk, J.M., Hamsten, M., von Feilitzen, K., Forsberg, M., Persson, L., Johansson, F., Zwahlen, M., von Heijne, G., Nielsen, J., Pontén, F., 2015. Tissue-based map of the human proteome. *Science* 347, 1260419. <https://doi.org/10.1126/science.1260419>.

- Uhlen, M., Karlsson, M.J., Zhong, W., Tebani, A., Pou, C., Mikes, J., Lakshmikanth, T., Forsström, B., Edfors, F., Odeberg, J., Mardinoglu, A., Zhang, C., von Feilitzen, K., Mulder, J., Sjöstedt, E., Hober, A., Oksvold, P., Zwahlen, M., Ponten, F., Lindskog, C., Sivertsson, Å., Fagerberg, L., Brodin, P., 2019. A genome-wide transcriptomic analysis of protein-coding genes in human blood cells. *Science* 366, eaax9198. <https://doi.org/10.1126/science.aax9198>.
- Van Agthoven, J.F., Xiong, J.P., Alonso, J.L., Rui, X., Adair, B.D., Goodman, S.L., Arnaout, M.A., 2014. Structural basis for pure antagonism of integrin $\alpha V\beta 3$ by a high-affinity form of fibronectin. *Nat. Struct. Mol. Biol.* 21, 383–388.
- Walls, A.C., Park, Y.J., Tortorici, M.A., Wall, A., McGuire, A.T., Veesele, D., 2020. Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell*. <https://doi.org/10.1016/j.cell.2020.02.058>.
- Wan, Y., Shang, J., Graham, R., Baric, R.S., Li, F., 2020. Receptor recognition by the novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS coronavirus. *J. Virol.* 94, e00127–20. <https://doi.org/10.1128/JVI.00127-20>.
- Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., Wang, B., Xiang, H., Cheng, Z., Xiong, Y., Zhao, Y., Li, Y., Wang, X., Peng, Z., 2020a. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 323, 1061–1069. <https://doi.org/10.1001/jama.2020.1585>.
- Wang, H., Yang, P., Liu, K., Guo, F., Zhang, Y., Zhang, G., Jiang, C., 2008. SARS coronavirus entry into host cells through a novel clathrin- and caveolae-independent endocytic pathway. *Cell Res.* 18, 290–301.
- Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M., Shi, Z., Hu, Z., Zhong, W., Xiao, G., 2020b. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res.* 30, 269–271.
- Wang, Y.-X., Ma, J.-R., Wang, S.-Q., Zeng, Y.-Q., Zhou, C.-Y., Ru, Y.-H., Zhang, L., Lu, Z.-G., Wu, M.-H., Li, H., 2020c. Utilizing integrating network pharmacological approaches to investigate the potential mechanism of Ma Xing Shi Gan Decoction in treating COVID-19. *Eur. Rev. Med. Pharmacol. Sci.* 24 (6), 3360–3384. https://doi.org/10.26355/eurrev_202003_20704.
- Wrapp, D., Wang, N., Corbett, K. S., Goldsmith, J. A., Hsieh, C. L., Abiona, O., Graham, B. S., McLellan, J. S., 2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science*. 2020, abb2507. 10.1126/science.abb2507.
- Wunderlich, V., Sydow, G., 1982. Disintegration of retroviruses by chelating agents. *Arch. Virol.* 73, 171–183.
- Xu, H., Zhong, L., Deng, J., Peng, J., Hongxia, D., Zeng, X., Li, T., Chen, Q., 2020a. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int. J. Oral Sci.* 12 <https://doi.org/10.1038/s41368-020-0074-x>.
- Xu, Z., Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L., Tai, Y., Bai, C., Gao, T., Song, J., Xia, P., Dong, J., Zhao, J., Wang, F.-S., 2020b. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir. Med.* 8, 420–422.
- Yuan, M., Wu, N.C., Zhu, X., Lee, C.-C.D., So, R.T.Y., Lv, H., Mok, C.K.P., Wilson, I.A., 2020. A highly conserved cryptic epitope in the receptor-binding domains of SARS-CoV-2 and SARS-CoV. *Science* 2020. <https://doi.org/10.1126/science.abb7269>.
- Zhang, H., Kang, Z., Gong, H., Xu, D., Wang, J., Li, Z., Cui, X., Xiao, J., Meng, T., Zhou, W., Liu, J., Xu, H., 2020. The digestive system is a potential route of 2019-nCoV infection: a bioinformatics analysis based on single-cell transcriptomes. *bioRxiv*. 2020, 927806; doi: <https://doi.org/10.1101/2020.01.30.927806>.
- Zhao, Y., Zhao, Z., Wang, Y., Zhou, Y., Ma, Y., Zuo, W., 2020. Single-cell RNA expression profiling of ACE2, the putative receptor of Wuhan 2019-nCoV. *BioRxiv*. <https://doi.org/10.1101/2020.01.26.919985>.