

Technical Note

Open Access

## A sequencing strategy for identifying variation throughout the prion gene of BSE-affected cattle

Michael L Clawson\*<sup>1</sup>, Michael P Heaton<sup>1</sup>, John W Keele<sup>1</sup>,  
Timothy PL Smith<sup>1</sup>, Gregory P Harhay<sup>1</sup>, Juergen A Richt<sup>2</sup> and  
William W Laegreid<sup>1,3</sup>

Address: <sup>1</sup>United States Department of Agriculture (USDA), Agricultural Research Service (ARS), US Meat Animal Research Center (USMARC), Clay Center, NE 68933, USA, <sup>2</sup>USDA, ARS, National Animal Disease Center, PO Box 70, Ames, IA 50010, USA and <sup>3</sup>University of Illinois at Champaign-Urbana, Department of Pathobiology, 2001 S. Lincoln Avenue, Urbana, IL 61802, USA

Email: Michael L Clawson\* - [mike.clawson@ars.usda.gov](mailto:mike.clawson@ars.usda.gov); Michael P Heaton - [mike.heaton@ars.usda.gov](mailto:mike.heaton@ars.usda.gov);  
John W Keele - [john.keele@ars.usda.gov](mailto:john.keele@ars.usda.gov); Timothy PL Smith - [tim.smith@ars.usda.gov](mailto:tim.smith@ars.usda.gov); Gregory P Harhay - [gregory.harhay@ars.usda.gov](mailto:gregory.harhay@ars.usda.gov);  
Juergen A Richt - [juergen.richt@ars.usda.gov](mailto:juergen.richt@ars.usda.gov); William W Laegreid - [laegreid@uiuc.edu](mailto:laegreid@uiuc.edu)

\* Corresponding author

Published: 23 June 2008

Received: 18 March 2008

BMC Research Notes 2008, 1:32 doi:10.1186/1756-0500-1-32

Accepted: 23 June 2008

This article is available from: <http://www.biomedcentral.com/1756-0500/1/32>

© 2008 Clawson et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Classical and atypical bovine spongiform encephalopathies (BSEs) are cattle prion diseases. Distinct bovine prion gene (*PRNP*) alleles have been associated with classical and atypical BSE susceptibility. However, the full extent of *PRNP* allele association with BSE susceptibility is not known. A systematic sequence-based genotyping method that detects variation throughout *PRNP* would be useful for: 1) detecting rare *PRNP* alleles that may be present in BSE-affected animals and 2) testing *PRNP* alleles for an association with either classical or atypical BSE susceptibility.

**Findings:** We improved a Sanger-based sequencing strategy for detecting bovine *PRNP* variation through all exons, introns, and part of the promoter (25.2 kb). Our current method can detect 389 known and other potentially unknown *PRNP* polymorphisms that may be present in BSE-affected cattle. We determined *PRNP* genotypes for the first U.S. BSE case and her sire. Previously unknown *PRNP* polymorphisms were not detected in either animal and all *PRNP* genotypes support the sire-daughter relationship.

**Conclusion:** The methodologies described here characterize variation throughout *PRNP*. Consequently, rare *PRNP* alleles that may be present in BSE-affected cattle can be detected.

### Introduction

Transmissible spongiform encephalopathies (TSEs), also known as prion diseases, are a class of neurodegenerative disorders that occur in humans, ruminants, cats, and mink [1]. Three distinct TSEs afflict cattle: classical bovine spongiform encephalopathy (BSE), atypical H-type BSE, and atypical L-type BSE [2-4]. Bovine prion gene (*PRNP*)

alleles are associated with classical and atypical BSE susceptibility [5-8]. However, the extent of bovine *PRNP* alleles that predict BSE susceptibility has not yet been determined.

Classical BSE is acquired by cattle through the consumption of meat and bone meal contaminated with the infec-

tious prion agent [9]. Classical BSE susceptibility is associated with the deletion alleles of two bovine *PRNP* insertion/deletion polymorphisms, one within the promoter region and the other in intron 1 [5-7]. Both of these polymorphisms have alleles in linkage disequilibrium (LD) with the alleles of at least 43 other *PRNP* polymorphisms in *Bos taurus* cattle [10]. Thus, additional *PRNP* alleles are likely to associate with classical BSE susceptibility with comparable if not greater significance than the indel alleles.

Atypical BSEs have recently been identified in Asian, North American, and European cattle [11]. Two *PRNP* alleles are implicated with atypical BSE susceptibility. A non-synonymous polymorphism (E211K) was found by one of us (J.A.R.) within the prion coding region of an H-type atypical BSE case identified in the U.S. (2006). The E211K polymorphism is homologous to the human E200K polymorphism, a risk factor for genetic Creutzfeldt-Jakob disease [12], and is the suspected cause of the atypical BSE case. The 211 K allele is exceedingly rare and has not been found in other atypical BSE cases or healthy cattle [10,13,14]. However, a *PRNP* haplotype that is associated with atypical BSE susceptibility was found in H- and L-type atypical cases from Canada, France, and the U.S., and may have widespread involvement with atypical BSE [8]. The haplotype spans part of *PRNP* intron 2, the entire coding region of exon 3, and part of the three prime untranslated region of exon 3 (13 kb) [8,10]. Alleles that may be causative for atypical BSE susceptibility, including those that may be within *PRNP*, are thought to be linked with the implicated haplotype in atypical BSE cases [8].

Efforts to identify *PRNP* alleles that predict classical or atypical BSE susceptibility require improved methods to: 1) efficiently extract DNA from available tissue samples and 2) identify variation throughout *PRNP*. The brainstem obex is the specified sample for BSE testing [15]. If an obex sample is BSE positive and the carcass from which

it originated is destroyed before additional samples are collected, the obex becomes the sole source of transmissible material and DNA from the BSE-afflicted animal. Thus, the ability to obtain DNA from obex tissue is a critical prerequisite for *PRNP* genotyping. The ability to accurately genotype known *PRNP* polymorphisms and identify additional *PRNP* polymorphisms that may be present in BSE-afflicted animals is also important. Here, we provide an efficient method of DNA extraction from bovine obex and report a Sanger-based sequencing strategy that comprehensively identifies variation throughout *PRNP* (25.2 kb). We determined the *PRNP* sequence and haplotypes of the U.S. case of classical BSE (2003) and her sire, show that both animals do not have previously unrecognized *PRNP* variation, and that their parent-offspring relationship is supported throughout *PRNP*.

**DNA isolation method**

For DNA isolation, ten cattle obexes (0.7 – 1.9 grams) were individually homogenized in PBS (weight per volume = 25%). Three hundred and fifty microliters of the homogenate, containing 87.5 mg of obex, was digested overnight in proteinase K. One half of the digestion was frozen while the other half was subjected to standard Tris-saturated phenol:chloroform:isoamyl alcohol (25:24:1, pH 8) and chloroform extractions, followed by salt precipitation, ethanol washing, and re-suspension in TE (see additional file 1 for supplementary methods on DNA isolation from cattle obex tissue and quantification). The procedure yielded a mean of 12.1 µg DNA (high = 17.34 µg, low = 7.64 µg, Table 1). Each extraction yielded more than 3× the 2.4 µg of DNA typically used to comprehensively amplify and sequence *PRNP*. The DNAs were obtained from the equivalent of 43.75 mg of obex, or less than 7% of each obex. Thus, DNA sufficient for *PRNP* genotyping can be obtained from a small portion of the obex tissue.

**Table 1: Cattle obex DNA yield and quality**

Sample <sup>a</sup>	DNA Yield (µg)	DNA concentration (µg/mL) <sup>b</sup>	260/280 nm
1	15.55	155.54	1.87
2	17.34	173.39	1.87
3	10.78	107.82	1.87
4	13.34	133.39	1.84
5	7.80	77.97	1.83
6	7.64	76.40	1.83
7	17.81	178.07	1.86
8	9.76	97.55	1.80
9	10.09	100.86	1.83
10	10.70	106.98	1.84

<sup>a</sup>Ten cattle obex tissues (44 mg each).

<sup>b</sup>0.1 mL suspension.

Mean yield = 12.1 µg, SD = 3.7, 95% CI = ± 2.3.

**Development of oligonucleotides for PRNP amplification and sequence genotyping**

We improved a set of oligonucleotides that can be used to: 1) collectively amplify *PRNP* (25.2 kb) on 24 overlapping amplicons (Figure 1b) and 2) genotype *PRNP* variation by sequence [10]. This new set is comprised of 189 oligonucleotides, of which 168 have been previously described [10]. All of the oligonucleotides are designed around 389 known *PRNP* polymorphisms (48 amplification/sequencing, 141 sequencing; see additional file 2 for a table of oligonucleotides for *PRNP* amplification and sequencing). Ten *PRNP* amplicons described previously [10] have been modified with redesigned oligonucleotides due to unsatisfactory performance, including one that covers the InDel within intron 1 that is associated with classical BSE susceptibility and another that covers the entire coding region of exon 3. Strict PCR conditions and/or reagents for many of these amplicons are required (see additional file #3 for supplementary PCR information). All 189 oligonucleotides of the set have been tested for amplification and/or sequencing performance on 192 cattle DNAs representing 21 breeds (see additional file #3 for breed information) [10,16]. Three hundred and eighty-eight *PRNP* polymorphisms have been observed in these cattle DNAs [10]. None of the oligonucleotides hybridize to genomic sites containing the 388 polymorphisms or E211K (Figure 2).


Additional *PRNP* variation is predicted to exist in cattle and some of it could affect the hybridization of oligonucleotides within the set. This likelihood is accounted for by overlapping *PRNP* amplicons coupled with overlapping *PRNP* nucleotide coverage by sequencing oligonucleotides (Figure 2). Thus, the set will facilitate the detection

of both known and unknown *PRNP* polymorphisms, including those within primer hybridization sites.


Complete PCR coverage of *PRNP* (25.2 kb) from a BSE-positive animal can be conducted in a single 96-well plate using a thermocycler capable of zoned heating cycles (see additional file #4 for a 96-well PCR design). Two 55- $\mu$ L PCRs for each of the 24 *PRNP* amplicons yield enough template to sequence *PRNP* at least twice with each of the 189 oligonucleotides on an ABI capillary 3730 sequencer. This coverage can yield approximately eight-fold coverage of each *PRNP* nucleotide base throughout the 25.2-kb region, except for several small problematic regions (Figure 2). Each amplification reaction requires 50 ng of DNA. Consequently, thorough coverage of *PRNP* can be achieved with 2.4  $\mu$ g of genomic DNA. All of the sequencing reactions required for duplicate coverage of *PRNP* can fit on a single 384-well plate.

**Complete PRNP sequences of the U.S. classical BSE case of 2003 and her sire**


The oligonucleotide set was used to amplify and sequence *PRNP* from: 1) the U.S. case of classical BSE (2003), a Holstein cow imported from Canada, and 2) her Holstein sire. All sequence nucleotides were scored with Phred, aligned with Phrap, edited in Consed, and genotyped with Polyphred software [17]. Unambiguous genotypes for 388 *PRNP* polymorphisms were scored for the U.S. BSE case and her sire (see additional file #5 for a complete list of genotypes and GenBank files [EU130450](#) and [EU130451](#)). Sequences from both animals did not reveal any previously unknown *PRNP* polymorphisms. Both animals are homozygous for the deletion alleles of the two InDels previously associated with classical BSE sus-



**Figure 1**  
**PRNP coverage. (a) Sequence-based physical map of bovine PRNP.** Orange and yellow arrows represent untranslated and protein coding regions, respectively. **(b) PRNP amplicons.** Grey rectangles represent 24 overlapping amplicons. Numbers assigned to the amplicons are above the rectangles. Numbers of polymorphisms observed in U.S. cattle in genomic regions spanned by the amplicons are shown inside the rectangles.



**Figure 2**  
**Physical maps of the 24 BTAPRNP amplicons that collectively span bovine PRNP (25.2 kb).** Alternate splicing of Exon I is annotated on DS13a2 and DS13b. Amplicon sizes may vary by the presence or absence of Insertion/Deletion alleles.



**Figure 3**  
**PRNP haplotypes of the U.S. classical BSE case and her sire. (a) Physical map of haplotype networks one and two that span PRNP regions of high and low LD, respectively. (b) PRNP haplotype relationships.** Circles represent PRNP haplotypes observed in a group of 86 U.S. Holsteins. Circle areas are proportional to haplotype frequencies. Rectangles represent haplotypes with a frequency of less than 2.5% in the Holstein group. Vertical brackets denote haplotypes with deletion alleles from a 23-bp and/or 12-bp InDel previously associated with classical BSE susceptibility. **(c) PRNP diplotypes of the U.S. classical BSE case and her sire.** The frequencies of diplotypes "Ab" and "Ae" in the Holstein group are 0.264 and 0.030, respectively.

ceptibility. All of the *PRNP* polymorphism genotypes support the sire-offspring relationship.

#### **PRNP haplotypes of the BSE case and her sire**

Nineteen haplotype tagging polymorphisms (htSNPs) [10] were used to phase *PRNP* haplotypes of the classical BSE case and her sire (see additional file #6 for a description of the phasing method [18,19]). The htSNPs define *PRNP* haplotypes within two independent Median-Joining networks that account for high and low LD regions observed in *Bos taurus* cattle populations [10] (Figure 3, see additional file #7 for haplotype sequences). Nine htSNPs define haplotypes within a region of high LD that spans the *PRNP* promoter through a small portion of intron 2 (network one). Ten other htSNPs define haplotypes within a *PRNP* region of low LD that span most of intron 2 through the 3'UTR (network two). The htSNPs collectively define haplotype diversity within and across *PRNP* and have been previously used to test *PRNP* variation for an association with atypical BSE susceptibility [8].

*PRNP* haplotypes of the classical BSE case and her sire were compared to each other and to those from 86 U.S. Holsteins, of which 82 were part of the USMARC Dairy Cattle Panel (MDCP1.5) [10] and 4 were part of the USMARC Beef Cattle Discovery Panel 2.1 (MBCDP2.1) [16]. The classical BSE case and her sire were both homozygous for a haplotype in network one that is common in U.S. Holstein (haplotype A, Figure 3). The two InDel deletion alleles that associate with BSE susceptibility are present on this haplotype. However, while the sire was homozygous for a common *PRNP* haplotype in network two (haplotype b), the classical BSE case was heterozygous for the same haplotype and one much less common (haplotype e). The frequencies of haplotype combinations "Ab" and "Ae" in the Holstein DNAs are 0.264 and 0.030, respectively. The significance of these haplotype combinations and classical BSE susceptibility is unknown. *PRNP* haplotype combinations such as those described here can be tested for association with either classical or atypical BSEs in larger studies.

#### **Conclusion**

BSE has grown in complexity with the identification of classical and atypical types. *PRNP* alleles associate with classical and atypical BSE susceptibility. The methodologies described here characterize variation throughout *PRNP*. Consequently, rare *PRNP* alleles that may be present in BSE-affected cattle can be detected.

#### **Competing interests**

The authors declare that they have no competing interests.

#### **Authors' contributions**

MLC conceived the project, developed DNA extraction procedures, designed and tested oligonucleotides, analyzed sequence, and wrote the manuscript. MPH designed oligonucleotides, developed DNA extraction procedures, and contributed DNA samples to the study. JWK assisted in sequence analyses. TPLS conducted sequencing and sequence analyses. GPH assisted in haplotype analyses. JAR contributed DNA samples to the study and improved oligonucleotide design. WWL designed oligonucleotides and assisted in sequence analyses. All authors read and approved the final manuscript.

#### **Additional material**

##### **Additional file 1**

*Supplementary Methods – DNA isolation from cattle obex and quantification. Detailed technical and reagent information regarding DNA isolation from cattle obex tissue.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S1.doc>]

##### **Additional file 2**

*Supplementary Table – Oligonucleotides for PRNP amplification and DNA sequencing [USMARC Bovine PRNP Oligonucleotide Set Version 4 (MBPRNPOSv4)]. Complete list of oligonucleotides used for PRNP amplification and DNA sequencing.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S2.xls>]

##### **Additional file 3**

*Supplementary Methods – Cattle DNAs used for oligonucleotide testing, PCR reagents, and thermocycling conditions. Complete list of cattle breed DNAs used for oligonucleotide testing and technical information on PCR reagents and thermocycling conditions for PRNP amplification.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S3.doc>]

##### **Additional file 4**

*Supplementary Figure – PRNP PCRs in 96 well format. Illustration of PRNP amplification in a 96 well plate.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S4.ppt>]

##### **Additional file 5**

*Supplementary Table – PRNP genotypes of the U.S. classical BSE case and her sire. Side-by-side comparison of PRNP genotypes of the U.S. classical BSE case and her sire.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S5.xls>]

### Additional file 6

Supplementary Methods – PRNP haplotype reconstructions. Method for determining PRNP haplotypes in the group of 86 Holsteins, the classical BSE case, and her sire.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S6.doc>]

### Additional file 7

Supplementary Table – PRNP haplotype sequences in Networks 1 and 2. PRNP haplotype sequences represented in Networks 1 and 2 of Figure 3. Note: This table was originally published in PLoS ONE [8].

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1756-0500-1-32-S7.xls>]

### Acknowledgements

We thank Gennie Schuller-Chavez for outstanding technical support; Dr. Mohammad Koohmaraie for coordinating the collection of cattle obex samples at a Nebraska slaughter plant; Drs. Shuna Jones and Dee Griffin for collecting the obex samples; Tammy Sorensen and Jacky Carnahan for technical support; Drs. Jeff Gawronski and Dan Nonneman for critically reviewing the manuscript; and Joan Rosch for secretarial support. This research was supported by the USDA National Research Initiative, Competitive Grant No. 2005-35212-15890 (MLC, MPH, WWL) and the Agricultural Research Service (all authors).

### References

- Prusiner SB: **Prions**. *Proc Natl Acad Sci U S A* 1998, **95**:13363-13383.
- Wells GA, Scott AC, Johnson CT, Gunning RF, Hancock RD, Jeffrey M, Dawson M, Bradley R: **A novel progressive spongiform encephalopathy in cattle**. *Vet Rec* 1987, **121**:419-420.
- Biacabe AG, Laplanche JL, Ryder S, Baron T: **Distinct molecular phenotypes in bovine prion diseases**. *EMBO Rep* 2004, **5**(1):110-114.
- Casalone C, Zanusso G, Acutis P, Ferrari S, Capucci L, Tagliavini F, Monaco S, Caramelli M: **Identification of a second bovine amyloidotic spongiform encephalopathy: molecular similarities with sporadic Creutzfeldt-Jakob disease**. *Proc Natl Acad Sci U S A* 2004, **101**(9):3065-3070.
- Sander P, Hamann H, Pfeiffer I, Wemheuer W, Brenig B, Groschup MH, Ziegler U, Distl O, Leeb T: **Analysis of sequence variability of the bovine prion protein gene (PRNP) in German cattle breeds**. *Neurogenetics* 2004, **5**(1):19-25.
- Juling K, Schwarzenbacher H, Williams JL, Fries R: **A major genetic component of BSE susceptibility**. *BMC Biol* 2006, **4**:33.
- Haase B, Doherr MG, Seuberlich T, Drogemuller C, Dolf G, Nicken P, Schiebel K, Ziegler U, Groschup MH, Zurbriggen A, Leeb T: **PRNP promoter polymorphisms are associated with BSE susceptibility in Swiss and German cattle**. *BMC Genet* 2007, **8**:15.
- Clawson ML, Richt JA, Baron T, Biacabe AG, Czub S, Heaton MP, Smith TPL, Laegreid WW: **Association of a bovine prion gene haplotype with atypical BSE**. *PLoS ONE* 2008, **3**(3):e1830.
- Wilesmith JW, Ryan JB, Atkinson MJ: **Bovine spongiform encephalopathy: epidemiological studies on the origin**. *Vet Rec* 1991, **128**:199-203.
- Clawson ML, Heaton MP, Keele JW, Smith TPL, Harhay GP, Laegreid WW: **Prion gene haplotypes of U.S. cattle**. *BMC Genet* 2006, **7**:51.
- Ducrot C, Arnold M, de Koeijer A, Heim D, Calavas D: **Review on the epidemiology and dynamics of BSE epidemics**. *Vet Res* 2008, **39**:15.
- Kovacs GG, Puopolo M, Ladogana A, Pocchiari M, Budka H, van Duijn C, Collins SJ, Boyd A, Giulivi A, Coulthart M, Delasnerie-Laupretre N, Brandel JP, Zerr I, Kretzschmar HA, de Pedro-Cuesta J, Calero-Lara M, Glatzel M, Aguzzi A, Bishop M, Knight R, Belay G, Will R, Mitrova E: **Genetic prion disease: the EUROCJD experience**. *Hum Genet* 2005, **118**(2):166-174.
- Heaton MP, Leymaster KA, Freking BA, Hawk DA, Smith TPL, Keele JW, Snelling WM, Fox JM, Chitko-McKown CG, Laegreid WW: **Prion gene sequence variation within diverse groups of U.S. sheep, beef cattle, and deer**. *Mamm Genome* 2003, **14**(11):765-777.
- Heaton MP, Keele JW, Harhay GP, Richt JA, Koohmaraie M, Wheeler TL, Shackelford SD, Casas E, King DA, Sonstegard TS, Van Tassell CP, Neibergs HL, Chase CC Jr, Kalbfleisch TS, Smith TP, Clawson ML, Laegreid WW: **Prevalence of the prion protein gene E211K variant in U.S. cattle**. *BMC Vet Res* 2008, **4**:25.
- USDA: **Procedures Manual Bovine Spongiform Encephalopathy (BSE) Ongoing Surveillance Plan**. U.S. Department of Agriculture Animal and Plant Health Inspection Service Veterinary Services; 2006.
- Heaton MP, Chitko-McKown CG, Grosse WM, Keele JW, Keen JE, Laegreid WW: **Interleukin-8 haplotype structure from nucleotide sequence variation in commercial populations of U.S. beef cattle**. *Mamm Genome* 2001, **12**:219-226.
- Stephens M, Sloan JS, Robertson PD, Scheet P, Nickerson DA: **Automating sequence-based detection and genotyping of SNPs from diploid samples**. *Nat Genet* 2006, **38**(3):375-381.
- Stephens M, Smith NJ, Donnelly P: **A new statistical method for haplotype reconstruction from population data**. *Am J Hum Genet* 2001, **68**:978-989.
- Stephens M, Scheet P: **Accounting for decay of linkage disequilibrium in haplotype inference and missing-data imputation**. *Am J Hum Genet* 2005, **76**:448-462.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

