

## Erlotinib hydrochloride: an anticancer agent

S. Selvanayagam,<sup>a\*</sup> B. Sridhar<sup>b</sup> and K. Ravikumar<sup>b</sup>

<sup>a</sup>Department of Physics, Kalasalingam University, Krishnankoil 626 190, India, and

<sup>b</sup>Laboratory of X-ray Crystallography, Indian Institute of Chemical Technology, Hyderabad 500 007, India

Correspondence e-mail: s\_selvanayagam@rediffmail.com


Received 11 April 2008; accepted 23 April 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(C-C) = 0.004$  Å;  $R$  factor = 0.073;  $wR$  factor = 0.151; data-to-parameter ratio = 18.2.

In the cation of the title compound,  $C_{22}H_{24}N_3O_4^+ \cdot Cl^-$ , an active ingredient of the anticancer drug also known as Tarceva, the quinazoline ring system is planar within 0.044 (3) Å. The dihedral angle formed by the mean planes of the two six-membered quinazoline rings is 3.2 (1)°. Both N-bound H atoms participate in N—H···Cl bonds, which link the ions into infinite chains running along the  $b$  axis. C—H···O interactions involving neighboring cations provide additional stabilization of these aggregates.

### Related literature

For related literature, see: Herbst *et al.* (2005); Minna & Dowell (2005); Li *et al.* (2007); Xia (2005). For bond-length data, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

$C_{22}H_{24}N_3O_4^+ \cdot Cl^-$   
 $M_r = 429.89$   
Monoclinic,  $P2_1/c$

$a = 14.5351$  (15) Å  
 $b = 18.4863$  (19) Å  
 $c = 8.1222$  (8) Å

$\beta = 102.966$  (2)°  
 $V = 2126.8$  (4) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.21$  mm<sup>-1</sup>  
 $T = 293$  (2) K  
 $0.26 \times 0.22 \times 0.20$  mm

#### Data collection

Bruker APEX area-detector diffractometer  
Absorption correction: none  
24377 measured reflections

5001 independent reflections  
3649 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.050$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.073$   
 $wR(F^2) = 0.151$   
 $S = 1.16$   
5001 reflections  
275 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.33$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.25$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N3—H3···Cl1	0.86	2.46	3.277 (2)	160
N1—H1···Cl1 <sup>i</sup>	0.86	2.23	3.066 (2)	165
Cl1—H1···O4 <sup>i</sup>	0.93	2.46	3.372 (3)	167

Symmetry code: (i)  $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97 and PARST (Nardelli, 1995).

SS thanks the Vice Chancellor and the management of Kalasalingam University, Anand Nagar, Krishnankoil, for their support and encouragement.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YA2075).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Bruker (2001). SAINT and SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Herbst, R. S., Prager, D., Hermann, R., Fehrenbacher, L., Johnson, B. E., Sandler, A., Kris, M. G., Tran, H. T., Klein, P., Li, X., Ramies, D., Johnson, D. H. & Miller, V. A. (2005). *J. Clin. Oncol.* **23**, 5892–5899.
- Li, Z., Xu, M., Xing, S., Ho, W. T., Ishii, T., Li, Q., Fu, X. & Zhao, Z. J. (2007). *J. Biol. Chem.* **282**, 3428–3432.
- Minna, J. D. & Dowell, J. (2005). *Nat. Rev. Drug Discov.* pp. S14–S15.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Xia, M. (2005). *Acta Cryst. E* **61**, o3380–o3382.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o931 [doi:10.1107/S1600536808011707]

## Erlotinib hydrochloride: an anticancer agent

**S. Selvanayagam, B. Sridhar and K. Ravikumar**

### Comment

Erlotinib hydrochloride, (I), also known under its tradename Tarceva, is a potent reversible epidermal growth factor receptor tyrosine kinase inhibitor with single agent activity in patients with non-small lung cancer, pancreatic cancer and several other types of cancer (Herbst *et al.*, 2005). It is the first drug to demonstrate an increase in survival in phase III trials in patients with advanced non-small cell lung cancer (Minna & Dowell, 2005). It has recently been shown to be potent inhibitor of JAK2 V617F activity; JAK2 V617F is a mutant of tyrosine kinase JAK2 (Li *et al.*, 2007). As no crystal structure of the title compound has yet been published, we have undertaken the single-crystal X-ray diffraction study and report here its results.

The X-ray study confirmed the molecular structure and atomic connectivity for (I), as illustrated in Fig. 1. The C21—C22 bond length [1.170 (4) Å] is consistent with its acetylenic character, as evidenced by literature value of 1.174 (11) Å (see Allen *et al.*, 1987). The geometry of the quinazoline ring system is comparable to that in the reported related structure (Xia, 2005).

The bicyclic system is effectively planar with a maximum deviation of 0.044 (3) Å for the C1 atom. The dihedral angle formed by the mean planes of two six membered rings of quinazoline moiety is 3.2 (1)°. The benzene ring C15—C20 and its attached ethynyl group are coplanar with a maximum deviation -0.011 (3) Å for the C20 atom. The dihedral angle between this ring and quinazoline ring system is 34.4 (1)°. The short contacts H3···H4 (2.12 Å) and H7···H9A (2.14 Å) result in substantial widening of the C2—N3—C15 and C6—O1—C9 bond angles [127.4 (2)° and 117.1 (2)°, respectively].

Both N-bound H atoms (HN1 and H3) participate in H-bonds with the Cl1 anion (Table 2). These bonds link cations and anions into the infinite chains running along the *b* axis of the crystal. The C1—H1···O4 interactions involving neighboring cations provide additional stabilization for these chains (Fig. 2).

### Experimental

In order to obtain crystals suitable for X-ray study, commercially available erlotinib hydrochloride was dissolved in a methanol-water solution (90:10v/v); the solvents were then allowed to evaporate slowly.

### Refinement

The acetylenic H22 atom was located in a difference Fourier map and refined isotropically [C22—H22 0.95 (3) Å]; all other H atoms were positioned geometrically with C—H distances of 0.93–0.97 Å, N—H 0.86 Å and were included in the refinement in the riding motion approximation with  $U_{\text{iso}} = 1.5U_{\text{eq}}(\text{C})$  for methyl H and  $1.2U_{\text{eq}}$  for all other H atoms.

# supplementary materials

---

## Figures




Fig. 1. The structure and atom-numbering scheme for (I); displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radius.




Fig. 2. Molecular packing of (I) viewed along the *c* axis; H-bonds are shown as dashed lines; H atoms, not involved in H-bonds, have been omitted.

## Erlotinib hydrochloride

### Crystal data

$C_{22}H_{24}N_3O_4^+ \cdot Cl^-$	$F_{000} = 904$
$M_r = 429.89$	$D_x = 1.343 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 14.5351 (15) \text{ \AA}$	Cell parameters from 9448 reflections
$b = 18.4863 (19) \text{ \AA}$	$\theta = 2.2\text{--}23.4^\circ$
$c = 8.1222 (8) \text{ \AA}$	$\mu = 0.21 \text{ mm}^{-1}$
$\beta = 102.966 (2)^\circ$	$T = 293 (2) \text{ K}$
$V = 2126.8 (4) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.26 \times 0.22 \times 0.20 \text{ mm}$

### Data collection

Bruker APEX area-detector diffractometer	3649 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.050$
Monochromator: graphite	$\theta_{\max} = 28.0^\circ$
$T = 293(2) \text{ K}$	$\theta_{\min} = 1.8^\circ$
$\omega$ scans	$h = -18 \rightarrow 19$
Absorption correction: none	$k = -24 \rightarrow 24$
24377 measured reflections	$l = -10 \rightarrow 10$
5001 independent reflections	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.073$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.151$	$w = 1/[\sigma^2(F_o^2) + (0.0551P)^2 + 0.7563P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.16$	$(\Delta/\sigma)_{\max} = 0.001$
5001 reflections	$\Delta\rho_{\max} = 0.33 \text{ e \AA}^{-3}$
275 parameters	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.50200 (5)	0.34427 (3)	0.62635 (9)	0.0469 (2)
O1	0.27725 (11)	0.61695 (9)	0.9544 (2)	0.0395 (4)
O2	0.10485 (14)	0.70220 (13)	0.9170 (3)	0.0648 (6)
O3	0.29876 (12)	0.48732 (9)	0.8521 (2)	0.0417 (4)
O4	0.20059 (13)	0.31485 (10)	0.6875 (3)	0.0514 (5)
N1	0.57019 (14)	0.69146 (10)	0.8273 (3)	0.0395 (5)
HN1	0.5603	0.7355	0.8527	0.047*
N2	0.67238 (14)	0.60941 (11)	0.7423 (3)	0.0408 (5)
N3	0.63506 (13)	0.48880 (10)	0.7058 (3)	0.0350 (5)
H3	0.5945	0.4550	0.7065	0.042*
C1	0.64787 (18)	0.67467 (13)	0.7780 (4)	0.0435 (7)
H1	0.6887	0.7123	0.7678	0.052*
C2	0.61060 (16)	0.55508 (12)	0.7464 (3)	0.0318 (5)
C3	0.52182 (16)	0.56803 (12)	0.7918 (3)	0.0310 (5)
C4	0.45169 (16)	0.51536 (12)	0.7961 (3)	0.0321 (5)
H4	0.4604	0.4681	0.7637	0.039*
C5	0.37123 (16)	0.53333 (13)	0.8474 (3)	0.0324 (5)
C6	0.35795 (16)	0.60529 (13)	0.9019 (3)	0.0326 (5)
C7	0.42426 (17)	0.65734 (12)	0.8963 (3)	0.0346 (5)
H7	0.4158	0.7044	0.9305	0.042*
C8	0.50471 (16)	0.63882 (12)	0.8385 (3)	0.0320 (5)

## supplementary materials

---

C9	0.26703 (18)	0.68616 (14)	1.0300 (4)	0.0417 (6)
H9A	0.2742	0.7248	0.9530	0.050*
H9B	0.3151	0.6919	1.1333	0.050*
C10	0.17127 (19)	0.68926 (15)	1.0675 (4)	0.0448 (6)
H10A	0.1574	0.6439	1.1166	0.054*
H10B	0.1689	0.7277	1.1478	0.054*
C11	0.0123 (3)	0.7093 (3)	0.9450 (7)	0.1109 (16)
H11A	-0.0314	0.7178	0.8392	0.166*
H11B	0.0106	0.7492	1.0197	0.166*
H11C	-0.0047	0.6656	0.9949	0.166*
C12	0.30023 (17)	0.41876 (13)	0.7695 (4)	0.0422 (6)
H12A	0.3512	0.3889	0.8318	0.051*
H12B	0.3093	0.4256	0.6558	0.051*
C13	0.2062 (2)	0.38355 (15)	0.7650 (4)	0.0510 (7)
H13A	0.1974	0.3784	0.8793	0.061*
H13B	0.1560	0.4142	0.7029	0.061*
C14	0.1801 (3)	0.3188 (2)	0.5094 (5)	0.0765 (11)
H14A	0.1774	0.2709	0.4631	0.115*
H14B	0.2286	0.3460	0.4740	0.115*
H14C	0.1203	0.3423	0.4699	0.115*
C15	0.72087 (16)	0.46773 (12)	0.6616 (3)	0.0327 (5)
C16	0.71529 (18)	0.41268 (14)	0.5436 (3)	0.0396 (6)
H16	0.6575	0.3912	0.4968	0.048*
C17	0.7960 (2)	0.38986 (16)	0.4959 (4)	0.0474 (7)
H17	0.7922	0.3529	0.4169	0.057*
C18	0.88191 (19)	0.42120 (15)	0.5641 (4)	0.0473 (7)
H18	0.9356	0.4060	0.5296	0.057*
C19	0.88874 (17)	0.47541 (13)	0.6842 (3)	0.0395 (6)
C20	0.80763 (17)	0.49842 (13)	0.7347 (3)	0.0371 (6)
H20	0.8118	0.5341	0.8168	0.045*
C21	0.9793 (2)	0.50773 (16)	0.7563 (4)	0.0490 (7)
C22	1.0541 (2)	0.5325 (2)	0.8100 (5)	0.0653 (9)
H22	1.115 (2)	0.5534 (16)	0.849 (4)	0.063 (9)*

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0508 (4)	0.0276 (3)	0.0642 (5)	-0.0062 (3)	0.0173 (3)	-0.0023 (3)
O1	0.0339 (9)	0.0328 (9)	0.0559 (11)	-0.0003 (7)	0.0190 (8)	-0.0044 (8)
O2	0.0438 (12)	0.0814 (16)	0.0696 (15)	0.0176 (11)	0.0135 (10)	0.0081 (12)
O3	0.0353 (9)	0.0315 (9)	0.0628 (12)	-0.0071 (7)	0.0207 (9)	-0.0071 (8)
O4	0.0525 (12)	0.0327 (10)	0.0694 (14)	-0.0093 (9)	0.0145 (10)	-0.0046 (9)
N1	0.0373 (12)	0.0223 (10)	0.0611 (14)	-0.0008 (9)	0.0158 (10)	-0.0023 (10)
N2	0.0325 (11)	0.0270 (11)	0.0661 (15)	-0.0004 (9)	0.0181 (10)	0.0036 (10)
N3	0.0276 (10)	0.0264 (10)	0.0527 (13)	-0.0003 (8)	0.0127 (9)	0.0009 (9)
C1	0.0330 (14)	0.0291 (13)	0.0715 (19)	-0.0040 (11)	0.0184 (13)	0.0048 (12)
C2	0.0279 (12)	0.0266 (12)	0.0412 (14)	0.0007 (9)	0.0083 (10)	0.0030 (10)
C3	0.0280 (12)	0.0289 (12)	0.0361 (13)	0.0010 (9)	0.0073 (10)	0.0027 (10)

C4	0.0311 (12)	0.0232 (11)	0.0424 (14)	0.0007 (9)	0.0091 (10)	0.0008 (10)
C5	0.0299 (12)	0.0299 (12)	0.0377 (13)	-0.0032 (10)	0.0080 (10)	0.0028 (10)
C6	0.0297 (12)	0.0329 (13)	0.0361 (13)	0.0023 (10)	0.0095 (10)	0.0009 (10)
C7	0.0361 (13)	0.0237 (11)	0.0447 (14)	0.0023 (10)	0.0107 (11)	-0.0030 (10)
C8	0.0299 (12)	0.0263 (12)	0.0394 (13)	-0.0007 (9)	0.0068 (10)	0.0027 (10)
C9	0.0431 (15)	0.0349 (14)	0.0490 (16)	0.0055 (12)	0.0147 (12)	-0.0035 (12)
C10	0.0493 (16)	0.0390 (15)	0.0512 (16)	0.0085 (12)	0.0221 (13)	0.0014 (13)
C11	0.045 (2)	0.152 (4)	0.135 (4)	0.034 (2)	0.019 (2)	-0.013 (4)
C12	0.0350 (14)	0.0305 (13)	0.0636 (18)	-0.0034 (11)	0.0162 (12)	-0.0027 (12)
C13	0.0448 (16)	0.0404 (15)	0.073 (2)	-0.0095 (12)	0.0240 (15)	-0.0085 (14)
C14	0.093 (3)	0.066 (2)	0.077 (3)	-0.032 (2)	0.031 (2)	-0.0167 (19)
C15	0.0338 (13)	0.0260 (12)	0.0403 (14)	0.0051 (10)	0.0125 (11)	0.0077 (10)
C16	0.0385 (14)	0.0353 (14)	0.0443 (15)	0.0027 (11)	0.0079 (11)	0.0015 (11)
C17	0.0542 (17)	0.0441 (16)	0.0466 (16)	0.0070 (13)	0.0167 (13)	-0.0070 (13)
C18	0.0421 (16)	0.0517 (17)	0.0538 (17)	0.0127 (13)	0.0230 (13)	0.0022 (14)
C19	0.0330 (13)	0.0352 (14)	0.0533 (17)	0.0038 (11)	0.0158 (12)	0.0107 (12)
C20	0.0346 (13)	0.0285 (12)	0.0496 (16)	0.0018 (10)	0.0123 (11)	-0.0001 (11)
C21	0.0387 (16)	0.0495 (17)	0.064 (2)	0.0052 (13)	0.0237 (14)	0.0062 (14)
C22	0.0416 (19)	0.077 (2)	0.082 (2)	-0.0098 (17)	0.0233 (17)	-0.0084 (19)

*Geometric parameters (Å, °)*

O1—C6	1.352 (3)	C9—H9B	0.9700
O1—C9	1.441 (3)	C10—H10A	0.9700
O2—C10	1.397 (3)	C10—H10B	0.9700
O2—C11	1.419 (4)	C11—H11A	0.9600
O3—C5	1.361 (3)	C11—H11B	0.9600
O3—C12	1.436 (3)	C11—H11C	0.9600
O4—C13	1.412 (3)	C12—C13	1.507 (3)
O4—C14	1.412 (4)	C12—H12A	0.9700
N1—C1	1.317 (3)	C12—H12B	0.9700
N1—C8	1.378 (3)	C13—H13A	0.9700
N1—HN1	0.8600	C13—H13B	0.9700
N2—C1	1.309 (3)	C14—H14A	0.9600
N2—C2	1.353 (3)	C14—H14B	0.9600
N3—C2	1.337 (3)	C14—H14C	0.9600
N3—C15	1.427 (3)	C15—C16	1.388 (3)
N3—H3	0.8600	C15—C20	1.389 (3)
C1—H1	0.9300	C16—C17	1.382 (4)
C2—C3	1.440 (3)	C16—H16	0.9300
C3—C8	1.400 (3)	C17—C18	1.375 (4)
C3—C4	1.416 (3)	C17—H17	0.9300
C4—C5	1.368 (3)	C18—C19	1.386 (4)
C4—H4	0.9300	C18—H18	0.9300
C5—C6	1.429 (3)	C19—C20	1.399 (3)
C6—C7	1.370 (3)	C19—C21	1.444 (4)
C7—C8	1.397 (3)	C20—H20	0.9300
C7—H7	0.9300	C21—C22	1.170 (4)
C9—C10	1.491 (3)	C22—H22	0.95 (3)

## supplementary materials

---

C9—H9A	0.9700		
C6—O1—C9	117.09 (19)	H10A—C10—H10B	108.3
C10—O2—C11	111.7 (3)	O2—C11—H11A	109.5
C5—O3—C12	116.48 (18)	O2—C11—H11B	109.5
C13—O4—C14	112.8 (2)	H11A—C11—H11B	109.5
C1—N1—C8	120.4 (2)	O2—C11—H11C	109.5
C1—N1—HN1	119.8	H11A—C11—H11C	109.5
C8—N1—HN1	119.8	H11B—C11—H11C	109.5
C1—N2—C2	117.6 (2)	O3—C12—C13	106.5 (2)
C2—N3—C15	127.4 (2)	O3—C12—H12A	110.4
C2—N3—H3	116.3	C13—C12—H12A	110.4
C15—N3—H3	116.3	O3—C12—H12B	110.4
N2—C1—N1	125.4 (2)	C13—C12—H12B	110.4
N2—C1—H1	117.3	H12A—C12—H12B	108.6
N1—C1—H1	117.3	O4—C13—C12	111.1 (2)
N3—C2—N2	117.4 (2)	O4—C13—H13A	109.4
N3—C2—C3	121.2 (2)	C12—C13—H13A	109.4
N2—C2—C3	121.4 (2)	O4—C13—H13B	109.4
C8—C3—C4	117.6 (2)	C12—C13—H13B	109.4
C8—C3—C2	116.7 (2)	H13A—C13—H13B	108.0
C4—C3—C2	125.7 (2)	O4—C14—H14A	109.5
C5—C4—C3	120.7 (2)	O4—C14—H14B	109.5
C5—C4—H4	119.7	H14A—C14—H14B	109.5
C3—C4—H4	119.7	O4—C14—H14C	109.5
O3—C5—C4	125.2 (2)	H14A—C14—H14C	109.5
O3—C5—C6	114.5 (2)	H14B—C14—H14C	109.5
C4—C5—C6	120.3 (2)	C16—C15—C20	120.1 (2)
O1—C6—C7	124.4 (2)	C16—C15—N3	117.1 (2)
O1—C6—C5	115.7 (2)	C20—C15—N3	122.8 (2)
C7—C6—C5	119.9 (2)	C17—C16—C15	119.7 (2)
C6—C7—C8	119.2 (2)	C17—C16—H16	120.1
C6—C7—H7	120.4	C15—C16—H16	120.1
C8—C7—H7	120.4	C18—C17—C16	120.6 (3)
N1—C8—C7	119.4 (2)	C18—C17—H17	119.7
N1—C8—C3	118.3 (2)	C16—C17—H17	119.7
C7—C8—C3	122.2 (2)	C17—C18—C19	120.2 (2)
O1—C9—C10	108.1 (2)	C17—C18—H18	119.9
O1—C9—H9A	110.1	C19—C18—H18	119.9
C10—C9—H9A	110.1	C18—C19—C20	119.6 (2)
O1—C9—H9B	110.1	C18—C19—C21	120.0 (2)
C10—C9—H9B	110.1	C20—C19—C21	120.4 (2)
H9A—C9—H9B	108.4	C15—C20—C19	119.6 (2)
O2—C10—C9	108.7 (2)	C15—C20—H20	120.2
O2—C10—H10A	109.9	C19—C20—H20	120.2
C9—C10—H10A	109.9	C22—C21—C19	177.3 (3)
O2—C10—H10B	109.9	C21—C22—H22	178 (2)
C9—C10—H10B	109.9		
C2—N2—C1—N1	4.2 (4)	C1—N1—C8—C3	-1.7 (4)

C8—N1—C1—N2	−2.7 (4)	C6—C7—C8—N1	177.9 (2)
C15—N3—C2—N2	1.5 (4)	C6—C7—C8—C3	−2.4 (4)
C15—N3—C2—C3	−178.7 (2)	C4—C3—C8—N1	−177.0 (2)
C1—N2—C2—N3	178.4 (2)	C2—C3—C8—N1	4.1 (3)
C1—N2—C2—C3	−1.3 (4)	C4—C3—C8—C7	3.4 (4)
N3—C2—C3—C8	177.5 (2)	C2—C3—C8—C7	−175.5 (2)
N2—C2—C3—C8	−2.7 (3)	C6—O1—C9—C10	176.3 (2)
N3—C2—C3—C4	−1.2 (4)	C11—O2—C10—C9	−177.2 (3)
N2—C2—C3—C4	178.5 (2)	O1—C9—C10—O2	−76.9 (3)
C8—C3—C4—C5	−1.2 (3)	C5—O3—C12—C13	−170.3 (2)
C2—C3—C4—C5	177.5 (2)	C14—O4—C13—C12	−79.6 (3)
C12—O3—C5—C4	−11.3 (3)	O3—C12—C13—O4	−179.3 (2)
C12—O3—C5—C6	168.7 (2)	C2—N3—C15—C16	−146.2 (2)
C3—C4—C5—O3	178.3 (2)	C2—N3—C15—C20	35.6 (4)
C3—C4—C5—C6	−1.7 (4)	C20—C15—C16—C17	−1.7 (4)
C9—O1—C6—C7	−8.6 (3)	N3—C15—C16—C17	−179.9 (2)
C9—O1—C6—C5	172.4 (2)	C15—C16—C17—C18	−0.1 (4)
O3—C5—C6—O1	1.8 (3)	C16—C17—C18—C19	1.1 (4)
C4—C5—C6—O1	−178.3 (2)	C17—C18—C19—C20	−0.5 (4)
O3—C5—C6—C7	−177.3 (2)	C17—C18—C19—C21	179.5 (3)
C4—C5—C6—C7	2.7 (4)	C16—C15—C20—C19	2.4 (4)
O1—C6—C7—C8	−179.6 (2)	N3—C15—C20—C19	−179.6 (2)
C5—C6—C7—C8	−0.6 (4)	C18—C19—C20—C15	−1.3 (4)
C1—N1—C8—C7	177.9 (2)	C21—C19—C20—C15	178.7 (2)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N3—H3···Cl1	0.86	2.46	3.277 (2)	160
N1—HN1···C11 <sup>i</sup>	0.86	2.23	3.066 (2)	165
C1—H1···O4 <sup>i</sup>	0.93	2.46	3.372 (3)	167

Symmetry codes: (i)  $-x+1, y+1/2, -z+3/2$ .

## supplementary materials

---

Fig. 1




Fig. 2

