

Atf6 α impacts cell number by influencing survival, death and proliferation



Rohit B. Sharma, Jarin T. Snyder, Laura C. Alonso*

ABSTRACT

Background: A growing body of literature suggests the cell—intrinsic activity of Atf6 α during ER stress responses has implications for tissue cell number during growth and development, as well as in adult biology and tumorigenesis [1]. This concept is important, linking the cellular processes of secretory protein synthesis and endoplasmic reticulum stress response with functional tissue capacity and organ size. However, the field contains conflicting observations, especially notable in secretory cell types like the pancreatic beta cell.

Scope of review: Here we summarize current knowledge of the basic biology of Atf6 α , along with the pleiotropic roles Atf6 α plays in cell life and death decisions and possible explanations for conflicting observations. We include studies investigating the roles of Atf6 α in cell survival, death and proliferation using well-controlled methodology and specific validated outcome measures, with a focus on endocrine and metabolic tissues when information was available.

Major conclusions: The net outcome of Atf6 α on cell survival and cell death depends on cell type and growth conditions, the presence and degree of ER stress, and the duration and intensity of Atf6 α activation. It is unquestioned that Atf6 α activity influences the cell fate decision between survival and death, although opposite directions of this outcome are reported in different contexts. Atf6 α can also trigger cell cycle activity to expand tissue cell number through proliferation. Much work remains to be done to clarify the many gaps in understanding in this important emerging field.

© 2019 Published by Elsevier GmbH. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords Activating transcription factor 6; Pancreatic beta cell; Cell survival; Apoptosis; Replication

1. INTRODUCTION

The mammalian unfolded protein response (UPR) is an elegant cellular process originating in the endoplasmic reticulum (ER) which adapts ER protein folding capacity to meet protein folding load [2,3]. The ER is a critical multifunctional cellular organelle with roles including secretory protein synthesis, folding, quality control and targeting; calcium homeostasis; and glucose and lipid metabolism. Like many other biological systems, ER mass and function are actively determined by the relative rates of catabolic destruction and de novo synthesis [2,3].

The UPR, triggered by ER stressors such as excess unfolded proteins, redox imbalance, or calcium depletion, is the principal regulator of ER expansion. The biology of the UPR, which is conserved across phyla, has been extensively reviewed [2–6]. Three ER transmembrane proteins, Perk, Ire1 and Atf6 α , respond to ER stress and activate a coordinated translational and transcriptional program to slow new peptide entry into the ER, enhance ER associated degradation of misfolded protein, and increase biosynthesis of ER components to expand ER capacity. If the adaptation is successful, the cell resumes function at a new higher capacity.

However, if adaptation fails, unresolvable ER stress and chronic UPR activation lead to cell death [7].

The original understanding of the UPR was as a cell-autonomous mechanism by which a cell adapts secretory-pathway protein production capacity to demand. However, emerging evidence suggests important roles for the UPR at the level of complex tissues and even whole organism health and metabolism, by influencing tissue function and cell number through death, survival, and proliferation decisions [5,6]. Of the three ER transmembrane proteins sensing stress and initiating the UPR, Atf6 α may be the least well understood. Roles of Atf6 α in organogenesis and development have been reviewed [1]. Here we summarize current understanding of the roles played by Atf6 α in influencing cell proliferation and death, with attention paid to conflicting conclusions and possible explanations. When possible, we focus on data generated in primary tissues and cancers.

2. OVERVIEW OF ATF6 α BIOLOGY

Atf6 α was identified as an ER stress response mediator in 1998 in the laboratory of Kazutoshi Mori in Kyoto, Japan, using a yeast-one-hybrid

Diabetes Center of Excellence, Department of Medicine, University of Massachusetts Medical School, Worcester, MA, USA

*Corresponding author. AS7-2047, Division of Diabetes, 368 Plantation Street, Worcester, MA, 01605, USA. Fax: +508 856 3803. E-mail: Laura.Alonso@umassmed.edu (L.C. Alonso).

Abbreviations: Atf6 α , Activating transcription factor alpha; Bcl-2, B-cell lymphoma 2; cFos, human homolog of Finkel–Biskis–Jenkins murine osteogenic sarcoma virus oncogene; Chop, C/EBP homologous protein; ER, endoplasmic reticulum; ERAD, ER associated degradation; Jnk, c-Jun N-terminal Kinase; mTor, mammalian target of rapamycin; Rheb, Ras homolog, mTor binding; Runx2, Runt-related transcription factor 2; S1P, site 1 protease; S2P, site 2 protease; Vegf, Vascular endothelial growth factor

<https://doi.org/10.1016/j.molmet.2019.06.005>

screen with a promoter element found to be common to genes upregulated by the UPR [8]. Initially found to be an activating transcription factor (ATF) family member that weakly bound a cAMP response element [9] and an SRF-binding protein [10], Atf6 α is now known to be a member of a family of stress-responsive bZip transcription factors called OASIS factors [11]. The pace of Atf6 α -related discovery has accelerated recently; most papers on the function of this important factor have been published in the past 5 years.

2.1. ATF6 α gene structure

The human *ATF6 α* gene, conserved across plants and metazoans [12], is located on chromosome 1. *ATF6 α* contains only a single protein-coding transcript, with 16 exons and no known splice isoforms, which codes for the 670aa ATF6 α protein (Figure 1). The mouse *Atf6 α* gene is also encoded on chromosome 1, also contains 16 exons with no known splice variants, and encodes a 656aa protein.

2.2. ATF6 α protein size and intracellular localization

In unstressed human cells, ATF6 α protein is detected at a 90 kDa size, larger than the calculated molecular weight of 74.6 kDa [8]. However, during ER stress conditions, such as exposure to the glycosylation inhibitor tunicamycin or SERCA inhibitor thapsigargin, ATF6 α -directed antibodies detect an additional 50 kDa band [8,13]. Indirect immunofluorescence, cellular fractionation and tryptic digestion studies determined that p90ATF6 α is a type II transmembrane ER resident glycoprotein with a single-pass hydrophobic transmembrane domain of 21 aa near the middle of the protein [13,14]. In contrast, p50ATF6 α is a soluble, short-lived nuclear protein that is most easily detected in the presence of the ALLN protease inhibitor [13–16]. Studies with mutant constructs suggest that the presence or absence of the transmembrane region determines the localization of the N-terminal (p50ATF6 α , aa 1–373) domain, suggesting a model in which a cleavage event releases the cytosolic domain from the transmembrane and luminal domains, resulting in nuclear localization [13]. The structural elements of ATF6 α are summarized in Figure 1.

2.3. p90ATF6 α transit to Golgi during ER stress

Although full length p90ATF6 α contains two Golgi localization signals [17], during normal conditions it is found in the ER [13,14]. The 272aa ER luminal domain is sufficient to sense stress, relocate the protein to

the Golgi, and allow cleavage [18]. Stress sensing occurs through interaction with the Grp78 (also called BiP, Hspa5) chaperone [17]. In unstressed conditions ATF6 α is retained in the ER by interaction with Grp78, but during stress Grp78 is titrated away, releasing ATF6 α to proceed to the Golgi in a COPII dependent mechanism [19]. Transport is regulated; even with excess unfolded proteins in the ER, ATF6 α is selectively allowed to move to the Golgi while unfolded proteins are retained in the ER [20]. In a pancreatic beta cell line, ATF6 α translocation to Golgi required Sar1A, a small GTPase involved in COPII vesicle formation [21].

2.4. Mechanism of p90ATF6 α cleavage in the Golgi

The mechanism of the ATF6 α cleavage event was clarified by Brown and Goldstein [15]. Similarly to SREBP, p90ATF6 α undergoes regulated intramembrane proteolysis, in which S1P first cleaves the peptide on the luminal side of the transmembrane domain, and then S2P cleaves the peptide within the transmembrane domain, liberating the cytoplasmic soluble domain [15]. In support of this model, a serine protease inhibitor had no impact on transit of p90ATF6 α to the Golgi, but prevented ATF6 α cleavage and target activation [22]. A novel feedback loop has been described, in which ATF6 α transcriptional target and key Golgi calcium regulator Nucleobindin 1 inhibits S1P ATF6 α cleavage, without impacting transit to Golgi [23]. In neurons, Calsenilin, encoded by the *KCNIP3* gene, regulates ATF6 α processing and mediates protection by repaglinide on Huntington's disease [24].

2.5. Protein modifications impact ATF6 α behavior

Post-translational modifications also impact ATF6 α transport and activation by ER stress. Under unstressed conditions ATF6 α is found in monomeric, dimeric and multimeric complexes due to intra- and intermolecular disulfide bonds [25]. Only reduced monomeric ATF6 α transits to Golgi, but reduction was not sufficient to induce activation [25]. An siRNA screen for ATF6 α activators in cancer cells identified a novel mechanism in which the protein disulfide isomerase Pdia5 promotes a disulfide bond rearrangement in the luminal domain of ATF6 α , leading to ATF6 α packaging in CopII vesicles [26]. Intriguingly, the ATF6 α luminal domain has three conserved N-linked glycosylation sites, and glycosylation provides a negative signal, restricting ATF6 α responsiveness to stress [27]. Under-glycosylated forms of ATF6 α achieved by mutation of the glycosylation sites leads to reduced interaction with ER




Figure 1: Schematic of the ATF6 α protein, with known domain structures and functional elements. nATF6, nuclear ATF6. S1P, site 1 protease. S2P, site 2 protease. VP16, herpes simplex virus protein vsm65 activating domain. Data contributing to this figure are from [13,15–17,25,51].

chaperone calreticulin and increased transit to Golgi, cleavage, and transactivation of target genes [27]. The role of stress kinases is controversial; one study found that p38-Mapk phosphorylated Atf6 α at Thr-166, preventing cleavage and nuclear translocation [28]; in other conditions p38-Mapk phosphorylation of the N-terminal cytoplasmic domain was found to enhance Atf6 α transcriptional activation of targets such as Grp78 [29,30].

2.6. Atf6 α abundance is also regulated at the RNA level

In addition to the well-known protein-level activation by ER stress described above, Atf6 α is also itself increased at the RNA level by ER stress. In MEFs, Atf6 α mRNA was increased by tunicamycin exposure in sXbp dependent fashion [31]. On the other hand, thapsigargin was observed to increase Atf6 α mRNA in a feed-forward autoregulation loop via Atf6 α binding elements in the Atf6 α promoter [32]. The proteasome-regulating transcription factor Nrf1, and ERR γ , also positively regulate Atf6 α gene transcription through direct enhancer binding [33,34].

2.7. Nuclear p50Atf6 α regulates gene transcription through direct DNA binding

Cleavage of Atf6 α in the Golgi releases p50Atf6 α , a 373aa cytoplasmic fragment which contains a basic leucine zipper DNA binding domain and several putative nuclear localization signals, and localizes to the nucleus [8,13–15]. As such, ER stress-induced transit and cleavage of Atf6 α activates an ER-to-nucleus transcriptional regulatory program. The first identified DNA regulatory sequence bound by Atf6 α was a 19 nucleotide motif, CCAAT₉CCACG, called the ER stress response element (ERSE) [8]. This bipartite motif contains a CCAAT box, bound by general factors, and a CCACG box, bound by Atf6 α [8,35,36]. Subsequently, Atf6 α was found to also bind to TGACGTG(G), now called the UPRE, which contains a partial complement of the ERSE sequence [36]; however, the UPRE is now considered to be more responsive to sXbp1 than Atf6 α [37]. A third ER stress response element, ERSE-II, ATTGG-N-CCACG, contains the Atf6 α -binding CCACG box next to an inverted CCAAT box with a much shorter spacer of only one base pair [37].

ERSE motifs are found in known ER stress responsive genes Grp78, Grp94 and calreticulin, and mediate stress-responsive transcription of a luciferase construct [8]. Base-by-base mutation mapped the ERSE critical nucleotides [8]. The bipartite ERSE motif engages general transcription factors YY1 and NF-Y/CBP at the CCAAT box; Atf6 α binds at the CCACG box [8,35,36]. Atf6 α interacts with the C-subunit of NF-Y [38]; co-binding of NF-Y and YY1 added selectivity and strength to the Atf6 α transcriptional response [14]. In addition to SRF, NF-Y and YY1, Atf6 α may regulate gene transcription by interacting with PGC1alpha [34]. Mutation of the bZip DNA binding domain eliminated transactivation of a Grp78-ERSE luciferase reporter [16]. Atf6 α (1–373) transcription transactivating activity is mostly contained in a VP16-like domain at aa 1–38 and 52–93 [16].

2.8. Cross-talk between Atf6 α and other UPR pathways

Teasing out the role of Atf6 α in the UPR transcriptional response is complicated by overlapping functions of other UPR nuclear effectors, especially sXbp1 [39]. In general, the UPRE motif is activated by sXbp1, whereas ERSE and ERSE-II are activated by both sXbp1 and Atf6 α [37]. In the presence of NF-Y, sXbp1 can replace Atf6 α at the ERSE, but with lower binding efficiency [37]. Complicating matters, Atf6 α transcriptionally induces Xbp1 [40], but may suppress Ire1 mRNA levels [41]. Further confounding separation of roles, Atf6 α heterodimerizes with sXbp1, binding the UPRE with 8-fold higher affinity than sXbp1 homodimer [42].

Atf6 α has a homologue, Atf6 β , that shares similar biology: full length Atf6 β is ER membrane localized and upon ER stress transits to Golgi, undergoes cleavage by S1P/S2P and releases an N-terminal transcription factor [43]. Atf6 β exhibits some degree of functional redundancy with Atf6 α , since gene deletion of either Atf6 α or Atf6 β has minimal biological impact but deletion of both Atf6 α and Atf6 β is embryonic lethal [42]. However, in vitro studies show important differences. Atf6 α is solely responsible for the classical UPR-dependent gene regulation [42,44]; in fact, Atf6 β has been reported to antagonize some Atf6 α actions, such as transcriptional induction of Grp78 [45,46].

Atf6 α also has a complex relationship with UPR death effector Chop. Atf6 α transcriptionally induces *Chop* mRNA [47,48]. However, overexpression of Chop suppressed Atf6 α activation of target gene Grp78 in dose-dependent manner. Chop is also a bZip transcription factor and may heterodimerize with Atf6 α to suppress its activity [49]. On the other hand, CHIP assay showed that Chop binds to the Grp78 promoter, and a mutant Chop defective in DNA binding failed to suppress Atf6 α mediated Grp78 induction, favoring a model in which Chop inhibits Atf6 α gene regulation by competing for the regulatory motifs [49].

2.9. Canonical function of Atf6 α

The principal outcome of Atf6 α activation during ER stress is the expansion of functional ER capacity. Atf6 α transcriptionally upregulates many genes involved in protein folding, including ER-resident chaperones, foldases, calcium transport proteins, and oxidation/reduction regulators [1,39,42]. Atf6 α is also required for optimal clearance of misfolded proteins via Endoplasmic Reticulum-Associated Degradation (ERAD) genes and for expansion of the ER through induction of membrane synthesis [1,50]. Atf6 α promotes ER expansion via heavily redundant and overlapping roles with the other UPR pathways, such that disentangling one from the others is challenging [2,4,39].

2.10. Mechanisms turning the Atf6 α signal off

Full length p90Atf6 α is subject to ubiquitination and proteasomal degradation, more pronounced after ER stress induction [51]. WFS1, an ER transmembrane protein mutated in Wolfram syndrome, was found to suppress Atf6 α activation by inducing proteasome-mediated degradation of full-length Atf6 α [52]. Supporting a role for chronic activation of Atf6 α in Wolfram syndrome, Atf6 α levels were increased in both Wfs1-deleted mice and Wolfram patient samples [52]. In addition, p90Atf6 α was identified as a transmembrane target of ERAD, requiring both ERAD E3 ligase Sel1l and mannose trimming [50]. Nuclear Atf6 α is short-lived, requiring protease inhibition to detect, suggesting a rapid degradation mechanism [13–16]. Observing that protein stability of N-terminal Atf6 α mutants was correlated with transcriptional activity, the VP16-homologous domain identified by Christopher Glembotski's group was found to confer both transactivation and degradation capacity [16]. Intriguingly, the unspliced form of Xbp1 mRNA is also translated to a protein product, which accumulates in later stages of UPR recovery and may be responsible for targeting sXbp1 and nAtf6 α for proteasome mediated degradation, implicating an Ire1-derived off-signal in shutting down the sXbp1 and nAtf6 α transcriptional programs [53].

3. CONFLICTING ROLES: ATF6 α CAN PROMOTE BOTH CELL SURVIVAL AND CELL DEATH

Atf6 α has been reported to play numerous, disparate roles in processes regulating cell number (Figure 2). In its canonical role, Atf6 α

activation drives a multi-pronged and robust effort by the cell to restore protein folding capacity, export misfolded protein from the ER for degradation by the proteasome, and ultimately, promote cell survival in response to diverse insults to the ER folding environment [44]. This pro-survival function may drive the pathology of achromatopsia, a genetic human retinal disease resulting from *ATF6 α* mutations [54]. Mutations that impair ATF6 α activation by disrupting ER-Golgi trafficking, regulated intramembrane proteolytic cleavage or transcriptional activity were found to increase cell death in patient-derived fibroblasts, although a role for cell death in achromatopsia pathogenesis remains uncertain [54].

On the other hand, ATF6 α has also been found to increase cell death, through direct and indirect mechanisms, supporting a pro-apoptotic role for ATF6 α under some conditions. ATF6 α induces expression of Chop, which drives apoptosis through various mechanisms. A tissue highlighting the complicated coexisting pro- and anti-apoptotic functions of ATF6 α is the pancreatic beta cell. For example, pathogenesis of Wolfram Syndrome, a progressive neurological syndrome characterized by hearing loss, optic atrophy and diabetes, may be due to ATF6 α toxicity [52]. WFS1 provides feedback inhibition on ATF6 α signaling, via targeting full length ER ATF6 α for proteasomal degradation by recruiting the HRD1 E3 ligase [52]. Loss of function of WFS1 in Wolfram Syndrome causes pancreatic beta cell dysfunction, apoptosis and diabetes through dysregulated excess ATF6 α activity [52]. ATF6 α was also implicated in the transcriptional response to lipotoxicity leading to human beta cell death [55]. On the other hand, loss of ATF6 α was reported to cause beta cell death in a type 1 diabetes model, and cell death was decreased by restoration of ATF6 α expression [56]. Known mechanisms of pleiotropic pro-survival and/or pro-cell death ATF6 α functions are reviewed below.

4. ATF6 α PROMOTES CELL SURVIVAL

4.1. ATF6 α promotes cell survival by adapting protein folding capacity during ER stress

As noted above, the primary function of ATF6 α is as an ER membrane sensor, detecting misfolded proteins in the ER lumen and activating a transcriptional response to restore protein folding homeostasis. Efficient ER function requires the activity of many genes that ATF6 α controls, including protein folding chaperones, foldases, [Ca²⁺]-regulatory and -regulated ER proteins, redox regulators, as well as other miscellaneous genes with unclear roles in the UPR [2,4,13,14,57,58]. One of the earliest studies of ATF6 α revealed that ER Ca²⁺-ATPase inhibitor thapsigargin activates cleavage of ATF6 α to induce protein folding chaperones [13]. Overexpressed ATF6 α cooperates with general transcription factors NF-Y or YY1 to bind the ERSE and induce expression of ER chaperones GRP78, GRP94, and calreticulin, among many other genes later identified [14,47]. Conversely, deletion of ATF6 α in MEFs severely reduced the ER-stress dependent induction of a suite of ER chaperone genes: Grp78, Grp94, Grp170, p58IPK, and Erdj3 [57]. In addition to chaperones, ATF6 α also increases membrane phosphatidylcholine synthesis which may help to increase folding capacity by increasing ER volume [59].

Some ATF6 α -dependent chaperones are critically important for cell survival; chaperone deletion may induce apoptosis despite activation of UPR [60–62]. In addition to chaperone, calcium and UPR-inhibitory actions, Grp78 directly binds and inhibits ER-resident Bik pro-apoptotic activity, preventing binding of Bcl2 [63,64]. Homozygous loss of chaperone expression can cause embryonic lethality or severe developmental defects. For example, deletion of Grp78 is lethal very early in development, around the time of embryo implantation into the




Figure 2: Summary of ATF6 α -regulated pathways regulating cell fate decisions leading to cell death, cell survival or proliferation. S1P, site 1 protease. S2P, site 2 protease. Chop, C/EBP homologous protein. Jnk, c-Jun N-terminal kinase. Runx2, Runt-related transcription factor 2. Bcl-2, B-cell lymphoma 2. Vegf, Vascular endothelial growth factor. cFos, human homolog of Finkel–Biskis–Jinkins murine osteogenic sarcoma virus oncogene. Rheb, Ras homolog, mTor binding. ERAD, ER associated degradation. mTor, mammalian target of rapamycin. Please see the text for citations.

uterine wall, with excess apoptosis in the inner cell mass [60]. Loss of function in chaperone p58IPK causes diabetes and neurodegenerative disorders in mice and is linked to a similar syndrome in rare familial cases [65]. Intriguingly, in addition to ER-resident chaperone expression, Atf6 α may also control extracellular protein folding via increased transcription and secretion of chaperone ERdj3, which functions both intracellularly and extracellularly to prevent aggregation of unfolded proteins [66]. Overexpressed ERdj3 reduced extracellular amyloid beta aggregation, and secreted ERdj3 in conditioned media prevented vacuolization in a neuroblastoma line treated with toxic prion protein. As such, Atf6 α -mediated induction of chaperones may be important for both intracellular and extracellular proteostasis.

4.2. Atf6 α influences ER redox status

Atf6 α pro-survival functions also include maintenance of healthy ER redox status. Atf6 α controls the expression of several ER-resident proteins that catalyze the formation and breakage of disulfide bonds during protein folding [67], such as Erp72, p5, & Ero1beta [42,57]. These oxidoreductases are oxygen-dependent enzymes that may link hypoxia to UPR activation [68]. Atf6 α has a well-supported role in promoting adaptation to hypoxic stress. Atf6 α knock-in reduced necrosis and apoptosis and improved heart function after ischemia/reperfusion injury [69]. Conversely DN-Atf6 α or siAtf6 α increased apoptosis in response to ischemia/reperfusion in cardiac myocytes [70]. Pdia6 is an Atf6 α target that promotes chemotherapeutic resistance in cancer and protects against ischemia/reperfusion injury in cardiomyocytes when overexpressed [71,72]. Atf6 α -dependent induction of antioxidant genes has also been reported to be protective in cardiac ischemic/reperfusion injury [73] but an antioxidant function of Atf6 α has not been confirmed in other studies. Besides the heart, Atf6 α also appears protective against ischemia/reperfusion injury in the brain. In a surgically-induced murine stroke model, overexpressed Atf6 α reduced infarct size and improved cognition, associated with upregulated Bcl2 and accelerated induction of autophagy [74]. Interestingly, the usually pro-apoptotic Atf6 α target Chop may actually be protective in hypoxic neurons [75].

4.3. Atf6 α targets improve clearance of ER misfolded proteins

Atf6 α also promotes cell survival by transcriptionally activating ER-associated degradation pathways (ERAD) that clear misfolded proteins from the ER. Atf6 α and Xbp1 knockout MEFs show an inability to induce canonical ERAD genes HerpUD1, Edem, and Hrd1 in response to ER stress, along with a decrease in viability compared to wild type cells [42]. ATF6 α also induces Derlin-3, an important ERAD component that facilitates extrusion of misfolded proteins into the cytosol for proteasomal degradation [76,77]. Derlin-3 improved clearance of misfolded protein from the ER and was necessary and sufficient to reduce apoptosis in rat cardiomyocytes that underwent simulated ischemia/reperfusion injury [76].

4.4. Atf6 α influences autophagy

ATF6 α may also improve cell survival by inducing autophagy, clearing misfolded proteins and damaged organelles. As noted in 4.2, overexpression of nuclear Atf6 α protected mice against stroke, with early induction of autophagy [74]. Orml3 knockout splenic B cells had decreased Atf6 α and Beclin1 expression, with reduced survival; overexpression of ATF6 α rescued Beclin1 expression, autophagy, and survival [78]. Atf6 α can also interact with C/EBP-beta to trigger autophagy through induction of Dapk1 in response to bacterial insult or IFN γ . In support of the importance of this observation, Atf6 α KO mice are susceptible to death from bacterial infection [79]. Phosphorylation

of Atf6 α by Ask1/Mkk3-p38Mapk pathway was necessary for its activation and subsequent interaction with C/EBP-beta to mediate IFN γ induced Dapk1 [28]. Similarly, in the mouse neuroblastoma line Neuro2, Japanese encephalitis virus infection induced Atf6 α and Xbp1-dependent autophagy, preventing apoptosis [80]. In this model system Xbp1, but not Atf6 α , was necessary for Beclin-1 induction, while Atg3 was Atf6 α -dependent. An intriguing hypothesis is that Atf6 α could have evolved in part as a defense mechanism against pathogen-induced ER stress. On the other hand, ER-stress induced autophagy was unaffected by knockdown of Atf6 α in the human neuroblastoma line SK-N-SH but was instead dependent on the Ire1-Jnk pathway [81], and siAtf6 α had no effect on Dengue-virus induced autophagy [82].

4.5. Atf6 α may affect survival through activation of mTor

Atf6 α was required for mTor activation in a *Drosophila* cell line [83]. Atf6 α promoted chemotherapeutic resistance of dormant squamous carcinoma through Akt-independent mTor activation via transcriptionally inducing mTor-activator Rheb [84]. Furthermore, in endothelial cells, Vegf induced Atf6 α was required for pro-survival Akt phosphorylation by mTor [85].

4.6. Atf6 α has a limited role under unstressed conditions

Surprisingly, considering the implied importance of this supposedly critical UPR component, under unstressed conditions Atf6 α -null mice are remarkably normal. Individual Atf6 α or Atf6 β -null mice appear viable and healthy, but deletion of both genes results in embryonic lethality, suggesting that Atf6 α and Atf6 β have redundant functions such that Atf6 β can replace Atf6 α under basal conditions [42,44]. One cell type that may require Atf6 α under unstressed conditions is the pancreatic beta cell; in normoglycemic conditions, anti-Atf6 α siRNA-treated insulinoma cells had a JNK-dependent increase in apoptosis [86]. Although Atf6 α knockdown in these cells reduced Atf6 α protein by only around 60%, this reduction was sufficient to reduce Grp78 mRNA and protein expression under basal conditions. Other Atf6 α targets Grp94 and ERAD component Herp were only blunted during induction of ER stress. On the other hand, whole body Atf6 α -null mice had no discernible impairment in glucose metabolism except under insulin-demand stress conditions [87].

4.7. Atf6 α is required for resilience in the face of stress

When exposed to stress conditions, Atf6 α -null cells have impaired stress resistance. Although Atf6 α knockout MEFs had no significant difference in chaperone expression or viability when cultured in normal growth medium, induction of chaperones was blunted after treatment with ER stress-inducing drugs, and viability was decreased [44]. Atf6 α is also critical for the acute adaptive response to ER stress in vivo. Challenging Atf6 α -null mice with intraperitoneal tunicamycin resulted in macroscopic liver damage and a dramatic 80% mortality (compared to 0% in wild type mice), along with reduced and delayed induction of ER-chaperone and ERAD genes. Pancreatic beta cells in Atf6 α -null mice also showed impaired function under insulin demand stress [87]. Taken together, it seems the impact of Atf6 α depletion requires acute stress to be revealed [44].

In contrast, it has been reported that Atf6 α is not required for induction of UPR genes by ER stress in stable anti-Atf6 α siRNA-expressing MEFs [31]. The authors postulated that transcription factors related to Atf6 α (Atf6 β or other family members) may compensate for Atf6 α loss. Although the authors reported that residual Atf6 α activity was unlikely, due to undetectable Atf6 α expression in the Atf6 α knockdown cells, nuclear Atf6 α is a potent and short-lived transcription factor [16].

Therefore, it is possible that some Atf6 α may have been present but not detected. This is consistent with another study using Atf6 α siRNA in liver Kupffer cells that showed Trn-induced upregulation of Xbp1, Chop, and Grp78 was unaffected by Atf6 α KD in which p50-Atf6 α was decreased but still detectable after knockdown [88].

5. IN OTHER CONTEXTS, ATF6 α INCREASES APOPTOSIS

Contradictory to its reported pro-survival role, in certain contexts Atf6 α has been shown to activate the intrinsic mitochondrial apoptosis pathway, possibly related to supra-physiological Atf6 α activation. Ectopically expressed Atf6 α directly bound an ERSE in the Bcl-2 promoter to repress its expression [89]. Overexpressed Atf6 α decreased viability of vascular endothelial cells exposed to thapsigargin [90]. Atf6 α may also regulate intrinsic apoptosis through downregulation of E2f1 expression, the loss of which was sufficient to increase expression of Puma and Noxa [91]. In mouse granulosa cells, knockdown of Atf6 α decreased p53 and apoptosis [92]. In contrast, knockdown of Xbp1 increased apoptosis [93]. Overexpression of nuclear Atf6 α in differentiating myoblasts upregulated a WW-domain binding protein (Wbp1) and downregulated anti-apoptotic Mcl-1, which was sufficient to increase apoptosis [94].

5.1. Atf6 α increases expression of Chop

The most widely supported pro-death signal downstream of Atf6 α is the transcription factor Chop (Ddit3, Gadd153). Atf6 α cooperates with the Perk/Atf4 pathway to induce maximal expression of Chop in response to ER stress [95]. Chop knockout MEFs have delayed apoptosis after exposure to tunicamycin [96]. Similarly, tunicamycin-induced apoptosis was reduced, but not eliminated, in the kidney proximal tubular epithelium of Chop-null mice. Chop loss of function is protective in models of diabetes [97–100], neurodegenerative disease [101,102], renal injury [103–107], and sepsis [108].

Chop induces cell death through several mechanisms. Like Atf6 α , Chop directly controls expression of components of the intrinsic apoptosis pathway. It is reported to downregulate pro-survival Bcl2 [109] and increase expression of pro-apoptotic Puma and Bim [110]. Via induction of Gadd34, Chop antagonizes the p-eIF2 α -mediated translation block to increase protein synthesis even though the protein folding capacity may not have been adequately restored [103,111]. Furthermore, Ero1 α is activated by Chop and contributes to oxidative stress in the ER which sensitizes cells to undergo apoptosis [103,111]. The Atf4/Chop pathway could define a switch from anti-apoptotic to pro-apoptotic signaling with longer stress duration, with the short half-life of pro-apoptotic mRNAs providing protection against cell death early during adaptive UPR [112].

5.2. Atf6 α -induced autophagy promotes cell death

As described above, Atf6 α -dependent autophagy can be protective in some cases but leads to autophagic cell death in other contexts. In MCF7 cells, siRNA silencing of ATF6 α , IRE1, or dnPERK, inhibited doxorubicin-induced autophagy and apoptosis [113]. Knockdown of ATF6 α also prevented berberine-induced elevation of GRP78 in cancer cell lines, which was required for induction of autophagy and cell death [114].

5.3. Atf6 α leads to inflammation

In addition to the well-studied effects on Chop and intrinsic apoptosis pathways, Atf6 α also modulates inflammatory and immunogenic cell death. Atf6 α promoted NF- κ B activation and pro-inflammatory cytokine expression in liver Kupffer cells to promote liver damage during

ischemia/reperfusion injury [88]. On the other hand, Atf6 α -induced sXbp1 prevents Tnf- α induced apoptosis in osteoarthritic cartilage [115], and Atf6 α was necessary for suppression of Tnf- α -induced NF- κ B activation via upregulation of C/EBP β and activation of mTor, which prevented Akt phosphorylation [116].

5.4. How can Atf6 α be both pro- and anti-cell survival?

The many seemingly contradictory findings with regard to the downstream effects of Atf6 α may have multiple explanations. Heterodimerization with other transcription factors could provide cell-type specificity for Atf6 α target genes, leading to different results in different cell types. Post-translational modifications may also provide context-specificity of Atf6 α activity. Cell-type specific epigenetic landscapes could prime or block activation of Atf6 α target genes. In addition, the ability of Atf6 β to compensate for Atf6 α loss may vary by cell type. Secretory cells such as pancreatic beta cells may be especially dependent on Atf6 α to cope with the high basal ER function requirement inherent in synthesizing proteins for systemic use. The duration and intensity of Atf6 α activation may clearly play a role in the switch between pro-survival and pro-apoptotic UPR signaling, and some differences in results may be due to excess overexpression. Specific underlying mechanisms explaining divergent effects of Atf6 α require further study.

6. ATF6 α ALSO MODULATES CELL PROLIFERATION

In addition to the canonical Atf6 α downstream response leading to expanded ER proteosynthetic capacity, and the extensive evidence that Atf6 α modulates cell survival, some studies have found that Atf6 α action also influences the decision to enter the cell cycle. Some of the work implicating Atf6 α in proliferation has been performed in cancer or cancer cell lines [117–122]; other observations are in primary tissues such as pancreatic beta cells, ovarian cells, chondrocytes and cardiomyocytes [46,92,123–125]. In most cases, Atf6 α activation increased cell cycle entry, whether as part of healthy tissue growth and adaptation or as a maladaptive response to disease. Some data show a role for Atf6 α in tissue hypertrophy without hyperplasia [125–127]. An anti-proliferative effect has been described as well [128]. Since proliferation of normal tissues may be part of healthy adaptation or maladaptation, and proliferation of cancer cells leads to disease, Atf6 α roles in proliferation defy categorization as uniformly beneficial or harmful. What little is known of the mechanisms by which Atf6 α activation influences cell proliferation is summarized below.

6.1. Decreasing Atf6 α signaling inhibits cancer cell proliferation

Endoplasmic reticulum stress, and the UPR, are well-established features of cancer biology related to insults such as hypoxia, poor nutrient availability relative to metabolic demand, and oxidative stress [129]. Although the majority of studies investigating ATF6 α in cancer cells have focused on oncogenic cell survival in the face of ER stress, there is evidence that ATF6 α promotes proliferation as well. ATF6 α , known to be a tumorigenic factor in hepatocellular carcinoma [130], drives proliferation and BrdU incorporation in hepatoma cells in a proteostasis-stress dependent manner [118]. Treating cells with a proteasome inhibitor oprozomib decreased proliferation by inhibiting regulated intramembrane proteolysis, thus reducing ATF6 α cleavage and decreasing ATF6 α target gene activation [118]. Similarly, the antiretroviral protease inhibitor nelfinavir was found to have anti-proliferative effects in prostate cancer and liposarcoma through an off-target inhibition of S1P/S2P activation of ATF6 α [121,122]. In addition, knockdown of ATF6 α in glioblastoma cells reduced

proliferation, although the observation was weakened by use of a nonspecific viability assay to quantify proliferation [120]. These studies suggest that decreasing ATF6 α expression or activation reduced proliferation, supporting a pro-proliferative role for ATF6 α in human cancers, although downstream mechanisms were not tested.

6.2. Increasing Atf6 α signaling drives cancer cell proliferation

ATF6 α is implicated in the pathogenesis of hepatocellular carcinoma, and overexpression of nATF6 α in an HCC cell line increased gene expression of cell cycle associated genes [131]. In two colorectal cancer tumor banks, increased ATF6 α expression was associated with reduced disease-free survival [117]. To explore the mechanism, a mouse model was generated with tissue specific overexpression of active nuclear nAtf6 α in intestinal epithelial cells. These mice developed spontaneous colon cancer by 12 weeks of age, with increased proliferation of epithelial cells [117]. The mechanism was determined to be through an increase in gut permeability leading to inflammatory bacterial penetration into the gut wall, which activated Stat3 and led to cancer [117]. In endothelial cells, ATF6 α was found to activate a novel target, α B-crystallin, which was pro-proliferative via a mechanism involving VEGF [119].

6.3. Atf6 α influences cell cycle entry in non-transformed cells

Emerging evidence suggests that UPR pathways in general, and Atf6 α specifically, have previously unrecognized roles in normal organ development and function [1]. Deletion of both Atf6 α and Atf6 β is embryonic lethal [42]. To date, most evidence implicating Atf6 α in proliferation in normal tissues comes from neuro-endocrine type cells such as neurons, ovarian granulosa cells and pancreatic beta cells, or mesenchymal cells such as cartilage, vascular smooth muscle and cardiomyocytes. The involvement of Atf6 α in expansion of tissue cell number in response to tissue load stress is an interesting paradigm linking organ structure with function [123]. In each case, tissue growth may be beneficial or maladaptive.

6.4. Atf6 α promotes proliferation in endocrine cells and neurons

Unresolved ER stress contributes to diabetic decompensation, both through tissue insulin resistance in type 2 diabetes [132] and impaired beta cell insulin secretory capacity in type 1 and type 2 diabetes [2]. ER stress negatively impacts insulin production capacity through both impaired insulin synthesis and increased beta cell death [2]. However, modest physiological levels of ER stress may play a role in the adaptive increase in beta cell number that occurs in response to insulin demand [123,133]. During increased insulin demand in various physiological conditions (misfolding of pro-insulin, high fat feeding, hyperglycemia) beta cells showing activation of the UPR were more likely to enter the cell cycle [123]. Stress-associated beta cell proliferation was lost if ER stress was reduced using molecular chaperones, or by chemical inactivation or gene knockdown of Atf6 α [123]. Conversely, overexpression of Atf6 α was sufficient to drive proliferation in glucose-permissive conditions [123]. The mechanism by which Atf6 α promotes proliferation in beta cells is not yet known, but a proliferative role for Atf6 α has been observed in other endocrine cells as well. In ovarian granulosa cells, knockdown of Atf6 α arrested cells in S-phase, and caused a reduction in mRNA of cell cycle drivers cyclin A1, cyclin B1 and cyclin D2 [92]. Atf6 α may also play an intriguing toxic-proliferation role in the Huntington's neurodegenerative disorder. Neurons are post-mitotic and cannot tolerate cell cycle entry. In both mouse models and human disease, ATF6 α processing was found to be altered, leading to accumulation of full length Atf6 α , loss of the small GTPase Rheb, and inappropriate

accumulation of cell cycle drivers which resulted in neuronal cell death [124].

6.5. Atf6 α promotes proliferation in mesenchymal cells such as cartilage and smooth muscle

Atf6 α and Atf6 β are expressed throughout the proliferating and hypertrophic zones of cartilage development [125]. Metaphyseal chondrodysplasia type Schmid (MCDS) is an ER stress-associated dwarfism syndrome caused by mutations in type X collagen [46]. Overexpression of mutant collagen increased both Atf6 α and Atf6 β activation in HeLa cells. Ablation of Atf6 α in vitro diminished cellular transcriptional response to stress, and ablation of Atf6 α in vivo in MCDS mice worsened the disease phenotype, with expansion of the growth plate hypertrophic zone, decreased bone growth and increased Irf1 and Perk signaling [46]. On the other hand, ablation of Atf6 β in MCDS mice decreased ER stress markers and decreased the proliferation rate of growth plate chondrocytes [46]. Runx2, a transcription factor important for cartilage development, transcriptionally activates the Atf6 α gene [127]. Atf6 α was found to physically interact with Runx2, and Atf6 α overexpression promoted bone length increase in a long term hMSC culture model [125,127]. On the other hand, in two cell transformed chondrocyte cell lines, overexpression of Atf6 α decreased cell cycle entry as determined by a flow cytometry assay [128]. In human mesenchymal stem cells, CRISPR-mediated deletion of ATF6 α decreased population doubling, Ki67 levels, and percent of cells in S-phase, while increasing senescence markers [61]. RNAseq demonstrated loss of a number of cell-cycle promoting genes, including the protooncogene FOS [61]. Surprisingly, knockdown of FOS recapitulated many of the effects of ATF6 α deletion in hMSCs [61]. Maladaptive smooth muscle proliferation in pulmonary arterial hypertension was found to be mediated by Atf6 α [134]. Mild hypoxic stress activated Atf6 α ; interventions that decreased Atf6 α cleavage and target activation suppressed proliferation in smooth muscle cells both in vitro and in vivo [134]. In addition, ER stressors that increased nuclear Atf6 α in pulmonary artery smooth muscle cells increased proliferation as measured by nucleoside analog or Ki67, an effect that might be due to increased accumulation of intracellular iron [135].

6.6. Atf6 α can also drive tissue growth through cellular hypertrophy

Beyond increase in cell number, Atf6 α may also increase tissue mass by inducing cellular hypertrophy in chondrocytes and cardiomyocytes. In mice, Atf6 α deletion in the chondrodysplasia model described above resulted in reduced bone growth related, in part, to loss of cell height in the hypertrophic chondrocytes [46]. Atf6 α signaling was found to be markedly induced in two different cardiac hypertrophy models: pressure overload, by transverse aortic constriction, and hypertrophy in response to free-wheel exercise [126]. Conditional deletion of Atf6 α in cardiac myocytes decreased the hypertrophic response and caused cardiac dysfunction in both models; the mechanism was found to be through Atf6 α induction of mTor activator Rheb [126].

7. SUMMARY AND CONCLUSIONS

Atf6 α is an important stress response protein, sensing luminal ER stress and transmitting a nuclear signal that has many cellular effects. Canonically, in concert with other UPR pathways, Atf6 α activates gene expression of numerous ER resident proteins, resulting in enhanced protein folding capacity, resilience to oxidative/reductive stress, and degradation of ER luminal misfolded proteins. Activation of Atf6 α can have both pro-survival and pro-death outcomes, depending on the cell

Review

type and other contextual cues that largely remain to be clarified. ATF6 α is also implicated in driving tissue growth, through both proliferative and hypertrophic responses, which can be beneficial or maladaptive depending on the context. The literature contains numerous conflicting conclusions and unanswered questions that require further experimental investigation.

FUNDING

This article is part of a supplement entitled 'Biomarkers of Beta-Cell Health and Dysfunction: Towards Personalised Diabetes Care. Proceedings of the 20th Servier-IGIS Symposium'. The Symposium and the publication of the Supplement have been made possible by an unrestricted educational grant from Institut la Conférence Hippocrate — Servier Group.

ACKNOWLEDGMENTS

LCA, JTS and RBS wrote the manuscript, edited and approved the manuscript. The authors have no conflicts of interest with the work contained in this manuscript. This work was supported by NIH/NIDDK: R01DK114686 (LCA), R01 DK113300 (LCA), NIH/NIGMS: R25GM113686 (JTS), the American Diabetes Association grant #1-18-IBS-233 (LCA) in collaboration with the Order of the Amaranth, and the George F. and Sybil H. Fuller Foundation. The funding sources had no involvement in the analysis of the literature, the writing of this review, or in the decision to submit the article for publication.

CONFLICT OF INTEREST

None declared.

REFERENCES

- [1] Hillary, R.F., FitzGerald, U., 2018. A lifetime of stress: ATF6 in development and homeostasis. *Journal of Biomedical Science* 25(1):48–57.
- [2] Back, S.H., Kaufman, R.J., 2012. Endoplasmic reticulum stress and type 2 diabetes. *Annual Review of Biochemistry* 81:767–793.
- [3] Ron, D., Harding, H.P., 2012. Protein-folding homeostasis in the endoplasmic reticulum and nutritional regulation. *Cold Spring Harbor Perspectives in Biology* 4(12):1–13.
- [4] Hetz, C., Chevet, E., Oakes, S.A., 2015. Proteostasis control by the unfolded protein response. *Nature Cell Biology* 17(7):829–838.
- [5] Hetz, C., Papa, F.R., 2018. The unfolded protein response and cell fate control. *Molecular Cell* 69(2):169–181.
- [6] Hetz, C., 2012. The unfolded protein response: controlling cell fate decisions under ER stress and beyond. *Nature Reviews Molecular Cell Biology* 13(2): 89–102.
- [7] Galluzzi, L., Vitale, I., Aaronson, S.A., Abrams, J.M., Adam, D., Agostinis, P., et al., 2018. Molecular mechanisms of cell death: recommendations of the nomenclature committee on cell death 2018. *Cell Death and Differentiation* 25(3):486–541.
- [8] Yoshida, H., Haze, K., Yanagi, H., Yura, T., Mori, K., 1998. Identification of the cis-acting endoplasmic reticulum stress response element responsible for transcriptional induction of mammalian glucose-regulated proteins. Involvement of basic leucine zipper transcription factors. *The Journal of Biological Chemistry* 273(50):33741–33749.
- [9] Hai, T.W., Liu, F., Coukos, W.J., Green, M.R., 1989. Transcription factor ATF cDNA clones: an extensive family of leucine zipper proteins able to selectively form DNA-binding heterodimers. *Genes and Development* 3(12B):2083–2090.

- [10] Zhu, C., Johansen, F.E., Prywes, R., 1997. Interaction of ATF6 and serum response factor. *Molecular and Cellular Biology* 17(9):4957–4966.
- [11] Kondo, S., Saito, A., Asada, R., Kanemoto, S., Imaizumi, K., 2011. Physiological unfolded protein response regulated by OASIS family members, transmembrane bZIP transcription factors. *IUBMB Life* 63(4):233–239.
- [12] Hollien, J., 2013. Evolution of the unfolded protein response. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1833(11):2458–2463.
- [13] Haze, K., Yoshida, H., Yanagi, H., Yura, T., Mori, K., 1999. Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Molecular Biology of the Cell* 10(11):3787–3799.
- [14] Li, M., Baumeister, P., Roy, B., Phan, T., Foti, D., Luo, S., et al., 2000. ATF6 as a transcription activator of the endoplasmic reticulum stress element: thapsigargin stress-induced changes and synergistic interactions with NF- κ B and YY1. *Molecular and Cellular Biology* 20(14):5096–5106.
- [15] Ye, J., Rawson, R.B., Komuro, R., Chen, X., Davé, U.P., Prywes, R., et al., 2000. ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Molecular Cell* 6(6):1355–1364.
- [16] Thuerauf, D.J., Morrison, L.E., Hoover, H., Glembotski, C.C., 2002. Coordination of ATF6-mediated transcription and ATF6 degradation by a domain that is shared with the viral transcription factor, VP16. *The Journal of Biological Chemistry* 277(23):20734–20739.
- [17] Shen, J., Chen, X., Hendershot, L., Prywes, R., 2002. ER stress regulation of ATF6 localization by dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Developmental Cell* 3(1):99–111.
- [18] Chen, X., Shen, J., Prywes, R., 2002. The luminal domain of ATF6 senses endoplasmic reticulum (ER) stress and causes translocation of ATF6 from the ER to the Golgi. *The Journal of Biological Chemistry* 277(15):13045–13052.
- [19] Schindler, A.J., Schekman, R., 2009. In vitro reconstitution of ER-stress induced ATF6 transport in COPII vesicles. *Proceedings of the National Academy of Sciences of the United States of America* 106(42):17775–17780.
- [20] Nadanaka, S., Yoshida, H., Kano, F., Murata, M., Mori, K., 2004. Activation of mammalian unfolded protein response is compatible with the quality control system operating in the endoplasmic reticulum. *Molecular Biology of the Cell* 15(6):2537–2548.
- [21] Fang, J., Liu, M., Zhang, X., Sakamoto, T., Taatjes, D.J., Jena, B.P., et al., 2015. COPII-dependent ER export: a critical component of insulin biogenesis and β -cell ER homeostasis. *Molecular Endocrinology (Baltimore, Md.)* 29(8): 1156–1169.
- [22] Okada, T., Haze, K., Nadanaka, S., Yoshida, H., Seidah, N.G., Hirano, Y., et al., 2003. A serine protease inhibitor prevents endoplasmic reticulum stress-induced cleavage but not transport of the membrane-bound transcription factor ATF6. *The Journal of Biological Chemistry* 278(33):31024–31032.
- [23] Tsukumo, Y., Tomida, A., Kitahara, O., Nakamura, Y., Asada, S., Mori, K., et al., 2007. Nucleobindin 1 controls the unfolded protein response by inhibiting ATF6 activation. *The Journal of Biological Chemistry* 282(40): 29264–29272.
- [24] Naranjo, J.R., Zhang, H., Villar, D., González, P., Dopazo, X.M., Morón-Oset, J., et al., 2016. Activating transcription factor 6 derepression mediates neuroprotection in Huntington disease. *The Journal of Clinical Investigation* 126(2):627–638.
- [25] Nadanaka, S., Okada, T., Yoshida, H., Mori, K., 2007. Role of disulfide bridges formed in the luminal domain of ATF6 in sensing endoplasmic reticulum stress. *Molecular and Cellular Biology* 27(3):1027–1043.
- [26] Higa, A., Taouji, S., Lhomond, S., Jensen, D., Fernandez-Zapico, M.E., Simpson, J.C., et al., 2014. Endoplasmic reticulum stress-activated transcription factor ATF6 α requires the disulfide isomerase PDIA5 to modulate chemoresistance. *Molecular and Cellular Biology* 34(10):1839–1849.
- [27] Hong, M., Luo, S., Baumeister, P., Huang, J.-M., Gogia, R.K., Li, M., et al., 2004. Underglycosylation of ATF6 as a novel sensing mechanism for

- activation of the unfolded protein response. *The Journal of Biological Chemistry* 279(12):11354–11363.
- [28] Gade, P., Manjogowda, S.B., Nallar, S.C., Maachani, U.B., Cross, A.S., Kalvakolanu, D.V., 2014. Regulation of the death-associated protein kinase 1 expression and autophagy via ATF6 requires apoptosis signal-regulating kinase 1. *Molecular and Cellular Biology* 34(21):4033–4048.
- [29] Luo, S., Lee, A.S., 2002. Requirement of the p38 mitogen-activated protein kinase signalling pathway for the induction of the 78 kDa glucose-regulated protein/immunoglobulin heavy-chain binding protein by azetidine stress: activating transcription factor 6 as a target for stress-induced phosphorylation. *The Biochemical Journal* 366(Pt 3):787–795.
- [30] Thuerauf, D.J., Arnold, N.D., Zechner, D., Hanford, D.S., DeMartin, K.M., McDonough, P.M., et al., 1998. p38 Mitogen-activated protein kinase mediates the transcriptional induction of the atrial natriuretic factor gene through a serum response element. A potential role for the transcription factor ATF6. *The Journal of Biological Chemistry* 273(32):20636–20643.
- [31] Lee, A.-H., Iwakoshi, N.N., Glimcher, L.H., 2003. XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Molecular and Cellular Biology* 23(21):7448–7459.
- [32] Namba, T., Ishihara, T., Tanaka, K., Hoshino, T., Mizushima, T., 2007. Transcriptional activation of ATF6 by endoplasmic reticulum stressors. *Biochemical and Biophysical Research Communications* 355(2):543–548.
- [33] Baird, L., Tsujita, T., Kobayashi, E.H., Funayama, R., Nagashima, T., Nakayama, K., et al., 2017. A homeostatic shift facilitates endoplasmic reticulum proteostasis through transcriptional integration of proteostatic stress response pathways. *Molecular and Cellular Biology* 37(4) e00439-16.
- [34] Misra, J., Kim, D.-K., Choi, W., Koo, S.-H., Lee, C.-H., Back, S.-H., et al., 2013. Transcriptional cross talk between orphan nuclear receptor ERR γ and transmembrane transcription factor ATF6 α coordinates endoplasmic reticulum stress response. *Nucleic Acids Research* 41(14):6960–6974.
- [35] Li, W.W., Hsiung, Y., Zhou, Y., Roy, B., Lee, A.S., 1997. Induction of the mammalian GRP78/BIP gene by Ca²⁺ depletion and formation of aberrant proteins: activation of the conserved stress-inducible grp core promoter element by the human nuclear factor YY1. *Molecular and Cellular Biology* 17(1):54–60.
- [36] Wang, Y., Shen, J., Arenzana, N., Tirasophon, W., Kaufman, R.J., Prywes, R., 2000. Activation of ATF6 and an ATF6 DNA binding site by the endoplasmic reticulum stress response. *The Journal of Biological Chemistry* 275(35):27013–27020.
- [37] Yamamoto, K., Yoshida, H., Kokame, K., Kaufman, R.J., Mori, K., 2004. Differential contributions of ATF6 and XBP1 to the activation of endoplasmic reticulum stress-responsive cis-acting elements ERSE, UPRE and ERSE-II. *Journal of Biochemistry* 136(3):343–350.
- [38] Yoshida, H., Okada, T., Haze, K., Yanagi, H., Yura, T., Negishi, M., et al., 2001. Endoplasmic reticulum stress-induced formation of transcription factor complex ERSF including NF-Y (CBF) and activating transcription factors 6 α and 6 β that activates the mammalian unfolded protein response. *Molecular and Cellular Biology* 21(4):1239–1248.
- [39] Shoulders, M.D., Ryno, L.M., Genereux, J.C., Moresco, J.J., Tu, P.G., Wu, C., et al., 2013. Stress-independent activation of XBP1s and/or ATF6 reveals three functionally diverse ER proteostasis environments. *Cell Reports* 3(4):1279–1292.
- [40] Yoshida, H., Matsui, T., Yamamoto, A., Okada, T., Mori, K., 2001. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 107(7):881–891.
- [41] Walter, F., O'Brien, A., Concannon, C.G., Düssmann, H., Prehn, J.H.M., 2018. ER stress signaling has an activating transcription factor 6 α (ATF6)-dependent “off-switch”. *The Journal of Biological Chemistry* 293(47):18270–18284.
- [42] Yamamoto, K., Sato, T., Matsui, T., Sato, M., Okada, T., Yoshida, H., et al., 2007. Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6 α and XBP1. *Developmental Cell* 13(3):365–376.
- [43] Haze, K., Okada, T., Yoshida, H., Yanagi, H., Yura, T., Negishi, M., et al., 2001. Identification of the G13 (cAMP-response-element-binding protein-related protein) gene product related to activating transcription factor 6 as a transcriptional activator of the mammalian unfolded protein response. *The Biochemical Journal* 355(Pt 1):19–28.
- [44] Wu, J., Rutkowski, D.T., Dubois, M., Swathirajan, J., Saunders, T., Wang, J., et al., 2007. ATF6 α optimizes long-term endoplasmic reticulum function to protect cells from chronic stress. *Developmental Cell* 13(3):351–364.
- [45] Thuerauf, D.J., Morrison, L., Glembotski, C.C., 2004. Opposing roles for ATF6 α and ATF6 β in endoplasmic reticulum stress response gene induction. *The Journal of Biological Chemistry* 279(20):21078–21084.
- [46] Forouhan, M., Mori, K., Boot-Handford, R.P., 2018. Paradoxical roles of ATF6 α and ATF6 β in modulating disease severity caused by mutations in collagen X. *Matrix Biology Journal of the International Society for Matrix Biology* 70:50–71.
- [47] Yoshida, H., Okada, T., Haze, K., Yanagi, H., Yura, T., Negishi, M., et al., 2000. ATF6 activated by proteolysis binds in the presence of NF-Y (CBF) directly to the cis-acting element responsible for the mammalian unfolded protein response. *Molecular and Cellular Biology* 20(18):6755–6767.
- [48] Okada, T., Yoshida, H., Akazawa, R., Negishi, M., Mori, K., 2002. Distinct roles of activating transcription factor 6 (ATF6) and double-stranded RNA-activated protein kinase-like endoplasmic reticulum kinase (PERK) in transcription during the mammalian unfolded protein response. *The Biochemical Journal* 366(Pt 2):585–594.
- [49] Matsuda, T., Kido, Y., Asahara, S., Kaisho, T., Tanaka, T., Hashimoto, N., et al., 2010. Ablation of C/EBP β alleviates ER stress and pancreatic β cell failure through the GRP78 chaperone in mice. *The Journal of Clinical Investigation* 120(1):115–126.
- [50] Horimoto, S., Ninagawa, S., Okada, T., Koba, H., Sugimoto, T., Kamiya, Y., et al., 2013. The unfolded protein response transducer ATF6 represents a novel transmembrane-type endoplasmic reticulum-associated degradation substrate requiring both mannose trimming and SEL1L protein. *The Journal of Biological Chemistry* 288(44):31517–31527.
- [51] Hong, M., Li, M., Mao, C., Lee, A.S., 2004. Endoplasmic reticulum stress triggers an acute proteasome-dependent degradation of ATF6. *Journal of Cellular Biochemistry* 92(4):723–732.
- [52] Fonseca, S.G., Ishigaki, S., Oslowski, C.M., Lu, S., Lipson, K.L., Ghosh, R., et al., 2010. Wolfram syndrome 1 gene negatively regulates ER stress signaling in rodent and human cells. *The Journal of Clinical Investigation* 120(3):744–755.
- [53] Yoshida, H., Uemura, A., Mori, K., 2009. pXBP1(U), a negative regulator of the unfolded protein response activator pXBP1(S), targets ATF6 but not ATF4 in proteasome-mediated degradation. *Cell Structure and Function* 34(1):1–10.
- [54] Chiang, W.-C., Chan, P., Wissinger, B., Vincent, A., Skorczyk-Werner, A., Krawczyński, M.R., et al., 2017. Achromatopsia mutations target sequential steps of ATF6 activation. *Proceedings of the National Academy of Sciences of the United States of America* 114(2):400–405.
- [55] Cnop, M., Abdulkarim, B., Bottu, G., Cunha, D.A., Igoillo-Esteve, M., Masini, M., et al., 2014. RNA sequencing identifies dysregulation of the human pancreatic islet transcriptome by the saturated fatty acid palmitate. *Diabetes* 63(6):1978–1993.
- [56] Engin, F., Yermalovich, A., Nguyen, T., Nguyen, T., Hummasti, S., Fu, W., et al., 2013. Restoration of the unfolded protein response in pancreatic β cells protects mice against type 1 diabetes. *Science Translational Medicine* 5(211):211ra156.
- [57] Adachi, Y., Yamamoto, K., Okada, T., Yoshida, H., Harada, A., Mori, K., 2008. ATF6 is a transcription factor specializing in the regulation of quality control proteins in the endoplasmic reticulum. *Cell Structure and Function* 33(1):75–89.

- [58] Belmont, P.J., Tadimalla, A., Chen, W.J., Martindale, J.J., Thuerauf, D.J., Marcinko, M., et al., 2008. Coordination of growth and endoplasmic reticulum stress signaling by regulator of calcineurin 1 (RCAN1), a novel ATF6-inducible gene. *The Journal of Biological Chemistry* 283(20):14012–14021.
- [59] Maiuolo, J., Bulotta, S., Verderio, C., Benfante, R., Borgese, N., 2011. Selective activation of the transcription factor ATF6 mediates endoplasmic reticulum proliferation triggered by a membrane protein. *Proceedings of the National Academy of Sciences of the United States of America* 108(19):7832–7837.
- [60] Luo, S., Mao, C., Lee, B., Lee, A.S., 2006. GRP78/BiP is required for cell proliferation and protecting the inner cell mass from apoptosis during early mouse embryonic development. *Molecular and Cellular Biology* 26(15):5688–5697.
- [61] Wang, S., Hu, B., Ding, Z., Dang, Y., Wu, J., Li, D., et al., 2018. ATF6 safeguards organelle homeostasis and cellular aging in human mesenchymal stem cells. *Cell Discovery* 4:2.
- [62] Pan, Z., Erkan, M., Streit, S., Friess, H., Kleeff, J., 2009. Silencing of GRP94 expression promotes apoptosis in pancreatic cancer cells. *International Journal of Oncology* 35(4):823–828.
- [63] Fu, Y., Li, J., Lee, A.S., 2007. GRP78/BiP inhibits endoplasmic reticulum BIK and protects human breast cancer cells against estrogen starvation-induced apoptosis. *Cancer Research* 67(8):3734–3740.
- [64] Zhou, H., Zhang, Y., Fu, Y., Chan, L., Lee, A.S., 2011. Novel mechanism of anti-apoptotic function of 78-kDa glucose-regulated protein (GRP78): endocrine resistance factor in breast cancer, through release of B-cell lymphoma 2 (BCL-2) from BCL-2-interacting killer (BIK). *The Journal of Biological Chemistry* 286(29):25687–25696.
- [65] Synofzik, M., Haack, T.B., Kopajtic, R., Gorza, M., Rapaport, D., Greiner, M., et al., 2014. Absence of BiP co-chaperone DNAJC3 causes diabetes mellitus and multisystemic neurodegeneration. *American Journal of Human Genetics* 95(6):689–697.
- [66] Genereux, J.C., Qu, S., Zhou, M., Ryno, L.M., Wang, S., Shoulders, M.D., et al., 2015. Unfolded protein response-induced ERdj3 secretion links ER stress to extracellular proteostasis. *The European Molecular Biology Organization Journal* 34(1):4–19.
- [67] Ellgaard, L., Sevier, C.S., Billeid, N.J., 2018. How are proteins reduced in the endoplasmic reticulum? *Trends in Biochemical Sciences* 43(1):32–43.
- [68] Koritzinsky, M., Levitin, F., van den Beucken, T., Rumantir, R.A., Harding, N.J., Chu, K.C., et al., 2013. Two phases of disulfide bond formation have differing requirements for oxygen. *The Journal of Cell Biology* 203(4):615–627.
- [69] Martindale, J.J., Fernandez, R., Thuerauf, D., Whittaker, R., Gude, N., Sussman, M.A., et al., 2006. Endoplasmic reticulum stress gene induction and protection from ischemia/reperfusion injury in the hearts of transgenic mice with a tamoxifen-regulated form of ATF6. *Circulation Research* 98(9):1186–1193.
- [70] Doroudgar, S., Thuerauf, D.J., Marcinko, M.C., Belmont, P.J., Glembotski, C.C., 2009. Ischemia activates the ATF6 branch of the endoplasmic reticulum stress response. *The Journal of Biological Chemistry* 284(43):29735–29745.
- [71] Tufo, G., Jones, A.W.E., Wang, Z., Hamelin, J., Tajeddine, N., Esposti, D.D., et al., 2014. The protein disulfide isomerases PDIA4 and PDIA6 mediate resistance to cisplatin-induced cell death in lung adenocarcinoma. *Cell Death and Differentiation* 21(5):685–695.
- [72] Vekich, J.A., Belmont, P.J., Thuerauf, D.J., Glembotski, C.C., 2012. Protein disulfide isomerase-associated 6 is an ATF6-inducible ER stress response protein that protects cardiac myocytes from ischemia/reperfusion-mediated cell death. *Journal of Molecular and Cellular Cardiology* 53(2):259–267.
- [73] Jin, J.-K., Blackwood, E.A., Azizi, K., Thuerauf, D.J., Fahem, A.G., Hofmann, C., et al., 2017. ATF6 decreases myocardial ischemia/reperfusion damage and links ER stress and oxidative stress signaling pathways in the heart. *Circulation Research* 120(5):862–875.
- [74] Yu, Z., Sheng, H., Liu, S., Zhao, S., Glembotski, C.C., Warner, D.S., et al., 2017. Activation of the ATF6 branch of the unfolded protein response in neurons improves stroke outcome. *Journal of Cerebral Blood Flow and Metabolism — Official Journal of the International Society of Cerebral Blood Flow and Metabolism* 37(3):1069–1079.
- [75] Halterman, M.W., Gill, M., DeJesus, C., Ogihara, M., Schor, N.F., Federoff, H.J., 2010. The endoplasmic reticulum stress response factor CHOP-10 protects against hypoxia-induced neuronal death. *The Journal of Biological Chemistry* 285(28):21329–21340.
- [76] Belmont, P.J., Chen, W.J., San Pedro, M.N., Thuerauf, D.J., Gellings Lowe, N., Gude, N., et al., 2010. Roles for endoplasmic reticulum-associated degradation and the novel endoplasmic reticulum stress response gene Derlin-3 in the ischemic heart. *Circulation Research* 106(2):307–316.
- [77] Oda, Y., Okada, T., Yoshida, H., Kaufman, R.J., Nagata, K., Mori, K., 2006. Derlin-2 and Derlin-3 are regulated by the mammalian unfolded protein response and are required for ER-associated degradation. *The Journal of Cell Biology* 172(3):383–393.
- [78] Dang, J., Bian, X., Ma, X., Li, J., Long, F., Shan, S., et al., 2017. ORMDL3 facilitates the survival of splenic B cells via an ATF6 α -endoplasmic reticulum stress-beclin1 autophagy regulatory pathway. *Journal of Immunology (Baltimore, Md.: 1950)* 199(5):1647–1659.
- [79] Gade, P., Ramachandran, G., Maachani, U.B., Rizzo, M.A., Okada, T., Prywes, R., et al., 2012. An IFN- γ -stimulated ATF6-C/EBP- β -signaling pathway critical for the expression of Death Associated Protein Kinase 1 and induction of autophagy. *Proceedings of the National Academy of Sciences of the United States of America* 109(26):10316–10321.
- [80] Sharma, M., Bhattacharyya, S., Sharma, K.B., Chauhan, S., Asthana, S., Abidin, M.Z., et al., 2017. Japanese encephalitis virus activates autophagy through XBP1 and ATF6 ER stress sensors in neuronal cells. *The Journal of General Virology* 98(5):1027–1039.
- [81] Ogata, M., Hino, S., Saito, A., Morikawa, K., Kondo, S., Kanemoto, S., et al., 2006. Autophagy is activated for cell survival after endoplasmic reticulum stress. *Molecular and Cellular Biology* 26(24):9220–9231.
- [82] Lee, Y.-R., Kuo, S.-H., Lin, C.-Y., Fu, P.-J., Lin, Y.-S., Yeh, T.-M., et al., 2018. Dengue virus-induced ER stress is required for autophagy activation, viral replication, and pathogenesis both in vitro and in vivo. *Scientific Reports* 8(1):489.
- [83] Allen, D., Seo, J., 2018. ER stress activates the TOR pathway through Atf6. *Journal of Molecular Signaling* 13:1–9.
- [84] Schewe, D.M., Aguirre-Ghiso, J.A., 2008. ATF6 α -Rheb-mTOR signaling promotes survival of dormant tumor cells in vivo. *Proceedings of the National Academy of Sciences of the United States of America* 105(30):10519–10524.
- [85] Karali, E., Bellou, S., Stellas, D., Klinakis, A., Murphy, C., Fotsis, T., 2014. VEGF Signals through ATF6 and PERK to promote endothelial cell survival and angiogenesis in the absence of ER stress. *Molecular Cell* 54(4):559–572.
- [86] Teodoro, T., Odisho, T., Sidorova, E., Volchuk, A., 2012. Pancreatic β -cells depend on basal expression of active ATF6 α -p50 for cell survival even under nonstress conditions. *American Journal of Physiology Cell Physiology* 302(7):C992–C1003.
- [87] Usui, M., Yamaguchi, S., Tanji, Y., Tominaga, R., Ishigaki, Y., Fukumoto, M., et al., 2012. Atf6 α -null mice are glucose intolerant due to pancreatic β -cell failure on a high-fat diet but partially resistant to diet-induced insulin resistance. *Metabolism* 61(8):1118–1128.
- [88] Rao, J., Yue, S., Fu, Y., Zhu, J., Wang, X., Busuttil, R.W., et al., 2014. ATF6 mediates a pro-inflammatory synergy between ER stress and TLR activation in the pathogenesis of liver ischemia-reperfusion injury. *American Journal of Transplantation — Official Journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 14(7):1552–1561.

- [89] Zhang, Z., Wu, S., Muhammad, S., Ren, Q., Sun, C., 2018. miR-103/107 promote ER stress-mediated apoptosis via targeting the Wnt3a/ β -catenin/ATF6 pathway in preadipocytes. *Journal of Lipid Research* 59(5):843–853.
- [90] Huang, J., Wan, L., Lu, H., Li, X., 2018. High expression of active ATF6 aggravates endoplasmic reticulum stress-induced vascular endothelial cell apoptosis through the mitochondrial apoptotic pathway. *Molecular Medicine Reports* 17(5):6483–6489.
- [91] Pagliarini, V., Giglio, P., Bernardoni, P., De Zio, D., Fimia, G.M., Piacentini, M., et al., 2015. Downregulation of E2F1 during ER stress is required to induce apoptosis. *Journal of Cell Science* 128(6):1166–1179.
- [92] Xiong, Y., Chen, H., Lin, P., Wang, A., Wang, L., Jin, Y., 2017. ATF6 knockdown decreases apoptosis, arrests the S phase of the cell cycle, and increases steroid hormone production in mouse granulosa cells. *American Journal of Physiology Cell Physiology* 312(3):C341–C353.
- [93] Wang, N., Zhao, F., Lin, P., Zhang, G., Tang, K., Wang, A., et al., 2017. Knockdown of XBP1 by RNAi in mouse granulosa cells promotes apoptosis, inhibits cell cycle, and decreases estradiol synthesis. *International Journal of Molecular Sciences* 18(6):1–13.
- [94] Morishima, N., Nakanishi, K., Nakano, A., 2011. Activating transcription factor-6 (ATF6) mediates apoptosis with reduction of myeloid cell leukemia sequence 1 (Mcl-1) protein via induction of WW domain binding protein 1. *The Journal of Biological Chemistry* 286(40):35227–35235.
- [95] Ma, Y., Brewer, J.W., Diehl, J.A., Hendershot, L.M., 2002. Two distinct stress signaling pathways converge upon the CHOP promoter during the mammalian unfolded protein response. *Journal of Molecular Biology* 318(5):1351–1365.
- [96] Zinszner, H., Kuroda, M., Wang, X., Batchvarova, N., Lightfoot, R.T., Remotti, H., et al., 1998. CHOP is implicated in programmed cell death in response to impaired function of the endoplasmic reticulum. *Genes and Development* 12(7):982–995.
- [97] Oyadomari, S., Koizumi, A., Takeda, K., Gotoh, T., Akira, S., Araki, E., et al., 2002. Targeted disruption of the Chop gene delays endoplasmic reticulum stress-mediated diabetes. *The Journal of Clinical Investigation* 109(4):525–532.
- [98] Song, B., Scheuner, D., Ron, D., Pennathur, S., Kaufman, R.J., 2008. Chop deletion reduces oxidative stress, improves beta cell function, and promotes cell survival in multiple mouse models of diabetes. *The Journal of Clinical Investigation* 118(10):3378–3389.
- [99] Kaufman, R.J., Back, S.H., Song, B., Han, J., Hassler, J., 2010. The unfolded protein response is required to maintain the integrity of the endoplasmic reticulum, prevent oxidative stress and preserve differentiation in β -cells. *Diabetes Obesity and Metabolism* 12(Suppl. 2):99–107.
- [100] Gurlo, T., Rivera, J.F., Butler, A.E., Cory, M., Hoang, J., Costes, S., et al., 2016. CHOP contributes to, but is not the only mediator of, IAPP induced β -cell apoptosis. *Molecular Endocrinology* (Baltimore, Md.) 30(4):446–454.
- [101] Silva, R.M., Ries, V., Oo, T.F., Yarygina, O., Jackson-Lewis, V., Ryu, E.J., et al., 2005. CHOP/GADD153 is a mediator of apoptotic death in substantia nigra dopamine neurons in an in vivo neurotoxin model of parkinsonism. *Journal of Neurochemistry* 95(4):974–986.
- [102] Prasanthi, J.R.P., Larson, T., Schommer, J., Ghribi, O., 2011. Silencing GADD153/CHOP gene expression protects against Alzheimer's disease-like pathology induced by 27-hydroxycholesterol in rabbit hippocampus. *Public Library of Science one* 6(10):e26420.
- [103] Marciniak, S.J., Yun, C.Y., Oyadomari, S., Novoa, I., Zhang, Y., Jungreis, R., et al., 2004. CHOP induces death by promoting protein synthesis and oxidation in the stressed endoplasmic reticulum. *Genes and Development* 18(24):3066–3077.
- [104] Dong, B., Zhou, H., Han, C., Yao, J., Xu, L., Zhang, M., et al., 2014. Ischemia/reperfusion-induced CHOP expression promotes apoptosis and impairs renal function recovery: the role of acidosis and GPR4. *Public Library of Science one* 9(10):e110944.
- [105] Shao, D., Ni, J., Shen, Y., Liu, J., Zhou, L., Xue, H., et al., 2015. CHOP mediates XBP1S-induced renal mesangial cell necrosis following high glucose treatment. *European Journal of Pharmacology* 758:89–96.
- [106] Noh, M.R., Woo, C.-H., Park, M.-J., Kim, J., Park, K.M., 2018. Ablation of C/EBP homologous protein attenuates renal fibrosis after ureteral obstruction by reducing autophagy and microtubule disruption. *Biochimica et Biophysica Acta — Molecular Basis of Disease* 1864(5 Pt A):1634–1641.
- [107] Yang, J.-R., Yao, F.-H., Zhang, J.-G., Ji, Z.-Y., Li, K.-L., Zhan, J., et al., 2014. Ischemia-reperfusion induces renal tubule pyroptosis via the CHOP-caspase-11 pathway. *American Journal of Physiology Renal Physiology* 306(1):F75–F84.
- [108] Ferlito, M., Wang, Q., Fulton, W.B., Colombani, P.M., Marchionni, L., Fox-Talbot, K., et al., 2014. Hydrogen sulfide [corrected] increases survival during sepsis: protective effect of CHOP inhibition. *Journal of Immunology* (Baltimore, Md.: 1950) 192(4):1806–1814.
- [109] McCullough, K.D., Martindale, J.L., Klotz, L.O., Aw, T.Y., Holbrook, N.J., 2001. Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state. *Molecular and Cellular Biology* 21(4):1249–1259.
- [110] Ghosh, A.P., Klocke, B.J., Ballestas, M.E., Roth, K.A., 2012. CHOP potentially co-operates with FOXO3a in neuronal cells to regulate PUMA and BIM expression in response to ER stress. *Public Library of Science one* 7(6):e39586.
- [111] Han, J., Murthy, R., Wood, B., Song, B., Wang, S., Sun, B., et al., 2013. ER stress signalling through eIF2 α and CHOP, but not IRE1 α , attenuates adipogenesis in mice. *Diabetologia* 56(4):911–924.
- [112] Rutkowski, D.T., Arnold, S.M., Miller, C.N., Wu, J., Li, J., Gunnison, K.M., et al., 2006. Adaptation to ER stress is mediated by differential stabilities of pro-survival and pro-apoptotic mRNAs and proteins. *Public Library of Science Biology* 4(11):e374.
- [113] Lépine, S., Allegood, J.C., Edmonds, Y., Milstien, S., Spiegel, S., 2011. Autophagy induced by deficiency of sphingosine-1-phosphate phosphohydrolase 1 is switched to apoptosis by calpain-mediated autophagy-related gene 5 (Atg5) cleavage. *The Journal of Biological Chemistry* 286(52):44380–44390.
- [114] La, X., Zhang, L., Li, Z., Yang, P., Wang, Y., 2017. Berberine-induced autophagic cell death by elevating GRP78 levels in cancer cells. *Oncotarget* 8(13):20909–20924.
- [115] Guo, F.-J., Xiong, Z., Lu, X., Ye, M., Han, X., Jiang, R., 2014. ATF6 upregulates XBP1S and inhibits ER stress-mediated apoptosis in osteoarthritic cartilage. *Cellular Signalling* 26(2):332–342.
- [116] Nakajima, S., Hiramatsu, N., Hayakawa, K., Saito, Y., Kato, H., Huang, T., et al., 2011. Selective abrogation of BiP/GRP78 blunts activation of NF- κ B through the ATF6 branch of the UPR: involvement of C/EBP β and mTOR-dependent dephosphorylation of Akt. *Molecular and Cellular Biology* 31(8):1710–1718.
- [117] Coleman, O.I., Lobner, E.M., Bierwirth, S., Sorbie, A., Waldschmitt, N., Rath, E., et al., 2018. Activated ATF6 induces intestinal dysbiosis and innate immune response to promote colorectal tumorigenesis. *Gastroenterology* 155(5):1539–1552.e12.
- [118] Vandewynckel, Y.-P., Coucke, C., Laukens, D., Devisscher, L., Paridaens, A., Bogaerts, E., et al., 2016. Next-generation proteasome inhibitor oprozomib synergizes with modulators of the unfolded protein response to suppress hepatocellular carcinoma. *Oncotarget* 7(23):34988–35000.
- [119] Ruan, Q., Han, S., Jiang, W.G., Boulton, M.E., Chen, Z.J., Law, B.K., et al., 2011. α B-crystallin, an effector of unfolded protein response, confers anti-VEGF resistance to breast cancer via maintenance of intracrine VEGF in endothelial cells. *Molecular Cancer Research MCR* 9(12):1632–1643.
- [120] Dadey, D.Y.A., Kapoor, V., Khudanyan, A., Urano, F., Kim, A.H., Thotala, D., et al., 2016. The ATF6 pathway of the ER stress response contributes to enhanced viability in glioblastoma. *Oncotarget* 7(2):2080–2092.

Review

- [121] Guan, M., Fousek, K., Jiang, C., Guo, S., Synold, T., Xi, B., et al., 2011. Nelfinavir induces liposarcoma apoptosis through inhibition of regulated intramembrane proteolysis of SREBP-1 and ATF6. *Clinical Cancer Research — An Official Journal of the American Association for Cancer Research* 17(7):1796–1806.
- [122] Guan, M., Fousek, K., Chow, W.A., 2012. Nelfinavir inhibits regulated intramembrane proteolysis of sterol regulatory element binding protein-1 and activating transcription factor 6 in castration-resistant prostate cancer. *The FEBS Journal* 279(13):2399–2411.
- [123] Sharma, R.B., O'Donnell, A.C., Stamateris, R.E., Ha, B., McCloskey, K.M., Reynolds, P.R., et al., 2015. Insulin demand regulates β cell number via the unfolded protein response. *The Journal of Clinical Investigation* 125(10):3831–3846.
- [124] Fernandez-Fernandez, M.R., Ferrer, I., Lucas, J.J., 2011. Impaired ATF6 α processing, decreased Rheb and neuronal cell cycle re-entry in Huntington's disease. *Neurobiology of Disease* 41(1):23–32.
- [125] Xiong, Z., Jiang, R., Zhang, P., Han, X., Guo, F.-J., 2015. Transmission of ER stress response by ATF6 promotes endochondral bone growth. *Journal of Orthopaedic Surgery and Research* 10(141):1–12.
- [126] Blackwood, E.A., Hofmann, C., Santo Domingo, M., Bilal, A.S., Sarakki, A., Stauffer, W., et al., 2019. ATF6 regulates cardiac hypertrophy by transcriptional induction of the mTORC1 activator, Rheb. *Circulation Research* 124(1):79–93.
- [127] Guo, F., Han, X., Wu, Z., Cheng, Z., Hu, Q., Zhao, Y., et al., 2016. ATF6a, a Runx2-activable transcription factor, is a new regulator of chondrocyte hypertrophy. *Journal of Cell Science* 129(4):717–728.
- [128] Han, X., Zhang, P., Jiang, R., Xia, F., Li, M., Guo, F.-J., 2014. Explore on the effect of ATF6 on cell growth and apoptosis in cartilage development. *Histochemistry and Cell Biology* 142(5):497–509.
- [129] Cubillos-Ruiz, J.R., Bettigole, S.E., Glimcher, L.H., 2017. Tumorigenic and immunosuppressive effects of endoplasmic reticulum stress in cancer. *Cell* 168(4):692–706.
- [130] Wu, H., Wei, L., Fan, F., Ji, S., Zhang, S., Geng, J., et al., 2015. Integration of Hippo signalling and the unfolded protein response to restrain liver overgrowth and tumorigenesis. *Nature Communications* 6:6239–6254.
- [131] Arai, M., Kondoh, N., Imazeki, N., Hada, A., Hatsuse, K., Kimura, F., et al., 2006. Transformation-associated gene regulation by ATF6 α during hepatocarcinogenesis. *FEBS Letters* 580(1):184–190.
- [132] Ozcan, U., Cao, Q., Yilmaz, E., Lee, A.-H., Iwakoshi, N.N., Ozdelen, E., et al., 2004. Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. *Science (New York, N.Y.)* 306(5695):457–461.
- [133] Eizirik, D.L., Cnop, M., 2010. ER stress in pancreatic beta cells: the thin red line between adaptation and failure. *Science Signaling* 3(110):pe7.
- [134] Dromparis, P., Paulin, R., Stenson, T.H., Haromy, A., Sutendra, G., Michelakis, E.D., 2013. Attenuating endoplasmic reticulum stress as a novel therapeutic strategy in pulmonary hypertension. *Circulation* 127(1):115–125.
- [135] Wang, G., Liu, S., Wang, L., Meng, L., Cui, C., Zhang, H., et al., 2017. Lipocalin-2 promotes endoplasmic reticulum stress and proliferation by augmenting intracellular iron in human pulmonary arterial smooth muscle cells. *International Journal of Biological Sciences* 13(2):135–144.