



Article

Vitamin D Attenuates Oxidative Damage and Inflammation in Retinal Pigment Epithelial Cells

Ali Mohammad Tohari ^{1,2}, Reem Hasaballah Alhasani ², Lincoln Biswas ² , Sarita Rani Patnaik ², James Reilly ², Zhihong Zeng ^{3,*} and Xinhua Shu ^{2,4,*}

¹ Department of Clinical Biochemistry, King Fahad Hospital, PO Box 204, Jazan 91991, Saudi Arabia

² Department of Biological and Biomedical Sciences, Glasgow Caledonian University, Glasgow G4 0BA, UK

³ Department of Bioengineering and Environmental Science, Changsha University, Changsha 410022, China

⁴ Department of Vision Science, Glasgow Caledonian University, Glasgow G4 0BA, UK

* Correspondence: z20181201@ccsu.edu.cn (Z.Z.); Xinhua.Shu@gcu.ac.uk (X.S.)

Received: 21 June 2019; Accepted: 22 August 2019; Published: 24 August 2019



Abstract: Age-related macular degeneration (AMD), the most common visual disorder in elderly people, is characterized by the formation of deposits beneath the retinal pigment epithelium (RPE) and by dysfunction of RPE and photoreceptor cells. The biologically active form of vitamin D, 1,25-(OH)₂D₃ (VITD), is categorized as a multifunctional steroid hormone that modulates many transcriptional processes of different genes and is involved in a broad range of cellular functions. Epidemiological and genetic association studies demonstrate that VITD may have a protective role in AMD, while single nucleotide polymorphisms in the vitamin D metabolism gene (*CYP24A1*) increase the risk of AMD. However, the functional mechanisms of VITD in AMD are not fully understood. In the current study, we investigated the impact of VITD on H₂O₂-induced oxidative stress and inflammation in human RPE cells. We demonstrate that exposure to H₂O₂ caused significantly reduced cell viability, increased production of reactive oxygen species (ROS), lowered expression of antioxidant enzymes and enhanced inflammation. VITD exposure notably counteracted the above H₂O₂-induced effects. Our data suggest that VITD protects the RPE from oxidative damage and elucidate molecular mechanisms of VITD deficiency in the development of AMD.

Keywords: vitamin D; oxidative stress; inflammation; retinal pigment epithelial cells; age-related macular degeneration

1. Introduction

Age-related macular degeneration (AMD) is an accelerating neurodegenerative ocular disease that affects people above 55 years of age [1]. Epidemiologic studies have shown that there is an increasing number of new AMD cases diagnosed worldwide annually [2]. Wong et al. (2014) reported that the incidence of AMD is 7.1% in the African, 6.81% in the Asian, 11.19% in the Caucasian and 9.87% in the Hispanic population [3]. Several risk factors, such as smoking, have been associated with increased incidence of AMD [2]. A number of studies have suggested that the incidence of AMD is also linked to genetic defects [1,2]. It has been proposed that the AMD genetic association is strongly related to genetic defects in *ARMS2/HTRA1* and *CFH* located on, respectively, chromosome 10q26 and 1q32 [1,2]. Although the pathological mechanisms of AMD are not fully understood, many studies have suggested that AMD is an inflammatory process that involves increased production of many inflammatory mediators such as tumor necrosis factor- α (TNF- α), interleukins-1 β , 6, 8 and 33 (IL-1 β , IL-6, IL-8 and IL-33) [1,4], vascular endothelial growth factor A (VEGFA) [5] and activation of the alternative complement pathway [6].

Vitamin D is a fat-soluble secosteroid molecule that is found in two forms: vitamin D₂ and vitamin D₃ [7]. The latter is more abundant than vitamin D₂ in biological systems; thus, vitamin D₃ is commonly referred to simply as vitamin D. Humans and higher mammals can obtain vitamin D from various nutritional sources such as meats and fungi [7]. In addition, vitamin D can be produced endogenously. The endogenous synthesis of vitamin D starts in the epidermal skin layers where the vitamin D precursor 7-dehydrocholesterol (7-DHC) is converted to pre-vitamin D₃ through a photoisomerization process activated by exposure to type B ultraviolet radiation (UVB) between 280 and 320 nm, then incorporated with its carrier globin, vitamin D binding protein (DBP), and transported to the liver [7]. The pre-vitamin D₃ is transformed to 25-dihydroxyvitamin D₃ through the hepatic cytochrome P450 hydroxylase enzymes microsomal CYP2R1 and mitochondrial CYP2A1, attached to DBP and transported to the kidney [7]. Finally, 25-dihydroxyvitamin D₃ is converted to the active form 1,α,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) in renal cells by mitochondrial 1,α, CYP27B1. This step is tightly regulated by parathyroid hormone (PTH) and by circulating levels of calcium and phosphate, as well as by 1,25(OH)₂D₃ itself [7]. The molecule 1,α,25-dihydroxyvitamin D₃ is well recognized as a steroid hormone that regulates many cellular signalling activities through its nuclear vitamin D receptor (VDR) in target cells [7,8]. Vitamin D deficiency in the general population is currently considered a global health concern and represents a risk factor for the development of many diseases, particularly neurodegenerative diseases [9]. A large number of studies have shown that the likelihood of AMD is increased among those individuals with low circulating vitamin D levels [10–13].

IL-33 is a member of the IL-1 cytokine family that is expressed in many cell types of the nervous system [14]. IL-33 signalling is mediated through binding to its ST2 receptor in target cells and recruiting transcriptional cofactors such as IL-1R accessory protein (IL-1RAP), IL-1R-associated kinases 1/4 (IRAK1/4) and myeloid differentiation primary-response protein 88 (MYD88) into the transcription complex [14,15]. This cytokine has important roles in the inflammatory response in the modulation of neurodegenerative diseases, e.g., Alzheimer's disease [16] and AMD [4].

In the current study, we investigated the role of vitamin D in oxidative stress and inflammation. We found that vitamin D increased cell viability, reduced the formation of reactive oxygen species (ROS), and inhibited inflammation. Vitamin D also stimulated the expression of IL-33 at the mRNA and protein levels. Furthermore, IL-33 treatment markedly induced the expression of antioxidant genes. Our data suggest that the protective effects of vitamin D in AMD may be partially through the upregulation of IL-33.

2. Materials and Methods

2.1. Cell Culture

ARPE-19 cells (ATCC[®] CRL-2302[™], ATCC, Manassas, VA, USA) were grown in a T-25 cm² standard tissue culture flask in 5 mL of DMEM/F12 medium containing 2.4 mM L-Glutamine, 15 mM Hepes, 17.5 mM glucose, 10% FBS, 50 IU/mL of Penicillin and Streptomycin and 17.4 mM of 7.5% Sodium bicarbonate. The cells were maintained in a 5% CO₂ incubator at 37 °C. Once cell growth reached 80–90% confluency, the medium was removed and the cells were washed twice with PBS. The cells were then trypsinized with 1–1.5 mL of 0.5% trypsin–EDTA. Detached cells were resuspended in 4–5 mL of DMEM/F12 medium and seeded in 96- or 6-well tissue culture plates for 24 h at a density of 50,000 or 500,000 cells per well prior to cell treatment.

2.2. Cell Viability Assay

ARPE-19 cells were seeded at a density of 50,000 cells per well in a 96-well plate in DMEM/F12 medium for 24 h. The medium was then removed, and the cells were washed twice with 0.5 mL of 1× PBS/well. The cells were then treated with serial concentrations of H₂O₂ (150–2000 μM), 50 nM 1,α, 25(OH)₂D₃ (VITD), 0.1% ethanol for 6 and 24 h (or left untreated for the same time periods). In some experiments, ARPE-19 cells were treated with 750 μM H₂O₂, 750 μM H₂O₂ + 50 nM VITD, or left

untreated. All treatments were performed in serum-free DMEM/F12 medium. The cell viability assay was performed as previously described [17].

2.3. Caspase-3/7 Activity

ARPE-19 cells were seeded in a 96-well plate (20,000 cells/well) overnight and treated with 750 μM H_2O_2 , 750 μM H_2O_2 + 50 nM VITD, or 0.1% DMSO (control). Caspase-3/7 activity was measured by the Apo-ONE Homogeneous Caspase-3/7 kit (Promega, Southampton, UK) according to the manufacturer's instructions. Fluorescence was measured using a Fluostar Optima Plate Reader at an excitation wavelength of 499 nm and an emission wavelength of 521 nm. Caspase-3/7 activity was determined according to the following formula: % relative fluorescence = (fluorescence of treated cells – fluorescence of control cells)/fluorescence of control cells \times 100. All experiments were repeated three times.

2.4. Measurement of Reactive Oxygen Species (ROS)

ARPE-19 cells were cultured at a density of 25,000 cells per well in a 96-well plate and treated with 750 μM H_2O_2 , 750 μM H_2O_2 + 50 nM VITD, or 0.1% DMSO for 6 and 24 h. Production of ROS was measured using the 6-Carboxy-2',7'-Dichlorofluorescein diacetate (DCFH-DA) dye (Sigma, Dorset, UK). The medium was removed, and the cells were washed twice with 1 \times PBS, 200 μL /well. Then, 150 μL of 10 μM DCFH-DA in 1 \times PBS/well was added, and the plate was incubated for 45 min in a 5% CO_2 incubator at 37 $^\circ\text{C}$. Fluorescence was measured at 485 nm (excitation) and 525 nm (emission) using a Fluostar Optima microplate reader (BMG LABTECH Ltd, Bucks, UK). The ROS level was represented as the percentage of fluorescence intensity relative to the control after baseline fluorescence correction according to the following formula: (fluorescence of treated cells – fluorescence of control cells)/fluorescence of control cells \times 100. All experiments were repeated three times.

2.5. Enzyme-Linked Immunosorbent Assay (ELISA)

Total proteins from ARPE-19 cells, retina and RPE were extracted by homogenizing samples in 200 μL of T-PER buffer containing 1 \times protease Inhibitor Cocktail. The supernatants were collected by centrifugation at 10,000 rpm for 5 min. The concentration of IL-1 β , IL-8, IL-33, TNF α and VEGFA in control and treated human ARPE-19 cell media was measured using commercial ELISA kits from PeproTech and eBioscience according to the manufacturer's protocols. The levels of the examined cytokines were quantified using a standard curve constructed on the basis of the mean absorbance of standard solution concentrations of cytokines (X axis) against optical density (O.D.) for each standard solution (Y axis). The concentration of unknown samples (pg/mL) was calculated on the basis of the straight line equation obtained from the linear-regression trendline according to $Y = mx + c$ (where Y = O.D. of unknown sample, m = slope value, x = concentration of unknown sample and c = intercept). All experiments were repeated three times.

2.6. Biochemical Assay

Quantification of malondialdehyde (MDA) and glutathione (GSH) and measurement of CATALASE and superoxide dismutase (SOD) activities were performed using kits from CELL BIOLABS INC (San Diego, CA, USA) according to the manufacturer's instructions.

2.7. Mouse Samples

Mice used in this study were housed in the Animal Unit at Glasgow Caledonian University with free access to food and water and a 12 h light/dark environment. Mice at the age of 2 months, 9 months and 24 months were sacrificed, and the retina and RPE were dissected and stored at -80 $^\circ\text{C}$ for further analysis. Approval for animal use was granted by the Glasgow Caledonian University

Animal Ethics and Welfare Committee, in accordance with the UK home office animal care guidelines (Project licence P8C815DC9).

2.8. Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

Total RNAs from ARPE-19 cells and mouse tissues (retina and RPE) were extracted using Tri Reagent[®] (Sigma, Dorset, UK) following the manufacturer's instructions. The cDNA was synthesized using the High-Capacity cDNA Reverse Transcription kit (Thermo Fisher Scientific, Paisley, UK) as described by the manufacturer. Quantification of gene expression was performed using real-time PCR Platinum SYBR Green qPCR SuperMix-UDG with ROX assay, as described by the manufacturer. Briefly, 1.0 μ L of 50 ng/ μ L of cDNA was mixed with 7.5 μ L of Platinum SYBR Green qPCR SuperMix-UDG with ROX and 0.6 μ L of 10 μ M forward and reverse primers, and the reaction volume was scaled to 15 μ L with nuclease-free water. DNA amplification was carried out under the following conditions: 50 °C for 2 min (UDG incubation), followed by enzyme activation at 95 °C for 2 min, hold and then an amplification step of 40 cycles including DNA denaturation at 95 °C for 15 s, then primer annealing at 60 °C for 15 s. Fluorescence signals were detected at the end of the 60 °C step, and assay validity was assessed on the basis of the melting curve analysis following each run. Relative gene expression was determined according to the $2^{-\Delta\Delta Ct}$ formula. The primer sequences for qRT-PCR are available on request.

2.9. Immunostaining

ARPE-19 cells were fixed with methanol at -20 °C for 5 min, then washed with $1\times$ PBS twice. The cells were blocked with 2% BSA-PBS at room temperature for 30 min, then incubated with primary antibodies at 4 °C overnight. After washing three times (5 min each time), the cells were blocked again with 2% sheep serum in 2% BSA-PBS for 30 min. The cells were incubated with secondary antibodies at room temperature for 1 h, then washed 5 times with $1\times$ PBS (5 min each). The cells were mounted with DAPI solution and imaged under a confocal microscope.

2.10. Western Blot

Treated and control cells were lysed with Radio-Immuno Precipitation Assay (RIPA) buffer. Proteins were separated by SDS-PAGE and transferred to nitrocellulose membranes. The targeted proteins were detected using primary antibodies (NRF2, 1:1000; GAPDH, 1:1000) and secondary antibodies (1:10,000). The signals were quantified with the ImageStudio[™]Lite analysis software (LI-COR, Cambridge, UK).

2.11. Statistical Data Analysis

Data were analysed by one-way or two-way Anova followed by Bonferroni post-hoc test using GraphPad Prism version 6 software (GraphPad Software Inc., San Diego, CA, USA); $p < 0.05$ was considered significant. All experiments were repeated three times.

3. Results

3.1. VITD Treatment Improved Cell Viability and Reduced ROS Production and Apoptosis

In order to determine the appropriate concentration of H_2O_2 to induce a significant, though not excessive, toxic effect on cell viability, ARPE-19 cells were challenged with H_2O_2 at a range of concentrations (150 to 2000 μ M) for 6 or 24 h. We detected a significant reduction in cell viability in cells treated with 450, 600, 750 and 1000 μ M H_2O_2 for 6 and 24 h compared to the respective control cells. A concentration of H_2O_2 higher than 1000 μ M was determined to be highly toxic (Figure S1). Thus, 750 μ M H_2O_2 was used to treat ARPE-19 cells for all subsequent experiments in this study. We also chose 50 nM VITD for our current study on the basis of earlier in vitro studies [18,19]. We found that 50 nM VITD significantly increased the viability of treated ARPE-19 cells when compared to

untreated control cells (Figure S1). Previous studies reported that VITD treatment can protect cells and tissues from oxidative damage [20–23]. In the current study, treatment of ARPE-19 cells with 750 μM H_2O_2 for both 6 and 24 h caused a significant reduction in cell viability and a significant increase in ROS production (Figure 1A,B). The latter can activate apoptosis signalling pathways [24]. In the current study, caspase 3/7 activation, an indicator of apoptosis, was significantly increased in cells exposed to 750 μM H_2O_2 for both 6 and 24 h (Figure 1C). When H_2O_2 exposure was accompanied by a treatment with VITD, the above effects were reversed: there was a significant increase in cell viability, a significant decrease in ROS production and a significant decrease in caspase 3/7 activation, compared to the levels seen following treatment with H_2O_2 alone (Figure 1A–C). Increased ROS can promote lipid peroxidation, resulting in increased production of MDA and 4-hydroxynonal (4-HNE), which indicates local cell or tissue damage. We found that MDA production in ARPE-19 cells treated with H_2O_2 was significantly higher than that seen in untreated cells; co-treatment with VITD significantly counteracted this effect (Figure 1D).




Figure 1. Effects of vitamin D (VITD, $1,25(\text{OH})_2\text{D}_3$) on cell viability, reactive oxygen species (ROS) production, apoptosis and malondialdehyde (MDA) level. ARPE-19 cells were treated with H_2O_2 alone or with H_2O_2 + VITD for 6 h or 24 h. Cell viability (A), ROS production (B), Caspase 3/7, a biomarker of apoptosis (C), and MDA, a biomarker of lipid peroxidation, levels (D) were examined. Data are presented as the means \pm standard deviation (SD) of three independent experiments. UT: untreated control. ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

3.2. VITD Treatment Enhanced VDR Expression in Stressed ARPE-19 Cells

VDR is the principal nuclear receptor for VITD signalling in biological systems [7,8]. It is expressed in many cells including retinal and RPE cells [25–27]. Our current results show that VDR was expressed in untreated (control) ARPE-19 cells, in cells treated with H_2O_2 alone and in cells co-treated with H_2O_2 and VITD (Figure S2A). This expression was significantly reduced (upon both 6 and 24 h exposure) in ARPE-19 cells treated with 750 μM H_2O_2 compared to untreated cells; however, VDR expression was substantially increased (upon both 6 and 24 h exposure) in ARPE-19 cells co-treated with 750 μM H_2O_2 and 50 nM VITD compared to cells treated with 750 μM H_2O_2 alone (Figure S2B).

3.3. VITD Treatment Upregulated the Expression of Antioxidant Genes

Increased intracellular ROS production associated with many pathological conditions can cause major cellular injury, reducing cells' capacity to produce protective molecules such as anti-oxidant proteins [28]. Recent studies have shown that VITD treatment can modulate both gene expression and activity of many antioxidant molecules [29–31]. In the current study, we used qRT-PCR to examine the expression of *SOD1*, *SOD2*, *CATALASE (CAT)*, *GPX1*, *GPX2* and *GPX3* genes in ARPE-19 cells under our stress condition and VITD treatment. We found that, at both 6 and 24 h, the expression of *CAT*, *SOD1*, *SOD2*, *GPX2* and *GPX3* were noticeably reduced in ARPE-19 cells treated with 750 μM H_2O_2 compared to untreated cells; however, the expression of these antioxidant genes was significantly increased in ARPE-19 cells co-treated with 750 μM H_2O_2 and 50 nM VITD compared to cells treated with 750 μM H_2O_2 alone (Figure 2A). There was no significant difference at 6 or 24 h in *GPX1* expression in untreated cells, cells treated with 750 μM H_2O_2 alone, and cells co-treated with 750 μM H_2O_2 and 50 nM VITD (Figure 2A).




Figure 2. VITD treatment enhanced the expression of antioxidant genes. (A) ARPE-19 cells were exposed to H_2O_2 alone or H_2O_2 + VITD for 6 h or 24 h. The expression of *SOD1*, *SOD2*, *CATALASE (CAT)*, *GPX1*, *GPX2* and *GPX3* in treated and untreated cells was measured by qRT-PCR. (B) ARPE-19 cells were exposed to H_2O_2 alone or H_2O_2 + VITD for 24 h; the activities of SOD and CAT and GSH level were biochemically examined. Data are presented as the means \pm standard deviation (SD) of three independent experiments. UT: untreated control. ns: no significance. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

We also examined the effect of VITD treatment on SOD and CAT activities. The activities of SOD and CAT were significantly decreased in H₂O₂-treated cells compared with untreated control cells. The co-treatment of cells with 750 μM H₂O₂ and 50 nM VITD resulted in a significant increase in SOD and CAT activities compared to cells exposed to H₂O₂ alone (Figure 2B). GSH, an important intracellular antioxidant, can prevent ROS-induced damage. GSH level was significantly decreased in H₂O₂-treated cells compared to untreated control cells, while cells co-treated with VITD and H₂O₂ showed a significant increase in GSH (compared to cells treated with H₂O₂ alone (Figure 2B).

3.4. VITD Treatment Modulated the Expression of the Inflammatory Mediators

Recent studies have suggested that AMD is an inflammatory disease that is associated with increased levels of pro-inflammatory cytokines and the presence of macrophages and other immune cells [1]. VITD treatment has been shown to regulate the gene expression of many inflammatory mediators in various biological systems [32]. In this study, we investigated the expression of *IL-1β*, *IL-8*, *TNF-α* and *VEGFA* genes in ARPE-19 cells treated with 750 μM H₂O₂, in cells co-treated with 750 μM H₂O₂ and 50 nM VITD, and in untreated control cells. At both 6 and 24 h, the expression levels of *IL-1β*, *TNF-α*, *IL-8*, *IL-18* and *VEGFA* genes were significantly increased in ARPE-19 cells treated with H₂O₂ compared to untreated cells; by contrast, at both 6 and 24 h, their expression levels were substantially reduced in ARPE-19 cells co-treated with 750 μM H₂O₂ and 50 nM VITD compared to cells treated with 750 μM H₂O₂ alone (Figure 3A). We also used ELISA to measure the levels of secreted IL-1β, IL-8, TNF-α and VEGFA in the media of treated and control cells and found that all the examined cytokines were significantly increased in ARPE-19 cells treated with 750 μM H₂O₂ when compared to control cells; co-treatment with H₂O₂ and VITD markedly decreased the levels of these cytokines when compared to cells exposed to H₂O₂ alone (Figure 3B).




Figure 3. Cont.




Figure 3. VITD inhibited inflammation. (A) ARPE-19 cells were exposed to H₂O₂ alone or to H₂O₂ + VITD for 6 h or 24 h; the expression of IL-1 β , IL-8, TNF- α and VEGF in treated and untreated cells was measured by qRT-PCR. (B) ARPE-19 cells were exposed to H₂O₂ alone or to H₂O₂ + VITD for 24 h. The levels of IL-1 β , IL-8, TNF- α and VEGF were measured by ELISA. Data are presented as the means \pm standard deviation (SD) of three independent experiments. UT: untreated control. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

3.5. VITD Reversed the H₂O₂-Induced Change in NRF2 Expression

NRF2 is a redox-sensitive transcription factor that can bind antioxidant elements and activates the expression of antioxidant and detoxifying enzymes. VITD has been reported to control NRF2 expression [32]. In the current study, when ARPE-19 cells were treated with 750 μ M H₂O₂, NRF2 protein level was significantly decreased compared to untreated control cells; co-treatment with VITD and 750 μ M H₂O₂ resulted in a marked increase in NRF2 expression when compared to H₂O₂ treatment alone (Figure 4).




Figure 4. H₂O₂ treatment downregulated NRF2 expression, while VITD reversed the effect. (A) ARPE-19 cells were treated with H₂O₂ alone or H₂O₂ + VITD for 24 h; the protein levels of NRF2 in treated and control cells were examined by Western blotting. (B) NRF2 protein levels were quantified by normalizing with respect to GAPDH protein. Data are presented as the means \pm standard deviation (SD) of three independent experiments. UT: untreated control. * $p < 0.05$, **** $p < 0.0001$.

3.6. VITD Regulated IL-33 Expression

Previous studies have reported that IL-33 can function as both a proinflammatory and an inflammatory cytokine during the immune response [15], as anti-apoptotic and survival molecules [33] can protect from oxidative stress by increasing SOD activity [34] and can attenuate the development of autoimmune uveitis [35]. IL-33 and its receptor ST2 are abundantly expressed in many tissues such as the central nervous system and the retina [14,35,36]. We used immunostaining to detect IL-33 and ST2 expression in ARPE-19 cells and found that IL-33 was localised in the nucleus (Figure 5A), while ST2 was localised in the cytoplasm, predominantly around the nuclear membrane (Figure 5B). We also measured IL-33 expression in mouse retina and RPE: the expression of IL-33 decreased in the retina and RPE during ageing (Figure S3). We then examined whether H₂O₂ and VITD affect IL-33 expression. Using qRT-PCR, we measured the mRNA level of IL-33 in treated and control cells and found that IL-33 expression was significantly decreased in ARPE-19 cells treated with 750 μM of H₂O₂ compared to untreated cells, while it was significantly increased in ARPE-19 cells co-treated with 750 μM H₂O₂ and 50 nM VITD compared to cells treated with 750 μM H₂O₂ alone. VITD treatment alone for 6 and 24 h also significantly increased IL-33 expression when compared to control cells (Figure 5C). We also measured the protein level of IL-33 by ELISA: H₂O₂ treatment significantly decreased IL-33 level compared to control cells, while co-treatment with H₂O₂ and VITD notably counteracted this change. Exposure to VITD alone resulted in a significantly increased IL-33 level compared to the control cells (Figure 5D).




Figure 5. Localization of IL-33 (A) and ST-2 (B) in ARPE-19 cells using immunostaining with anti-IL-33 antibody and anti-ST2 antibody, respectively (63×). (C) Effects of VITD on IL-33 expression in ARPE-19 cells treated with VITD only, H₂O₂ alone or H₂O₂ + VITD for 6 h or 24 h. IL-33 mRNA levels

were quantified by qRT-PCR. (D) VITD treatment increased IL-33 protein production. ARPE-19 cells were incubated with VITD only, H₂O₂ alone or H₂O₂ + VITD for 24 h. Secreted IL-33 protein was measured by ELISA. Data are presented as the means ± standard deviation (SD) of three independent experiments. UT: untreated control. * $p < 0.05$, ** $p < 0.01$.

3.7. IL-33 Treatment Upregulated Antioxidant Gene Expression

There is growing evidence that IL-33 can protect cells from apoptosis and oxidative stress and can enhance cell survival in vitro and in vivo [33,37]. The protective function of IL-33 against apoptosis and oxidative stress is believed to occur through its action on the gene expression and production of anti-oxidant scavengers such as superoxide dismutases [34]. In ARPE-19 cells treated with rhIL-33 (10 or 50 ng/mL), the expression of *CAT*, *SOD1*, *SOD2*, *GPX2* and *GPX3* was significantly increased following 6 and 24 h treatments. However, there was no significant change in the expression of *GPX1* following treatment with either 10 or 50 ng/mL rhIL (Figure 6).




Figure 6. Effects of recombinant human IL-33 on the expression of antioxidant genes. ARPE-19 cells were treated with recombinant human IL-33 (rhIL-33) for 6 or 24 h. The expression of antioxidant genes (*SOD1*, *SOD2*, *CAT*, *GPX1*, *GPX2* and *GPX3*) was measured by qRT-PCR. Data are presented as the means ± standard deviation (SD) of three independent experiments; ns, no significance. * $p < 0.05$.

4. Discussion

It is well established that oxidative stress, inflammation and angiogenesis play a critical role in the development and progression of AMD [38]. VITD has multiple functions including the inhibition of oxidative stress, inflammation, macrophage activation and angiogenesis, thus offering therapeutic potential for AMD. Human RPE cell lines can synthesize active vitamin D (VITD), while VDR and vitamin D synthesis enzymes and metabolism enzymes are also expressed in the neuroretina, RPE and choroid [38]. Early epidemiological studies showed a connection between vitamin D and AMD, while genome-wide association studies demonstrated that the vitamin D metabolism gene (*CYP24A1*) is a risk factor for AMD [38–40]. Lee et al. (2012) reported that Vitamin D treatment resulted in decreased retinal inflammation and A β deposition and increased visual function in aged mice [41]. Previously, we reported that VITD can protect cone cells from oxidative damage [17]. Our current

data also demonstrate that VITD suppresses oxidative damage and inflammation in RPE cells, further supporting the hypothesis that VITD may play an important role in AMD.

Although the principal function of the active form of VITD [$1,\alpha,25\text{-(OH)}_2\text{D}_3$] is the regulation of calcium and phosphate homeostasis, it also acts as a steroid hormone, exerting its effect through VDR signalling in target cells. The VDR gene is expressed in a number of ocular structures such as the retina, the cornea and the RPE-choroid and in ocular cell lines such as ARPE-19 and human retinoblastoma (Y79) [25,39], thus demonstrating its regulatory function in the ocular system. VDR is one of the nuclear receptors of steroid hormones that mediate transcriptional processes in a variety of cells. Expression and production of VDR are influenced by factors such as sun exposure, hormones levels and age. VITD itself also regulates VDR expression through two gene expression enhancers containing vitamin D₃ response element (VDRE): CCAAT/enhancer binding protein- β (C/EBP β) and runt-related transcription factor-2 (RUNX2) located upstream of the VDR gene's transcriptional start site (TSS) [42,43]. We also confirmed that VITD stimulated VDR expression in ARPE-19 cells (data not shown) and found that H₂O₂ treatment significantly decreased VDR expression, while co-treatment with VITD reversed the H₂O₂-caused change (Figure S2).

ROS are highly reactive compounds that cause cellular damage through the oxidation of lipids, proteins and DNA, altering their biochemical characteristics and causing reduced cell viability and subsequent cell death [44]. Many studies have suggested that VITD can improve cell viability and protect against the deleterious effects of ROS in various cell lines [20,22,23]. Similarly, in the current study, a significant reduction of ARPE-19 cell viability was observed in cells treated with H₂O₂ at concentrations ranging between 450 and 2000 μM (Figure S1). The reduction in cell viability following H₂O₂ treatment was also associated with increased levels of free radicals (ROS) (Figure 1A,B). However, ROS levels were substantially decreased when ARPE-19 cells were treated with a combination of 750 μM of H₂O₂ and 50 nM VITD, suggesting that VITD suppressed H₂O₂-induced ROS production (Figure 1B). VITD treatment also inhibited lipid peroxidation and reduced cell death (Figure 1C,D). We also found that H₂O₂ exposure reduced the expression of antioxidant genes (*SOD1*, *SOD2*, *CAT*, *GPX1*, *GPX2*) and induced the production of inflammatory cytokines (IL-1 β , IL-8, TNF- α and VEGF). Co-treatment with VITD counteracted these H₂O₂-induced effects (Figure 2; Figure 3). These protective effects of VITD against H₂O₂-induced damage can be explained by its effect on the gene and protein expression of many antioxidants via upregulation of different transcription factors, e.g., NRF2, a key transcription factor that modulates the expression of many antioxidants [32]. Our data also demonstrated that H₂O₂ caused a significant decrease in NRF2 expression, while co-treatment with VITD markedly increased NRF2 expression (Figure 4). Although VITD plays a critical role in cell proliferation [45] and, as we found, VITD treatment significantly increased cell viability (Figure S1), the current study suggests that the protection of VITD against H₂O₂-induced toxicity is through antioxidant signalling pathways.

IL-33 is an important immune mediator in various disorders including neurodegenerative diseases [46]. IL-33 is abundantly expressed in many cell types. Recent studies have reported that IL-33 is expressed in human and mouse RPE and neuroretina [4,35,36]. Our data also demonstrated that both IL-33 and ST2 are expressed in ARPE-19 cells, an observation in accordance with previous reports [4]. Interestingly, IL-33 expression was notably decreased in aged mouse neuroretina and RPE (Figure S3). Liu et al. (2012) reported that amyloid- β 1-40 oligomer stimulated IL-33 expression in a human RPE (D407) cell line [47]. However, the data from the present study showed that the mRNA and protein levels of IL-33 were significantly lower in ARPE-19 cells treated with H₂O₂, while both IL-33 mRNA expression and protein levels were notably increased following VITD treatment (Figure 5C,D). Increased IL-33 expression has been reported in the vitreous and in retinal Müller cells and mononuclear cells of AMD patients, suggesting that IL-33 may play a pathogenic role in AMD [36]. However, Theodoropoulou et al. (2017) reported that IL-33 regulates tissue remodelling and suppresses murine choroidal neovascularization, a clinical feature of AMD, suggesting that IL-33 may confer some protection in AMD [4]. Additionally, glia-derived IL-33 has been shown to induce myeloid cell infiltration and enhance the recovery of the central nervous system from acute injury [14].

In an experimental autoimmune uveitis mouse model, IL-33 slows down disease progression with decreased retinal inflammation [35]. However, in injured mouse retina, Müller cells have been shown to increase IL-33 production, promoting the release of inflammatory cytokines and chemokines and the recruitment of the myeloid cells, which contribute to photoreceptor cell loss [36].

In the current study, we used co-treatment with VITD and H₂O₂ to assess the protection of VITD against H₂O₂-induced toxicity. Other researchers have evaluated VITD protection in in vitro cell lines by pre-treatment with VITD [20,23]. Ideally, we would also examine the protection of VITD against H₂O₂-induced toxicity by pre-treatment with VITD in ARPE-19 cells. Further preclinical tests in AMD animal models and AMD patient clinical trials are needed to verify the therapeutic potential of VITD for AMD.

5. Conclusions

In summary, VITD can possibly protect the retina and RPE from oxidative stress, inflammation and apoptosis through the suppression of pro-inflammatory mediators and by enhancing the antioxidant defence capacity.

Supplementary Materials: The following are available online at <http://www.mdpi.com/2076-3921/8/9/341/s1>, Figure S1: Effects of VITD and H₂O₂ on cell viability. ARPE-19 cells were exposed to VITD or H₂O₂ for 6 hours (A) or 24 hours (B). Cell viability was measured using MTT assay, Figure S2: (A) Expression of vitamin D receptor (VDR) was detected in untreated (UT) and treated (H₂O₂ alone or H₂O₂ + VITD) ARPE-19 cells by end-point PCR. PCR products were analyzed using agarose gel electrophoresis. (B) ARPE-19 cells were exposed to H₂O₂ or H₂O₂ + VITD for 6 hours or 24 hours; expression of VDR was measured by qRT-PCR. Figure S3: Expression of IL-33 in the retinas (A) and RPE (B) of mice at three age points was measured by qRT-PCR.

Author Contributions: X.S. designed the study. A.M.T., R.H.A., L.B. and S.R.P. performed the experiments. A.M.T. and X.S. analysed the data. J.R., Z.Z. and X.S. wrote the manuscript.

Funding: This work was supported by a PhD scholarship (A826) from Saudi Arabia Government. The work was also supported by the Rosetrees Trust (M160, M160-F1, M160-F2) and National Eye Research Centre (SAC037).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Cascella, R.; Ragazzo, M.; Strafella, C.; Missiroli, F.; Borgiani, P.; Angelucci, F.; Marsella, L.; Cusumano, A.; Novelli, G.; Ricci, F.; et al. Age-Related Macular Degeneration: Insights into Inflammatory Genes. *J. Ophthalmol.* **2014**, *2014*, 582842. [[CrossRef](#)] [[PubMed](#)]
2. Priya, R.R.; Chew, E.Y.; Swaroop, A. Genetic studies of age-related macular degeneration: Lessons, challenges, and opportunities for disease management. *Ophthalmology* **2012**, *119*, 2526–2536. [[CrossRef](#)] [[PubMed](#)]
3. Wong, W.L.; Su, X.; Li, X.; Cheung, C.M.; Klein, R.; Cheng, C.Y.; Wong, T.Y. Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: A systematic review and meta-analysis. *Lancet Glob. Health* **2014**, *2*, e106–e116. [[CrossRef](#)]
4. Theodoropoulou, S.; Copland, D.A.; Liu, J.; Wu, J.; Gardner, P.J.; Ozaki, E.; Doyle, S.L.; Campbell, M.; Dick, A.D. Interleukin-33 regulates tissue. *J. Pathol.* **2017**, *241*, 45–56. [[CrossRef](#)] [[PubMed](#)]
5. Holz, F.G.; Schmitz-Valckenberg, S.; Fleckenstein, M. Recent developments in the treatment of age-related macular degeneration. *J. Clin. Investig.* **2014**, *124*, 1430–1438. [[CrossRef](#)] [[PubMed](#)]
6. Toomey, C.B.; Kelly, U.; Saban, D.R.; Bowes Rickman, C. Regulation of age-related macular degeneration-like pathology by complement factor H. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E3040–E3049. [[CrossRef](#)]
7. Bikle, D.D. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem. Biol.* **2014**, *21*, 319–329. [[CrossRef](#)] [[PubMed](#)]
8. Krishnan, A.V.; Feldman, D. Mechanisms of the anti-cancer and anti-inflammatory actions of vitamin D. *Annu. Rev. Pharmacol. Toxicol.* **2011**, *51*, 311–336. [[CrossRef](#)]
9. Fernandes de Abreu, D.A.; Eyles, D.; Féron, F. Vitamin D, a neuro-immunomodulator: Implications for neurodegenerative and autoimmune diseases. *Psychoneuroendocrinology* **2009**, *34*, S265–S277. [[CrossRef](#)]
10. Graffe, A.; Beauchet, O.; Fantino, B.; Milea, D.; Annweiler, C. Vitamin and macular thickness in the elderly: An optical coherence tomography study. *Investig. Ophthalmol. Vis. Sci.* **2014**, *15*, 5298–5303. [[CrossRef](#)]

11. Itty, S.; Day, S.; Lyles, K.W.; Stinnett, S.S.; Vajzovic, L.M.; Mruthyunjaya, P. Vitamin deficiency in neovascular versus nonneovascular age-related macular degeneration. *Retina* **2014**, *34*, 1779–1786. [[CrossRef](#)]
12. Millen, A.E.; Volland, R.; Sondel, S.A.; Parekh, N.; Horst, R.L.; Wallace, R.B.; Hageman, G.S.; Chappell, R.; Blodi, B.A.; Klein, M.L.; et al. Vitamin D status and early age-related macular degeneration in postmenopausal women. *Arch. Ophthalmol.* **2011**, *129*, 481–489. [[CrossRef](#)]
13. Singh, A.; Falk, M.K.; Subhi, Y.; Sørensen, T.L. The association between plasma 25-hydroxyvitamin D and subgroups in age-related macular degeneration: A cross-sectional study. *PLoS ONE* **2013**, *29*, e70948. [[CrossRef](#)]
14. Gadani, S.P.; Walsh, J.T.; Smirnov, I.; Zheng, J.; Kipnis, J. The glia-derived alarmin IL-33 orchestrates the immune response and promotes recovery following CNS injury. *Neuron* **2015**, *85*, 703–709. [[CrossRef](#)]
15. Liew, F.; Pitman, N.; McInnes, I. Disease-associated functions of IL-33: The new kid in the IL-1 family. *Nat. Rev. Immunol.* **2010**, *10*, 103–110. [[CrossRef](#)]
16. Yu, J.T.; Song, J.H.; Wang, N.D.; Wu, Z.C.; Zhang, Q.; Zhang, N.; Zhang, W.; Xuan, S.Y.; Tan, L. Implication of gene polymorphism in Chinese patients with Alzheimer’s disease. *Neurobiol. Aging* **2012**, *33*, 1014.e11–1014.e14. [[CrossRef](#)]
17. Tohari, A.M.; Zhou, X.; Shu, X. Protection against oxidative stress by vitamin D in cone cells. *Cell Biochem. Funct.* **2016**, *34*, 82–94. [[CrossRef](#)]
18. Chen, Y.; Zhang, J.; Ge, X.; Du, J.; Deb, D.; Li, Y. Vitamin D receptor inhibits nuclear factor κ B activation by interacting with I κ B kinase β protein. *J. Biol. Chem.* **2013**, *288*, 19450–19458. [[CrossRef](#)]
19. Huang, Y.; Ho, Y.; Lai, C.; Chiu, C.; Wang, J. 1,25-Dihydroxyvitamin D3 attenuates endotoxin-induced production of inflammatory mediators by inhibiting MAPK activation in primary cortical neuron-glia cultures. *J. Neuroinflamm.* **2015**, *12*, 147. [[CrossRef](#)]
20. Bao, B.Y.; Ting, H.J.; Hsu, J.W.; Lee, Y.F. Protective role of 1 alpha, 25-dihydroxyvitamin D3 against oxidative stress in human prostate epithelial cells. *Int. J. Cancer* **2008**, *122*, 2699–2706. [[CrossRef](#)]
21. Hamden, K.; Carreau, S.; Jamoussi, K.; Miladi, S.; Lajmi, S.; Aloulou, D.; Ayadi, F.; Elfeki, A. 1 Alpha dihydroxyvitamin D3: Therapeutic and preventive effects against oxidative stress, hepatic, pancreatic and renal injury in alloxan-induced diabetes in rats. *J. Nutr. Sci. Vitamin.* **2009**, *55*, 215–222. [[CrossRef](#)]
22. Peng, X.; Vaishnav, A.; Murillo, G.; Alimirah, F.; Torres, K.E.; Mehta, R.G. Protection against cellular stress by 25-hydroxyvitamin D3 in breast epithelial cells. *J. Cell Biochem.* **2010**, *110*, 1324–1333. [[CrossRef](#)]
23. Polidoro, L.; Properzi, G.; Marampon, F.; Gravina, G.L.; Festuccia, C.; Di Cesare, E.; Scarsella, L.; Ciccarelli, C.; Zani, B.M.; Ferri, C. Vitamin D protects human endothelial cells from H₂O₂ oxidant injury through the Mek/Erk-Sirt1 axis activation. *J. Cardiovasc. Transl. Res.* **2013**, *6*, 221–231. [[CrossRef](#)]
24. Redza-Dutordoir, M.; Averill-Bates, D.A. Activation of apoptosis pathways by reactive oxygen species. *Biochim. Biophys. Acta* **2016**, *1863*, 2977–2992. [[CrossRef](#)]
25. Alsalem, J.; Patel, D.; Susarla, R.; Coca-Prados, M.; Bland, R.; Walker, E.; Rauz, S.; Wallace, G. Characterization of vitamin D production by human ocular barrier cells. *Investig. Ophthalmol. Vis. Sci.* **2014**, *55*, 2140–2141. [[CrossRef](#)]
26. Dwyer, M.; Kazmin, D.; Hu, P.; McDonnell, D.; Malek, G. Research resource: Nuclear receptor atlas of human retinal pigment epithelial cells: Potential relevance to age-related macular degeneration. *Mol. Endocrinol.* **2011**, *25*, 360–372. [[CrossRef](#)]
27. Ren, Z.; Li, W.; Zhao, Q.; Ma, L.; Zhu, J. The impact of 1,25-dihydroxy vitamin D3 on the expressions of vascular endothelial growth factor and transforming growth factor- β 1 in the retinas of rats with diabetes. *Diabetes Res. Clin. Pract.* **2012**, *98*, 474–480. [[CrossRef](#)]
28. Bhat, A.H.; Dar, K.B.; Anees, S.; Zargar, M.A.; Masood, A.; Sofi, M.A.; Ganie, S.A. Oxidative stress, mitochondrial dysfunction and neurodegenerative diseases: A mechanistic insight. *Biomed. Pharmacother.* **2015**, *74*, 101–110. [[CrossRef](#)]
29. Nakai, K.; Fujii, H.; Kono, K.; Goto, S.; Kitazawa, R.; Kitazawa, S.; Hirata, M.; Shinohara, M.; Fukagawa, M.; Nishi, S. Vitamin D activates the Nrf2-Keap1 antioxidant pathway and ameliorates nephropathy in diabetic rats. *Am. J. Hypertens.* **2013**, *27*, 586–595. [[CrossRef](#)]
30. Middleton, R.P.; Nelson, R.; Li, Q.; Blanton, A.; Labuda, J.A.; Vitt, J.; Inpanbutr, N. 1,25-Dihydroxyvitamin D3 and its analogues increase catalase at the mRNA, protein and activity level in a canine transitional carcinoma cell line. *Vet. Comp. Oncol.* **2015**, *13*, 452–463. [[CrossRef](#)]

31. Eelen, G.; Verlinden, L.; Meyer, M.; Gijbbers, R.; Pike, J.; Bouillon, R.; Verstuyf, A. 1,25-Dihydroxyvitamin D₃ and the aging-related forkhead box O and sestrin proteins in osteoblasts. *J. Steroid Biochem. Mol. Biol.* **2103**, 136, 112–119. [[CrossRef](#)]
32. Berridge, M.J. Vitamin D: A custodian of cell signalling stability in health and disease. *Biochem. Soc. Trans.* **2015**, *43*, 349–358. [[CrossRef](#)]
33. Seki, K.; Sanada, S.; Kudinova, A.; Steinhäuser, M.; Handa, V.; Gannon, J.; Lee, R. Interleukin-33 prevents apoptosis and improves survival after experimental myocardial infarction through ST2 signaling. *Circ. Heart Fail.* **2009**, *2*, 684–691. [[CrossRef](#)]
34. Zhang, H.F.; Xie, S.L.; Chen, Y.X.; Mai, J.T.; Wang, J.F.; Zhu, W.L.; Zhu, L.G. Altered serum levels of IL-33 in patients with advanced systolic chronic heart failure: Correlation with oxidative stress. *J. Transl. Med.* **2012**, *8*, 120. [[CrossRef](#)]
35. Barbour, M.; Allan, D.; Xu, H.; Pei, C.; Chen, M.; Niedbala, W.; Fukada, S.Y.; Besnard, A.G.; Alves-Filho, J.C.; Tong, X.; et al. IL-33 attenuates the development of experimental autoimmune uveitis. *Eur. J. Immunol.* **2014**, *44*, 3320–3329. [[CrossRef](#)]
36. Xi, H.; Katschke, K.J.; Li, Y.; Truong, T.; Lee, W.P.; Diehl, L.; Rangell, L.; Tao, J.; Arceo, R.; Eastham-Anderson, J.; et al. IL-33 amplifies an innate immune response in the degenerating retina. *J. Exp. Med.* **2016**, *213*, 189–207. [[CrossRef](#)]
37. Rui, T.; Zhang, J.; Xu, X.; Yao, Y.; Kao, R.; Martin, C. Reduction in IL-33 expression exaggerates ischaemia/reperfusion-induced myocardial injury in mice with diabetes mellitus. *Cardiovasc. Res.* **2012**, *94*, 370–378. [[CrossRef](#)]
38. Layana, A.; Minnella, A.; Garhöfer, G.; Aslam, T.; Holz, F.; Leys, A.; Silva, R.; Delcourt, C.; Souied, E.; Seddon, J. Vitamin D and Age-Related Macular Degeneration. *Nutrients* **2017**, *9*, 1120. [[CrossRef](#)]
39. Morrison, M.A.; Silveira, A.C.; Huynh, N.; Jun, G.; Smith, S.E.; Zacharaki, F.; Sato, H.; Loomis, S.; Andreoli, M.T.; Adams, S.M.; et al. Systems biology-based analysis implicates a novel role for vitamin D metabolism in the pathogenesis of age-related macular degeneration. *Hum. Genom.* **2011**, *5*, 538–568. [[CrossRef](#)]
40. Parekh, N.; Chappell, R.J.; Millen, A.E.; Albert, D.M.; Mares, J.A. Association between vitamin D and age-related macular degeneration in the Third National Health and Nutrition Examination Survey, 1988 through 1994. *Arch. Ophthalmol.* **2007**, *125*, 661–669. [[CrossRef](#)]
41. Lee, V.; Rekh, E.; Hoh Kam, J.; Jeffery, G. Vitamin D rejuvenates aging eyes by reducing inflammation, clearing beta and improving visual function. *Neurobiol. Aging* **2012**, *33*, 2382–2389. [[CrossRef](#)]
42. Christakos, S.; Dhawan, P.; Peng, X.; Obukhov, A.; Nowycky, M.; Benn, B.; Zhong, Y.; Liu, Y.; Shen, Q. New insights into the function and regulation of vitamin D target proteins. *J. Steroid Biochem. Mol. Biol.* **2007**, *103*, 405–410. [[CrossRef](#)]
43. Zella, L.; Kim, S.; Shevde, N.; Pike, J. Enhancers located within two introns of the vitamin D receptor gene mediate transcriptional autoregulation by 1,25-dihydroxyvitamin D₃. *Mol. Endocrinol.* **2006**, *20*, 1231–1247. [[CrossRef](#)]
44. Eshaq, R.; Wright, W.; Harris, N. Oxygen delivery, consumption, and conversion to reactive oxygen species in experimental models of diabetic retinopathy. *Redox Biol.* **2014**, *2*, 661–666. [[CrossRef](#)]
45. Samuel, S.; Sitrin, M.D. Vitamin D's role in cell proliferation and differentiation. *Nutr. Rev.* **2008**, *66*, S116–S124. [[CrossRef](#)]
46. Du, L.X.; Wang, Y.Q.; Hua, G.Q.; Mi, W.L. IL-33/ST2 pathway as a rational therapeutic target for CNS diseases. *Neuroscience* **2018**, *369*, 222–230. [[CrossRef](#)]
47. Liu, X.C.; Liu, X.F.; Jian, C.X.; Li, C.J.; He, S.Z. IL-33 is induced by amyloid- β stimulation and regulates inflammatory cytokine production in retinal pigment epithelium cells. *Inflammation* **2012**, *35*, 776–784. [[CrossRef](#)]

