





Macrophage reprogramming for therapy

Valentina M. T. Bart,¹ 
 Robert J. Pickering,^{2,3} 
 Philip R. Taylor^{1,4}  and
 Natacha Ipseiz¹ 

¹Systems Immunity Research Institute, Cardiff University, Cardiff, UK, ²Immunology Network, Adaptive Immunity Research Unit, GlaxoSmithKline, Stevenage, UK, ³Department of Medicine, School of Clinical Medicine, Addenbrooke's Hospital, University of Cambridge, Cambridge, UK and ⁴UK Dementia Research Institute at Cardiff, Cardiff University, Cardiff, UK

doi:10.1111/imm.13300

Received 2 October 2020; revised 2 December 2020; accepted 11 December 2020.

VMTB and RJP contributed equally as co-first author.

PRT and NI contributed equally as co-last author.

Correspondence: Philip R. Taylor, Systems Immunity Research Institute, Heath Park, Cardiff University, Cardiff, CF14 4XN, UK.
 Email: TaylorPR@cardiff.ac.uk
 Senior author: Natacha Ipseiz
 Email: ipseizn@cardiff.ac.uk

Introduction

Historically considered as broad-range phagocytes playing a relatively 'passive' role within the immune system, macrophages (MΦ) have since benefited from more recent intensive characterization. Present in almost every tissue,

Summary

Dysfunction of the immune system underlies a plethora of human diseases, requiring the development of immunomodulatory therapeutic intervention. To date, most strategies employed have been focusing on the modification of T lymphocytes, and although remarkable improvement has been obtained, results often fall short of the intended outcome. Recent cutting-edge technologies have highlighted macrophages as potential targets for disease control. Macrophages play central roles in development, homeostasis and host defence, and their dysfunction and dysregulation have been implicated in the onset and pathogenesis of multiple disorders including cancer, neurodegeneration, autoimmunity and metabolic diseases. Recent advancements have led to a greater understanding of macrophage origin, diversity and function, in both health and disease. Over the last few years, a variety of strategies targeting macrophages have been developed and these open new therapeutic opportunities. Here, we review the progress in macrophage reprogramming in various disorders and discuss the potential implications and challenges for macrophage-targeted approaches in human disease.

Keywords: macrophages; polarization; reprogramming.

MΦ have been divided into two broad subclasses: those derived from an embryonic progenitor and those from adult monocytes. Many tissue-resident MΦ have a prenatal origin (yolk sac- or fetal liver-derived), their development is dependent on at least one essential tissue-specific transcription factor, and they maintain themselves by

Abbreviations: AAV, adeno-associated virus; AD, Alzheimer's disease; ALS, amyotrophic lateral sclerosis; BBB, blood–brain barrier; BMDM, bone marrow-derived macrophage; CIA, collagen-induced arthritis; CNS, central nervous system; DC, dendritic cell; EAE, experimental autoimmune encephalomyelitis; EGFR, epidermal growth factor receptor; FOXO3, forkhead box transcription factor family O 3; GA, glatiramer acetate; GM-CSF, granulocyte–macrophage CSF; HDAC, histone deacetylase; ID3, inhibitor of DNA 3; IFN-γ, interferon-gamma; IL, interleukin; IRF5, interferon regulatory factor 5; KC, Kupffer cell; LXR-α, liver X receptor-alpha; mAb, monoclonal antibody; MBC, metastatic breast cancer; MCRPC, metastatic castration-resistant prostate cancer; M-CSF, macrophage colony-stimulating factor; M-CSFR, M-CSF receptor; MHCI, major histocompatibility class I; MI, myocardial infarction; mo-KC, monocyte-derived KC; MS, multiple sclerosis; MΦ, macrophage; NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; NO, nitric oxide; NRF2, nuclear factor erythroid 2-related factor 2; PDAC, pancreatic ductal adenocarcinoma; PGM, proneural glioblastoma multiforme; RA, rheumatoid arthritis; scRNAseq, single-cell RNA sequencing; TAM, tumour-associated macrophages; TGF-β1, transforming growth factor beta 1; TLR, Toll-like receptor; TNBC, triple negative breast cancer; TREM2, triggering receptor expressed on myeloid cells-2; TTP, tristetraprolin; VEGF, vascular endothelial growth factor; Ywhaz, tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta

self-renewal, while others are recruited from the peripheral monocyte pool.¹ The origins and replenishment of MΦ populations have been the subject of numerous reviews²⁻⁶ and are not the focus of the present one. Tissue-resident MΦ are the predominant type of MΦ present during steady state and are thought to monitor tissues, and maintain homeostasis, cellular communication and immune surveillance.^{6,7} They also participate in developmental processes during embryogenesis.⁸⁻¹¹ Upon inflammation, whether induced by infection or injury, MΦ are recruited in large numbers from circulating monocytes to the tissue and are often loosely classified as pro- or anti-inflammatory. Previously, based on the actions of interferon-gamma (IFN-γ) and interleukin (IL)-4 on MΦ activation and with analogy to the Th1/2 T-cell subsets, MΦ were divided into two subtypes: 'M1' MΦ, 'classically activated' by IFN-γ; and 'M2' MΦ, 'alternatively activated' by type 2 anti-inflammatory cytokines such as IL-10. Those two subtypes of MΦ exhibit distinct metabolic function, the 'M1' having an anaerobic profile, based on glycolysis and production of nitric oxide (NO), whereas the 'M2' MΦ have an aerobic one, based on oxidative phosphorylation and production of arginase.¹²⁻¹⁴ However, with current knowledge on MΦ origin, diversity and significant plasticity,^{15,16} the acceptance of this dogma has diminished. Recent cutting-edge technological developments such as single-cell RNA sequencing (scRNAseq), advanced animal genetic modification and intravital microscopy¹⁷⁻²⁰ have allowed for a greater appreciation of cellular diversity than previously acknowledged. MΦ not only have impressive variability in their gene expression but they are also phenotypically plastic, allowing them to adapt to their environment and ensure appropriate responses. Tissue-specific transcriptional programmes, instigated by local signals, enable phenotypic specialization in discrete microenvironmental niches controlled by differential transcription factor usage.^{21,22} Dysfunction of MΦ behaviour or phenotype has been associated with the development of many conditions such as neurodegeneration, arthritis, chronic inflammation, atherosclerosis and cancer,^{23,24} and the identification of distinct subpopulations of MΦ may be key in disease understanding and treatment.²⁵⁻²⁷

All MΦ rely on specific cytokine availability for survival, proliferation and phenotype, including macrophage colony-stimulating factor (M-CSF), granulocyte-macrophage CSF (GM-CSF), IL-34 and transforming growth factor beta 1 (TGF-β1), as single factor or in combination.^{6,28-30} MΦ programming strongly depends on the environment, which profoundly affects the cells at a transcriptomic and epigenetic level.^{21,22,31} MΦ also demonstrate remarkable immune memory or 'trained immunity' capacities³²⁻³⁵ based on epigenetic modification following stimulation and long-term priming or 'imprinting', as seen with apoptotic cell recognition leading to a relatively stable tolerant state.³⁶⁻³⁸ This immune memory appears critical during

transplant, when monocytes and MΦ acquire specific memory to major histocompatibility class I (MHC I).³⁹ However, once isolated for *ex vivo* study, the characteristic gene signature of tissue-resident MΦ, including epigenetic modification,^{22,40,41} is often lost, indicating the need to explore their function and reprogramme them preferentially *in vivo*, or with the *in vivo* context considered.

The aim of this review was to have a broad overview of current research and possible treatments directed at MΦ in a selection of disease contexts. We have confined the focus to specific organs and conditions detailed below, purely as exemplars of the kinds of approaches that are being considered. However, the potential for MΦ targeting should not be considered restricted to the provided examples.

Methods to reprogramme macrophages

MΦ exhibit a high degree of plasticity in response to environmental signals, many of which are tissue- and context-specific. This results in a variety of MΦ subtypes with different origins, which may play distinct roles in human disease and potentially provide unique opportunities for targeted therapies.^{26,27,42} Before considering detailed examples of therapeutic approaches in specific tissue and disease contexts, we briefly introduce some common methods used to reprogramme MΦ, from conventional approaches, such as targeted antibody treatments and small molecule drugs to cutting-edge technology of gene expression modification using viral vectors, artificial DNA carriers, naked DNA and cell therapy (Figure 1).

Targeted antibody treatments are among the easiest and most efficient methods to target not only MΦ surface receptors involved in the regulation of immune responses⁴³ but also circulating cytokines/growth factors, preventing their interaction.⁴⁴ As a result, antibodies can alter MΦ activation status. However, this technique is mostly systemic and can lead to numerous off-target effects.

Gene therapy aims to alter specific gene expression by inserting genetic material into the target cell. Free nucleic acids can be directly injected *in vivo*. While this is generally considered safe,⁴⁵ detection by MΦ can induce inflammatory signalling. Although possibly advantageous when reprogramming MΦ into pro-inflammatory phenotypes, it may counter anti-inflammatory states and raises concern of off-target effects. Free nucleic acids also lack cell-targeting specificity, an issue that can be solved by attachment to carrier molecules, such as coupling to peptides directly targeting MΦ cell surface receptors.⁴⁶

Nucleic acids can also be introduced using modified viral vectors that lack the genes necessary for replication. Lentiviral vectors stably integrate genetic material into the host cell genome, while adenoviruses and adeno-associated viruses (AAVs) only cause transient gene expression. Despite their high efficacy, viral vectors are associated




Figure 1. Summary of macrophage manipulation techniques for therapeutic purpose. These strategies can directly be applied *in vivo*, as well as *in vitro* followed by adaptive transfer of manipulated MΦ. Free nucleic acids (1) can be manufactured easily and are very successfully used in some tissues including lungs and skeletal muscle. However, they lack MΦ specificity and are rapidly cleared from the environment, mostly by circulating enzymes and kidney. Viral vectors (2) can be employed to deliver nucleic acids, preventing clearance from the system. Depending on the type of vector, gene manipulation can be long term (lentivirus) or transient (adenovirus). Viral vectors are highly efficient and can be modified to improve MΦ targeting. However, they do entail safety considerations for patients and manufacturing staff. Free small molecules and cytokines (3) are known to act on MΦ polarization. They are easy to administer but prone to degradation. They are also often not MΦ specific and can cause off-target effects and toxicity. Encapsulation of nucleic acids, small molecules and cytokines into nanovectors (4) prolongs their half-life in the organism, while surface modifications (4b) allow targeting of specific cell types. Antibodies (5) can manipulate MΦ polarization by directly binding Fc or other cell surface receptors. While they are generally safe, high doses are often required for therapeutic efficacy translating into high costs

with significant disadvantages: from a manufacturing perspective, viral vector production is costly and requires specific safety measurements.⁴⁷ From a clinical point of view, random lentiviral RNA insertion into the genome could cause tumour suppressor gene disruption triggering

malignancy.⁴⁸ Also, viral vectors bear the risk of potentially high immunogenicity.^{49,50} Adenoviruses' triggering of immune responses could, however, be exploited for the use in tumour contexts where immune activation could be beneficial.

Another option for delivery of nucleic acids are non-immunogenic nanoparticles (<100 nm), and organic (e.g. liposome, polymers) or inorganic (e.g. gold, silica) particles widely used in clinical applications⁵¹ that are readily ingested by M Φ . Different types of nanovectors vary in their advantages in clinical applications⁵² and can inherently favour polarization towards either end of the M Φ activation spectrum.⁵³

The use of nanoparticles is not restricted to nucleic acid delivery. M Φ can also be targeted with compounds acting on signalling pathways to promote polarization such as cyclooxygenase-2 inhibitors,⁵⁴ receptor tyrosine kinase inhibitors⁵⁵ and histone deacetylase (HDAC) inhibitors,⁵⁶ as well as small molecule Toll-like receptor (TLR) agonists⁵⁷ and possibly cytokines.⁵⁸ Free small molecules can be associated with systemic side effects, while unprotected molecules and cytokines are relatively unstable *in vivo*.⁵⁹ Such limitations can be overcome by encapsulation into nanovectors.^{60,61} In general, nanoparticles are well tolerated, small enough to cross physiological barriers including the blood–brain barrier (BBB) and easily modified to allow cell-targeted delivery. They do, however, need to be carefully manufactured considering possible toxicity associated with different materials and delivery routes, as well as inflammatory responses associated with uptake by M Φ .⁶²

Blood and bone marrow-derived M Φ (BMDM) can be reprogrammed *ex vivo* by the same aforementioned methods and adoptively transferred to individuals. This may help alleviate some of the off-target limitations of directly targeting M Φ *in vivo*.⁶³ To retain polarization stability, M Φ can be genetically engineered *ex vivo* to over- or under-express factors associated with polarization phenotypes. An intriguing option was proposed by Shields and colleagues: *ex vivo* attachment of IFN- γ -loaded phagocytosis-resistant ‘backpacks’ to BMDM enabled slow release of IFN- γ *in vivo*, allowing injected cells to maintain a pro-inflammatory phenotype while simultaneously polarizing tumour-infiltrating M Φ .⁶⁴

The type of protocol used for therapeutic intervention will depend on the tissue, M Φ subtype and pathology to be treated. Recently, a first-in-human phase 1 dose–escalation trial confirmed the safety of autologous M Φ therapy in end-stage liver disease, which was well tolerated and is currently undergoing efficacy measures in an ongoing phase 2 randomized controlled trial.⁶⁵

Tissue-specific consideration in macrophage reprogramming

Cardiac macrophages

Mouse cardiac M Φ are composed of four different subsets, distinguishable by their cell surface expression of typical markers such as MHCII, Ly6C, CCR2, CD11c,

MerTK, CD206 and CD64, and are derived from yolk sac and fetal liver.⁶⁶ Their self-renewal capacity decreases with age, and they are thought to be gradually replenished by monocyte-derived M Φ over time.⁶⁷ Cardiac M Φ play important roles in tissue homeostasis, angiogenesis and vascular remodelling during embryogenesis, as well as in the action potential propagation by being electronically coupled with cardiomyocytes.^{68–70} During inflammation or following injury such as myocardial infarction (MI), resident cardiac M Φ become activated and high numbers of monocyte-derived M Φ are recruited to the injured site where their heterogeneity is thought to impact MI outcome.^{71–73} Recruited cardiac M Φ participate in all phases of MI, from the first acute inflammatory wave to the reparative phase and the promotion of angiogenesis and muscle regeneration.^{74,75} The repair after MI is associated with scar formation and fibrosis, decreasing cardiac functionality.⁷⁶ Interestingly, the resident cardiac M Φ seem to limit adverse remodelling⁷⁷ and Gata6⁺ pericardial M Φ have recently been shown to enter the site of injury and prevent fibrosis.⁷⁸ MI is a leading cause of death worldwide.⁷⁹ Clinical trials mainly focus on decreasing systemic inflammation by suppressing the immune system and inflammation with corticosteroid,⁸⁰ methotrexate (phase 3, completed and recruiting) or blocking pro-inflammatory molecules such as IL-6 (tocilizumab, phase 2, active) and IL-1 β (Anakinra, completed⁸¹). However, the use of broad-range immunosuppressors is questionable, as a controlled first phase of inflammation has been shown to be essential to tissue regeneration.⁸² In this line, recent work has shown that a located injury such as MI not only modifies the local M Φ pool, but also affects the number and molecular signature of off-site M Φ throughout the organism (liver, lung and kidney),⁸³ placing the effect of systemic drugs in an even more central question. Recent research focusing on the modulation of cardiac M Φ phenotype and activity in animal models shows promising results encouraging further human clinical trials. Using nanoparticle-delivered siRNA, Courties et al. successfully silenced interferon regulatory factor 5 (IRF5) in cardiac M Φ in a mouse model of MI, which improved healing.⁸⁴ Similarly, the targeting of miRNA-21 in cardiac M Φ with nanoparticles containing mimics (intravenously injected) or adenovirus particles overexpressing miR-21 (intramyocardially injected) reduced inflammation, fibrosis and cardiac dysfunction.^{85,86} Finally, the intramyocardial transplantation of *in vitro* M-CSF- and IL-4-primed ‘reparative’ M Φ following MI in mice showed beneficial effects over transplantation of non-primed bone marrow mononuclear cells.⁸⁷ While human assays to modify M Φ *in vivo* has been less prominent, recent work explored the M Φ heterogeneity and identified at least two subsets (CCR2⁺ and CCR2⁻) present in human heart which are essential for tissue function.⁸⁸

Macrophages in the central nervous system

Resident myeloid cells in the central nervous system (CNS) include parenchymal microglia and different types of non-parenchymal M Φ .⁸⁹ Additionally, inflammatory conditions can trigger the influx of monocyte-derived M Φ ⁹⁰ through disruption of the BBB. Under physiological conditions, this layer of endothelial cells restricts the crossing of cells and compounds into the CNS as previously reviewed.⁹¹ Both dysregulated resident and infiltrating M Φ have been linked to the pathophysiology of several neurological disorders including Alzheimer's disease (AD),^{92,93} multiple sclerosis (MS)⁹⁴ and amyotrophic lateral sclerosis (ALS).⁹⁵ Currently, no cure exists for those conditions and potential preventative and early treatment options face the difficulty of drug delivery.

Multiple sclerosis

MS is an autoimmune, demyelinating disease of the brain and spinal cord. It affects young adults causing progressive neurological deterioration with common symptoms including numbness, burning sensations, visual impairment, loss of balance, bladder dysfunction, fatigue and depression.⁹⁶ While the aetiology of MS is incompletely understood, it is believed that CNS-infiltrating pro-inflammatory phagocytes are key drivers in tissue destruction.⁹⁴ Indeed, in MS patients such cells have been found to express reduced levels of SHP-1 causing increased activation of STAT1, STAT6 and NF- κ B.⁹⁷ This was associated with a pro-inflammatory phenotype characterized by elevated proteinases including ADAM8, which could disrupt the BBB and contribute to demyelination, as well as increased molecules associated with antigen presentation and costimulation.⁹⁸ This view is supported by studies of experimental autoimmune encephalomyelitis (EAE), the murine model for MS.⁹⁹ Depletion of infiltrating phagocytes protected mice from axonal damage,¹⁰⁰ while blocking microglial release of nitrite and pro-inflammatory chemokines and cytokines significantly reduced clinical signs in the EAE model.¹⁰¹ The polarization of microglia by molecular control of cytokines and costimulatory molecules by nuclear receptor has been shown to be essential for the onset and development of EAE.¹⁰² EAE has limitations concerning translation to human disease but does still play an important role in drug development.¹⁰³

Glatiramer acetate (GA) is a polymer approved for the treatment of relapsing–remitting MS. While its mode of action is attributed to T-cell manipulation, GA also increases microglial phagocytic activity and IL-10 production while decreasing TNF *in vitro*.¹⁰⁴ GA likely does not cross the BBB on its own. However, it can be taken up by dendritic cells (DCs) and released in the CNS, with GA uptake promoting trans-endothelial migration of

DCs.¹⁰⁵ Alternatively, GA could enter the CNS when the BBB is disrupted in active MS lesions. Studies report GA-mediated reductions in relapse rates;¹⁰⁶ however, it is still unclear whether its effect on microglia contributes to this and whether GA actually slows disease progression.¹⁰⁷ As GM-CSF has been implicated in disease induction in the EAE model, MOR103, a monoclonal antibody (mAb) directed against GM-CSF, has been tested in MS patients and was found to be well tolerated, although no assessment regarding efficacy was conducted at this stage (NCT01517282).

While infiltrating pro-inflammatory phagocytes are potentially detrimental in MS, microglial production of pro-inflammatory factors, especially TNF, might support remyelination,¹⁰⁸ calling for a specific targeting of peripheral M Φ . Indeed, preventing peripheral M Φ entry into the CNS was beneficial in EAE.¹⁰⁹

Therapeutics aiming to interfere with peripheral M Φ need to be carefully considered to not affect microglia populations. In EAE, scRNAseq recently identified eight different monocyte subsets, one of which expressing CXCL10⁺ was associated with a pathogenic gene signature. This subset had also been observed in other inflammatory models such as pathogen infection, and its depletion in EAE was associated with clinical improvement.¹¹⁰ Whether and how this finding translates into the human context remains to be investigated.

Alzheimer's disease

AD is a chronic neurodegenerative disorder associated with CNS inflammation and the main cause of dementia. Genome-wide association studies have uncovered a variety of loci increasing susceptibility to the development of late-onset Alzheimer's disease, many of which are associated with immunity.⁹³ In addition, rare coding variants, such as that of triggering receptor expressed on myeloid cells 2 (TREM2), have strongly been linked to increased risk of developing AD.¹¹¹ In line with this, in a murine AD model expressing a TREM2 risk variant (R47H), anti-human TREM2 mAb increased microglial proliferation, reduced neuroinflammation and was associated with cognitive benefits.¹¹² The same antibody was well tolerated in a phase I clinical trial (NCT03635047).

Activated microglia are thought protective early in disease but over time become dysfunctional causing inflammatory injury.¹¹³ Importantly, scRNAseq identified distinct microglia subsets in AD, including a type associated with neuroprotection and increased phagocytosis also found in a mouse model of ALS.²⁵

An important regulator in microglial differentiation is the M-CSF receptor (M-CSFR): depending on the murine model and dose of inhibitor used, inhibition of M-CSFR signalling resulted in blockade of microglial proliferation^{114,115} or microglial depletion¹¹⁶ inducing an anti-

inflammatory state and cognitive improvements not associated with alterations in amyloid plaques. Together, these and other studies suggest M-CSFR inhibition as a viable option for clinical trials. Importantly, M-CSFR expression is not restricted to microglia but important for all macrophages. In line with this, M-CSFR inhibition was shown to also affect circulating and tissue-resident macrophages and lymphocytes in other organs in a mouse model.¹¹⁷ In this paper, the M-CSFR inhibitor PLX5622 was administered through diet. While not investigated by the authors, it is likely that such a method of drug delivery would also affect the gut microbiome. Considering the link between intestinal dysbiosis and neurodegeneration,¹¹⁸ such a method could affect treatment outcomes and a more localized way of drug delivery, potentially by intranasal¹¹⁹ or direct intraventricular/intrathecal administration,¹²⁰ could minimize off-target effects.

Amyotrophic lateral sclerosis

ALS is characterized by motoneuron degeneration typically causing paralysis and death within five years of diagnosis. In murine models, the number of resident microglia increases throughout disease progression concomitant with a switch from an anti- to a pro-inflammatory phenotype.¹²¹ Simultaneously, peripheral monocytes from ALS patients have been suggested to be more readily activated into pro-inflammatory phenotypes compared with those of healthy individuals, further driving neuroinflammation after CNS infiltration.¹²²

Vascular endothelial growth factor (VEGF) has been suggested to play a neuroprotective role by reducing motor neuron death through downregulation of pro-inflammatory cytokine production.¹²³ Intrathecal injection of adeno-associated virus containing VEGF expressing plasmid induced VEGF expression in motor neurons in a mouse model of ALS. This prolonged survival of the mice by two modes of action: increasing anti-apoptotic factors such as Bcl-2 and decreasing pro-apoptotic ones such as Bax, caspase-3 and caspase-9 in neurons and a switch in the inflammation balance (reduction in the pro-inflammatory TNF, IL-1 β and CD68 while increase in anti-inflammatory TGF- β released by microglia).¹²⁴ Unfortunately, phase II clinical trials for intracerebroventricular administration of the common splicing isoform VEGF165 were terminated due to a lack of favourable benefit–risk ratio (NCT01384162).

The involvement of pro-inflammatory M Φ and microglia in the pathophysiology of neuroinflammatory conditions strongly suggests the targeting of CNS myeloid cells for re-education as a treatment option. However, to date no agent designed to manipulate M Φ activation states in the CNS has been approved for the treatment of a neurological condition. This is partly due to poor animal models, which do not fully mimic human pathophysiology. It

is essential to understand the heterogeneity of myeloid populations and individual contributions to neurological pathologies in the CNS, to specifically target disease-associated cell types. Imaging mass cytometry is already used to characterize key players in neurological diseases by applying antibodies to post-mortem sample.¹²⁵ Additionally, scRNAseq of human microglia from brain autopsy samples could also uncover cell-specific targets for intervention that could aid the development of therapeutics. With potential targets being discovered, it is vital to also reconsider options for drug delivery: compounds delivered systemically can eventually reach the brain but will also affect cells in other tissues and organs. In particular, in the case of macrophage manipulation, many receptors and targets are shared by different tissue-resident macrophage populations making it difficult to prevent off-target effects. This issue could be overcome by exploring intracerebral,¹²⁶ interstitial¹²⁷ or intranasal¹²⁸ delivery methods.

Liver macrophages

The murine liver comprises two distinct populations of tissue-resident M Φ , the Kupffer cells (KCs) that occupy the sinusoidal vascular space, and the phenotypically distinct liver capsular M Φ , which reside in the hepatic capsule.¹²⁹ Additionally, the liver may also contain monocyte-derived and peritoneal M Φ ,⁸⁶ which are recruited following inflammatory events or injury.^{130–132} Liver M Φ heterogeneity was recently reviewed.^{130,131} In mice, fetal liver monocytic precursors give rise to embryonic KCs, which at steady-state maintain the KC pool through self-renewal, independent of BM-derived progenitors.^{1,133,134} In contrast, liver capsular M Φ arise entirely from adult circulating monocytes.¹²⁹ More recently however, BM-derived monocytes were shown to populate the KC niche during postnatal liver development, contributing significantly to the adult KC population.¹³⁵ Following the loss of KCs after infection,¹³⁶ or experimental depletion,^{135,137,138} the KC pool is repopulated through proliferation of surviving KCs, as well as the recruitment and differentiation of blood monocytes into KCs (mo-KCs). These mo-KCs were shown to be highly functionally and transcriptionally homologous to their embryonic counterparts¹³⁵ after 30 days post-depletion, although other studies have shown that embryonically derived KCs may exhibit some phenotypic and functional differences to their monocyte-derived counterparts.^{137,139} The replenishment of the depleted KC niche by either mechanism however, appears to be context-dependent. Indeed, in an acute liver injury model, the depleted KC niche was replenished through proliferation of the surviving KCs without input from circulating monocytes.¹⁴⁰ Similar to the development and maintenance of other tissue-resident M Φ populations, recent studies have highlighted the critical role of transcription factors in governing KC

development and identity, with the loss of inhibitor of DNA 3 (ID3)¹⁴¹ or liver X receptor- α (LXR- α)¹⁴² resulting in KC deficiency in mice. Two recent studies in mice demonstrated that liver-derived signals orchestrate monocyte recruitment¹⁴³ and initiate and maintain KC identity through the induction of lineage-determining factors including ID3 and LXR- α by acting on pre-existing but poised enhancers^{143,144}. Additionally, leveraging of scRNAseq technology for profiling human liver M Φ has revealed distinct subpopulations of KCs with discrete gene expression signatures^{26,145-147}, but our understanding of their biology is still restricted compared with mouse liver M Φ .

Liver M Φ play a key role in the pathogenesis of acute and chronic liver diseases, including acute liver failure, alcoholic liver disease, non-alcoholic fatty liver disease (NAFLD), viral hepatitis and hepatocellular carcinoma. They are also critically required for the restoration of tissue homeostasis and resolution of liver disease.¹⁴⁸ These seemingly contrasting roles for liver M Φ highlight the importance to identify and understand the relative contributions that distinct subsets have in disease progression, as well as tissue repair, to help the development of improved-targeted therapeutics, as well as the definition of biomarkers indicating either disease progression or regression (for a recent review, see Ref 131).

A recent study described a subset of Trem2^{hi} M Φ enriched in mouse models of non-alcoholic steatohepatitis (NASH), as well as in human NASH livers, correlating with disease severity.¹⁴⁹ Interestingly, in a diet-induced mouse model of NASH, KC identity was significantly altered by the NASH diet. NASH-induced changes in KC enhancers and gene expression were driven by activator protein 1 and early growth response protein 1 inducing a scar-associated M Φ phenotype, with increased expression of both *Trem2* and *Cd9*.¹⁵⁰ Interestingly, an independent study also identified a scar-associated TREM2⁺CD9⁺ subpopulation of M Φ in humans, which differentiate from circulating monocytes and expand during liver fibrosis.²⁶ In another study, a subset of M Φ (MerTK⁺HLA-DR^{high}) was reported to expand during the resolution phase of acute liver disease, with a comparable population identified in mice during the resolution phase of an acute liver injury model.¹⁵¹ Further studies are needed to fully define liver M Φ subsets and their contributions to disease progression, as well as in tissue repair and the restoration of homeostasis. However, some approaches to target M Φ in liver disorders are already under investigation.

Glucocorticoid and antibody–drug conjugate

Glucocorticoid receptor signalling modulates inflammation in KCs, suggesting that glucocorticoid treatment could serve as a potential M Φ -directed treatment for liver diseases.¹⁵² Liposomal delivery of dexamethasone was

shown to significantly reduce liver injury and fibrosis in experimental models of both acute and chronic liver injuries.¹⁵³ Direct targeting of M Φ with an antibody–drug conjugate composed of dexamethasone linked to an antibody against CD163 (a scavenger receptor highly expressed in KCs and infiltrating monocytes/M Φ) was shown to reduce inflammation, hepatocyte ballooning and fibrosis in a mouse model of NASH, while having no apparent systemic side effects.¹⁵⁴

Cytokines modulation

The pro-inflammatory cytokine TNF plays a major part in the development of steatosis, inflammation and fibrosis in NAFLD. KCs have been identified as the main cellular source of TNF in mouse models of NAFLD. Mannose-modified trimethyl chitosan–cysteine-conjugated nanoparticles were used to deliver siRNA targeting TNF to M Φ and protected mice from inflammation-driven liver damage and lethality in an acute liver injury model.¹⁵⁵

Gene expression modification

Oxidative stress and the associated damage have been suggested to link obesity and liver disease. Recently, Azzimato and colleagues¹⁵⁶ showed that oxidative stress was triggered by obesity in mouse and human livers. In parallel, nuclear factor erythroid 2-related factor 2 (NRF2), a transcription factor regulating the antioxidant response, was reduced, leading to an impaired antioxidant response. miR-144 was greatly upregulated in the liver of obese and insulin-resistant mice and humans and was shown to target NRF2. Consequently, delivery of an antagomiR targeting miR-144 expression to M Φ *in vivo* with glucan-encapsulated RNA interference particle technology increased NRF2 protein levels, reduced oxidative stress and improved hepatic metabolism in insulin-resistant mice.

Small molecule inhibitors

Galectin-3 is a pleiotropic protein highly expressed by M Φ in the liver and upregulated in models of liver disease.^{157,158} Galectin-3 deficiency in mice was protective in a concanavalin A-induced liver injury model, reducing pro-inflammatory cytokines, as well as attenuating fibrosis.¹⁵⁹ Similarly, pharmacological inhibition of Galectin-3 in liver injury models significantly reduced fibrosis and led to a reversal in cirrhosis and is now under evaluation for NASH in clinical trials.^{157,158}

The dual CCR2/CCR5 inhibitor cenicriviroc was shown to reduce monocyte recruitment and liver injury in an acute liver failure model,¹⁶⁰ as well as ameliorating hepatic inflammation and fibrosis in experimental models of NASH,¹⁶¹ and is also currently under evaluation in

clinical trials for the treatment of liver fibrosis in NASH (NCT03059446 and NCT03028740).¹⁶²

Adoptive transfer

In an experimental model of acute liver injury, adoptive transfer of *ex vivo* IL-4/IL-13-polarized BMDM rapidly reduced liver injury and several mediators of inflammation.¹⁶³ Of note, the adoptive transfer of primary human-polarized monocyte-derived M Φ partially recapitulated the therapeutic effect observed with polarized mouse BMDM in the same mouse model.¹⁶³ Similarly, injection of BMDM or embryonic stem cell-derived M Φ reduced both fibrosis and improved liver regeneration in a hepatic injury and fibrosis model.^{164,165} Clinical trials in humans have demonstrated the safety of administration of large and frequent infusions of autologous M Φ . A recent first-in-human trial evaluated the safety of a single peripheral infusion of autologous M Φ in end-stage liver disease, which was well tolerated and led to a reduction in clinical scoring, and is currently undergoing efficacy measures in an ongoing phase 2 randomized controlled trial.⁶⁵ However, this study did not determine whether the infused M Φ migrated to and engrafted in the liver. Previous studies in mice, as well as a case study in humans, suggest that the administration via peripheral or central veins M Φ traffic from the pulmonary vasculature via the blood before engrafting in the liver and spleen.^{164,166-168} Significant challenges remain for the adoption of autologous M Φ therapies, such as their scalability, as well as ensuring that engrafted cells maintain the intended phenotype, which can be greatly impacted by the tissue microenvironment.

Macrophages in arthritis

Rheumatoid arthritis (RA) is a chronic, systemic inflammatory autoimmune disorder that primarily affects synovial joints, leading to irreversible bone and cartilage destruction. Multiple studies have implicated monocytes and M Φ in the initiation and progression of RA.¹⁶⁹ M Φ are the most abundant immune cell and are a source of pro-inflammatory cytokines associated with RA pathogenesis including TNF, IL-6 and IL-1 β ,¹⁷⁰ as well as chemoattractants and metalloproteinases. Conversely, synovial tissue M Φ in healthy and RA patients in sustained remission suggest that M Φ have a fundamental role in maintaining and/or reinstating synovial homeostasis.^{171,172} In a murine model of sterile inflammatory arthritis, non-classical Ly6C⁻ but not Ly6C⁺ monocytes were reported to be crucial for the initiation of arthritis, while tissue-resident synovial M Φ restricted the development of arthritis.¹⁷³ Murine synovial lining CX3CR1⁺ M Φ form an immunological barrier in the lining layer of the synovium of healthy joints.¹⁷⁴ Depleting them in a mouse model of

arthritis disrupted barrier function and accelerated the onset and magnitude of arthritis, whereas depletion of CSF1R⁺ monocytes and M Φ expedited the resolution of inflammation. Interestingly, comparison of scRNAseq data with human data sets from RA patients revealed significant overlap, suggesting cells similar to synovial lining M Φ may also exist in humans.¹⁷⁴ A recent study described synovial M Φ subsets enriched in active RA (MerTK⁻ CD206⁻) or sustained remission (MerTK⁺ CD206⁺), which are thought to contribute to RA pathogenesis or remission through the production of pro-inflammatory cytokines or lipid mediators, respectively.¹⁷⁵ Similarly, *IL1B*⁺ pro-inflammatory monocytes were enriched in synovial tissue from patients with RA, whereas *NUPRI*⁺ monocytes were inversely correlated with tissue inflammation.¹⁷⁶ Another study recently identified HBEGF⁺ inflammatory M Φ enriched in the human RA tissues,¹⁷⁷ which promote synovial fibroblast invasiveness in an epidermal growth factor receptor (EGFR)-dependent manner. Interestingly, a previous study showed that EGFR inhibition reduced the severity of established RA in mice.¹⁷⁸ Collectively, these studies demonstrate that distinct monocyte and M Φ subsets have defined roles in RA. Strategies are currently being developed to target M Φ and improve RA pathology.

Cytokines

Several lines of evidence suggest that M Φ and GM-CSF strongly influence the development and progression of RA.^{179,180} In patients with RA, GM-CSF is elevated in plasma, synovial fluid and synovial tissues, while administration of recombinant GM-CSF has been reported to exacerbate RA disease activity.¹⁸¹ Depletion of GM-CSF or blockade of GM-CSFR α in a mouse model of RA significantly reduced the number of M Φ in the inflamed synovium, decreased synovial inflammation and joint destruction.^{182,183} Therapeutic antibodies targeting GM-CSF and its receptor have been developed and evaluated in clinical trials.^{181,184} Collectively, these studies have shown that treatment is associated with rapid and sustained improvements in measures of RA disease outcomes, as well as being well tolerated in safety studies.¹⁸⁵ Of note, a phase III clinical trial is currently ongoing evaluating otilimab, a fully human anti-GM-CSF monoclonal antibody, in RA patients who have had an inadequate response to disease-modifying anti-rheumatic drugs and/or Janus kinase inhibitors (NCT04134728).

Gene therapy

In the collagen-induced arthritis (CIA) mouse model, Kong and colleagues identified three key pro-resolving factors that were elevated in mouse synovial tissues during the resolution phase of CIA.¹⁸⁶ Among them, tyrosine

3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (Ywhaz) was also shown to be elevated in rat synovial tissue in a CIA model during the resolution phase compared with the peak phase, as well as in RA patients who responded well to treatment with anti-rheumatic drugs. *Ywhaz* is highly expressed in Tregs and M Φ and is one of the most common proteins found in exosomes¹⁸⁷ from Tregs, although no data were shown for M Φ in this study. Interestingly, treatment of thioglycollate-elicited peritoneal M Φ with recombinant *Ywhaz* reduced the expression of *Tnf* and *Il6*, while enhancing *Il10* expression.¹⁸⁶ Furthermore, siRNA knockdown of *Ywhaz* or treatment with an anti-*Ywhaz* antibody enhanced the expression of *Tnf* and *Il6* following LPS stimulation, supporting a role of *Ywhaz* in modulating pro-inflammatory cytokine production in M Φ .¹⁸⁶ Intra-articular delivery of adenovirus expressing *Ywhaz* suppressed the production of pro-inflammatory cytokines and significantly reduced synovial inflammation and joint destruction in mice.¹⁸⁶ *Ywhaz* is a member of the 14-3-3 protein family, which modulate the activity of binding partners by controlling protein localization, stability, conformation and activity mainly through phosphoserine/threonine motifs.¹⁸⁸ Previous studies have suggested that *Ywhaz* can modulate the activity of forkhead box transcription factor family O 3 (FOXO3), a transcription factor regulating the expression of cytokines,¹⁸⁹ and tristetraprolin (TTP),¹⁹⁰ which regulates cytokine production by destabilizing target mRNA molecules.¹⁹¹ However, further studies are needed to understand how *Ywhaz* regulates the production of cytokines and the subsequent resolution of arthritis. In arthritic rats, M Φ phenotype has been manipulated *in vivo* by tuftsin-modified nanoparticle-mediated delivery of plasmid encoding IL-10, leading to a significant reduction in pro-inflammatory cytokine production, diminishing inflammation and preventing the progression of joint damage.¹⁹²

Tumour-associated macrophages

Cancers are heterogeneous tissues with tumour-infiltrating immune cells playing key roles in disease progression.¹⁹³ Tumour-associated macrophages (TAMs) are major components of the immune cells infiltrating solid tumours. They are a heterogeneous population often coexpressing pro- and anti-inflammatory markers.¹⁹⁴ While pro-inflammatory functions could aid tumour cell elimination,¹⁹⁵ immune-suppressive TAMs are linked to worse prognosis due to their contribution to tumour growth and metastasis.^{196,197} TAMs repolarization to produce inflammatory mediators is currently being evaluated in a number of clinical trials, examples of which are summarized in Table 1.¹⁹⁸⁻²⁰⁶ Several strategies to target TAMs have been adopted, and we will describe common ones below.

Receptor targeting

Just as in other tissues, the M-CSFR axis is an attractive target in the tumour context: in many pre-clinical models, the use of M-CSFR inhibitors is associated with M Φ depletion and tumour regression in a T-cell-dependent manner.²⁰⁷ However, blocking M-CSFR signalling in murine proneural glioblastoma multiforme (PGM) and hepatocellular carcinoma caused tumour regression without TAM depletion, with persisting TAMs associated with functional alterations.^{208,209} M-CSFR signalling can be blocked by the use of antibodies targeting either M-CSF or its receptor. While in some clinical trials antibody treatment achieved disease stabilization,²¹⁰ other studies report TAM reductions without anti-tumour activity.²¹¹

Another option is the use of small molecule inhibitors acting on the receptor tyrosine kinase domain (e.g. PLX3397, BLZ945). Those inhibitors also interfere with other receptors expressed in myeloid and tumour cells, which could enhance their efficacy. As it has been shown in a PGM model that tumours can acquire resistance to BLZ945-mediated inhibition,²¹² the combination of M-CSFR inhibitors with other immune- or chemotherapeutic agents could be beneficial and is being evaluated in clinical trials.

TLRs are involved in immune surveillance, and their agonists can drive pro-inflammatory mediator release.²¹³ Imiquimod (TLR7 agonist) is already approved for topical therapy in squamous and basal cell carcinoma, and clinical trials are ongoing for several other TLR agonists. Imiquimod's mechanism of action likely affects not only M Φ but also dendritic cells and neutrophils.²¹⁴ *in vivo*, many TLR agonists have short half-lives, and some free agonists have been associated with toxicity. Such issues could be prevented by encapsulation into nanovectors. While no such strategies are currently in clinical trials, murine models support the efficacy and safety of resiquimod-loaded nanoparticles⁶¹ and ferumoxylol-linked Poly (I:C).²¹⁵

Another example of receptor targeting exploits CD40, a costimulatory molecule expressed on the surface of myeloid cells and B cells, with key roles in immune regulation.²¹⁶ Ligation with an agonist CD40 mAb triggered T-cell-dependent anti-tumour immunity in murine models of pancreatic ductal adenocarcinoma (PDAC) associated with increased MHCII, CD80 and CD86 expression on M Φ and elevated serum levels of IL-12 and TNF.²⁰⁴ Depletion of M Φ in this context prevented tumour regression. In a clinical trial, CD40 mAb combined with chemotherapy showed a trend of increased overall survival, even though the sample size was too small ($n = 21$) for conclusive results.²⁰⁴ Several other clinical trials that investigate the effect of targeting CD40 in different types of tumours have already been summarized.²¹⁷ While some of these therapeutics are associated with positive tumour

Table 1. Selective examples of TAMs reprogramming compounds currently undergoing clinical trials updated

Compound	Target	Clinical phase	Clinicaltrials.gov identifier	Status	Results	Type of malignancy
Imiquimod, cyclophosphamide and radiotherapy	TLR7	Phase II	NCT01421017	Complete	No results available	Skin metastasis in breast cancer
Imiquimod together with Abraxane	TLR7	Phase II	NCT00821964	Complete	Pathologic clinical response in 71.4% of patients	Advanced breast cancer ²⁰⁰
Imiquimod	TLR7	Phase II	NCT00031759	Complete	No impact on recurrence of cervical dysplasia	Cervical cancer ²⁰¹
Imiquimod	TLR7	Phase III	NCT00941252	Complete	Histologic regression in 73% of patients	Cervical intraepithelial neoplasia ²⁰²
Imiquimod	TLR7	Phase III	NCT01861535	Active, not recruiting	-	Vulvar intraepithelial neoplasia
Imiquimod	TLR7	Phase III	NCT02394132	Recruiting	-	Complex lentigo maligna
Imiquimod	TLR7	Phase IV	NCT01161888	Complete	No results available	Lentigo malignant of the face
Resiquimod	TLR7/8	Phase I/II	NCT01676831	Complete	Significant improvements of treated lesions in 75% of patients, clearing of all treated lesions in 30%	Cutaneous T-cell lymphoma ²⁰³
852A	TLR7	Phase I	NCT00095160	Complete	No results available	Refractory solid organ tumours
852A	TLR7	Phase II	NCT00319748	Complete	Evidence of immune activation as evaluated by cytokine production	Breast, ovarian, endometrial and cervical cancers
852A	TLR7	Phase II	NCT00189332	Complete	No results available	Metastatic cutaneous melanoma
Imo-2055	TLR9	Phase II	NCT00729053	Complete	Treatment-emergent adverse events observed in > 90% of patients	Renal cell carcinoma
CD40 mAb CP-870,893	CD40	Phase I	NCT02225002	Complete	No results available	Advanced solid tumours
CD40 mAb CP-870,893 and gemcitabine	CD40	Phase I	NCT01456585	Complete	No results available	Pancreatic ductal adenocarcinoma ²⁰⁴
CD40 mAb CP-870,893 and chemotherapy	CD40	Phase I	NCT00711191	Complete	Partial response in 4/21 patients, stable diseases in 11/21 patients	Advanced cancer of the pancreas ²⁰⁵
Vorinostat, gefitinib	HDAC	Phase I	NCT02151721	Unknown	-	EGFR mutant lung cancer
IPI-549 alone and with nivolumab	PI3K γ	Phase I	NCT02637531	Recruiting	-	Advanced solid tumours
IPI-549 with Tecentriq and Abraxane/Avastin	PI3K γ	Phase II	NCT03961698	Recruiting	-	Breast cancer, renal cell carcinoma
IPI-549	PI3K γ	Phase II	NCT03795610	Recruiting	-	Locally advanced HPV + and HPV- head and neck squamous cell carcinoma
IPI-549 with nivolumab	PI3K γ	Phase II	NCT03980041	Recruiting	-	Advanced urothelial carcinoma
BLZ945 monotherapy or combination with PDR001	M-CSFR (+/- PD-1 blockade)	Phase I/II	NCT02829723	Recruiting	-	Advanced solid tumours

Table 1. (Continued)

Compound	Target	Clinical phase	Clinicaltrials.gov identifier	Status	Results	Type of malignancy
PLX3397 + radiation therapy + temozolomide	M-CSFR (+ cKit, Flt3)	Phase Ib/II	NCT01790503	Complete	Stable disease in 24/50 patients, complete response in 2/50, partial response in 5/50 patients	Glioblastoma
PLX3397 and sirolimus	M-CSFR (+ cKit, Flt3)	Phase I/II	NCT02584647	Recruiting	-	Unresectable sarcoma and malignant peripheral nerve sheath tumours
MCS110 with carboplatin and gemcitabine	M-CSF	Phase II	NCT02435680	Complete	No results available	Advanced triple negative breast cancer with high TAMs
MCS110 with PDR001	M-CSFR (+/- PD-1 blockade)	Phase Ib/II	NCT02807844	Complete	Partial response in 1/48, stable disease in 9/48 patients	Advanced malignancies ²⁰⁶
LY3022855	M-CSFR	Phase I	NCT02265536	Complete	Stable disease in 5/22 MBC and 3/7 MCPRC patients	Metastatic breast cancer (MBC) Metastatic castration-resistant prostate cancer (MCRPC) ²⁰⁷

outcomes, M Φ -specific effects of treatment are generally not investigated.

In a novel particle-based strategy, Shields and colleagues injected M Φ equipped with 'backpacks' of biodegradable polymers loaded with IFN- γ into mice bearing 4T1 tumours; they showed that *ex vivo*-polarized M Φ carrying these backpacks were able to not only maintain a pro-inflammatory phenotype in the tumour microenvironment, but also induce the same phenotype in resident TAMs, associated with reduced tumour growth, lung metastasis and improved survival.⁶⁴ As backpacks could also be filled with cytokines promoting anti-inflammatory polarization, such an approach could be useful in other disease contexts.

Intracellular pathway targeting

TAMs can also be targeted with compounds that interfere with internal signalling, including modifications of nucleic acids, cellular kinases or HDAC.

Vorinostat, a small molecule inhibitor of HDAC, is approved for cutaneous T-cell lymphoma due to its growth-inhibiting effect on tumour cells.²¹⁸ Vorinostat also affects TAMs in murine models: encapsulating vorinostat and a chemotherapeutic drug within liposomes that target both non-small-cell lung cancer cells and CD206-expressing M Φ suppressed tumour growth via the upregulation of iNOS, CD86 and TNF while downregulating CD206, arginase and IL-10 in M Φ .²¹⁹

Another internal target for influencing M Φ signalling is PI3K γ , a phosphoinositide 3-kinases subunit mainly

expressed by myeloid cells. In a murine model of PDAC, PI3K γ blockade with the PI3K γ/δ inhibitor TG100-115 reduced M Φ expression of Arginase1, TGF- β and IL-10 and increased IL-12 and IFN- γ .²²⁰ This induced tumour suppression and prevented metastasis. M Φ -specific effects were replicated in a clinical trial of the selective PI3K γ inhibitor IPI-549 as monotherapy and in combination with chemotherapy, although the study is still ongoing and no results regarding survival and tumour regression are available yet.²²¹

In mouse models of ovarian cancer, melanoma and glioblastoma, nanoparticle-mediated delivery of *in vitro*-transcribed mRNAs encoding IRF5 together with its activating kinase IKK β increased pro-inflammatory myeloid cells and caused tumour clearance in some animals.²²² Similar results were found with nanoparticles delivering siRNA targeting growth factors including VEGF and placental growth factor, which reduced M Φ CD206 expression and IL-10 production while increasing IL-12 and IFN- γ in tumour tissues.²²³ Recently, siRNA-loaded nanoparticle mediated silencing of signal transducer and activator of transcription 3 and hypoxia-inducible factor 1 α prevented TAM-mediated angiogenesis and decreased tumour size.²²⁴ *In vitro*, repolarization has also been achieved with nanoparticle delivery of plasmid DNA encoding the IL-12 gene²²⁵ and with overexpression of microRNA-155 in triple negative breast cancer TAMs.²²⁶ Another approach delivers CRISPR-Cas9 gene editing machinery into M Φ : knock out of signal regulatory protein α , which engages CD47 on cancer cells and prevents phagocytosis, significantly increased M Φ targeting of osteosarcoma cells *in vitro*.²²⁷

TAMs re-education into pro-inflammatory phenotypes can be beneficial to overcome tumour-induced immune suppression. Reprogramming strategies can be combined with existing immunotherapies, chemotherapy and radiation, which has been shown to be successful in a variety of clinical trials. Importantly, TAMs targeting should be localized to cancerous areas to prevent systemic inflammation, and only affect pro-tumoral M Φ types. To achieve this, further investigation into TAMs heterogeneity and careful design and delivery of potential therapeutics are required.

Discussion

The recent development of individual cell-based technologies has led to an explosion of knowledge about M Φ ontogeny and diversity. The identification of their subpopulations in homeostatic and disordered contexts is a constantly growing field that highlights a broad phenotypic repertoire consistent with the variety of stimuli they are exposed to. Their impact on disease development is still under intense investigation and seems to be subtype-dependent. While some subsets are thought to directly contribute to disease progression, others have been found to be protective. However, most investigation is performed in animal models and the translation of this to human patients is often less clear.

Past approaches to M Φ reprogramming did not appreciate M Φ heterogeneity and plasticity, and how microenvironmental niche influences the phenotype. Early adoptive transfer studies had limited/no efficacy because of the impact of the *in vivo* microenvironment on M Φ phenotype once transferred.²²⁸ Novel experimental avenues, however, have succeeded in maintaining the phenotype of adoptively transferred cells despite environmental factors.⁶⁴ Other strategies have focused on direct *in vivo* modulation of M Φ phenotypes, also giving promising results. While different approaches to M Φ reprogramming have been described, each is associated with pitfalls and benefits influencing their utility for specific tissues. Where possible, future therapeutic approaches should consider tailoring of strategy towards a specific tissue microenvironment, as well as a specific disease-associated subset of cells, to improve efficacy and minimize off-target effects.

As M Φ reprogramming is a recent approach to therapy, important questions still need to be answered: How stably can different subsets of M Φ be re-educated? Is long-term reprogramming of long-lived resident M Φ safe? As M Φ are in constant interaction with their environment and other immune cells, including other M Φ subtypes, will modifying one particular subtype have secondary unintended impacts on the tissue? An alternative approach may be secondary targeting of M Φ populations via manipulation of their environmental cues

through alteration of the communication between tissue and M Φ . This could be achieved by targeting the cells that support M Φ growth and/or polarization, for example. Either way, it is clear that further growth in understanding of M Φ phenotypic heterogeneity in a given microenvironmental/disease context is required to appreciate the potential of targeting the M Φ in disease and capitalize on the advances that have begun to be made in this area.

Acknowledgments

P.R.T is a Wellcome Trust Investigator Award (107964/Z/15/Z) and an MRC UK Dementia Research Institute Professor. R.J.P was an employee of GlaxoSmithKline at the time of writing. R.J.P is currently an employee of the University of Cambridge.

Competing interest statement

The authors declare no conflict of interest.

Data Availability Statement

Not applicable.

References

- 1 Hashimoto D, Chow A, Noizat C, Teo P, Beasley MB, Leboeuf M, et al. Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity* 2013;**38**:792–804.
- 2 Davies LC, Jenkins SJ, Allen JE, Taylor PR. Tissue-resident macrophages. *Nat Immunol.* 2013;**14**:986–95.
- 3 Davies LC, Taylor PR. Tissue-resident macrophages: then and now. *Immunology* 2015;**144**:541–8.
- 4 Ginhoux F, Guilliams M. Tissue-resident macrophage ontogeny and homeostasis. *Immunity* 2016;**44**:439–49.
- 5 Gordon S, Martinez-Pomares L. Physiological roles of macrophages. *Pflugers Arch.* 2017;**469**:365–74.
- 6 Kierdorf K, Prinz M, Geissmann F, Gomez PE. Development and function of tissue resident macrophages in mice. *Semin Immunol.* 2015;**27**:369–78.
- 7 Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma *in vivo*. *Science* 2005;**308**(5726):1314–8.
- 8 Michaelson MD, Bieri PL, Mehler MF, Xu H, Arezzo JC, Pollard JW, et al. CSF-1 deficiency in mice results in abnormal brain development. *Development* 1996;**122**:2661–72.
- 9 DeFalco T, Bhattacharya I, Williams AV, Sams DM, Capel B. Yolk-sac-derived macrophages regulate fetal testis vascularization and morphogenesis. *Proc Natl Acad Sci U S A.* 2014;**111**:E2384–E2393.
- 10 Diez-Roux G, Argilla M, Makarenkova H, Ko K, Lang RA. Macrophages kill capillary cells in G1 phase of the cell cycle during programmed vascular regression. *Development* 1999;**126**:2141–7.
- 11 Pollard JW. Trophic macrophages in development and disease. *Nat Rev Immunol.* 2009;**9**:259–70.
- 12 Mills CD, Kincaid K, Alt JM, Heilman MJ, Hill AM. M-1/M-2 macrophages and the Th1/Th2 paradigm. *J Immunol.* 2000;**164**:6166–73.
- 13 Modolell M, Corraliza IM, Link F, Soler G, Eichmann K. Reciprocal regulation of the nitric oxide synthase/arginase balance in mouse bone marrow-derived macrophages by TH1 and TH2 cytokines. *Eur J Immunol.* 1995;**25**:1101–4.
- 14 Viola A, Munari F, Sanchez-Rodriguez R, Scolaro T, Castegna A. The metabolic signature of macrophage responses. *Front Immunol.* 2019;**10**:1462.
- 15 Porcheray F, Viaud S, Rimaniol AC, Leone C, Samah B, Dereuddre-Bosquet N, et al. Macrophage activation switching: an asset for the resolution of inflammation. *Clin Exp Immunol.* 2005;**142**:481–9.

- 16 Xue J, Schmidt SV, Sander J, Draffehn A, Krebs W, Quester I, et al. Transcriptome-based network analysis reveals a spectrum model of human macrophage activation. *Immunity* 2014;**40**:274–88.
- 17 McArdle S, Mikulski Z, Ley K. Live cell imaging to understand monocyte, macrophage, and dendritic cell function in atherosclerosis. *J Exp Med*. 2016;**213**:1117–31.
- 18 Tay TL, Mai D, Dautzenberg J, Fernandez-Klett F, Lin G, Sagar, et al. A new fate mapping system reveals context-dependent random or clonal expansion of microglia. *Nat Neurosci*. 2017;**20**:793–803.
- 19 Li C, Menoret A, Farragher C, Ouyang Z, Bonin C, Holvoet P, Vella AT, Zhou B. Single-cell transcriptomics-based MacSpectrum reveals macrophage activation signatures in diseases. *JCI Insight*. 2019;**4**:10. <http://doi.org/10.1172/jci.insight.126453>
- 20 Bian Z, Gong Y, Huang T, Lee CZW, Bian L, Bai Z, et al. Deciphering human macrophage development at single-cell resolution. *Nature* 2020;**582**:571–6.
- 21 Lavin Y, Winter D, Blecher-Gonen R, David E, Keren-Shaul H, Merad M, et al. Tissue-resident macrophage enhancer landscapes are shaped by the local microenvironment. *Cell* 2014;**159**:1312–26.
- 22 Gosselin D, Link VM, Romanoski CE, Fonseca GJ, Eichenfield DZ, Spann NJ, et al. Environment drives selection and function of enhancers controlling tissue-specific macrophage identities. *Cell* 2014;**159**:1327–40.
- 23 Wynn TA, Chawla A, Pollard JW. Macrophage biology in development, homeostasis and disease. *Nature* 2013;**496**:445–55.
- 24 Schultze JL, Schmieder A, Goerd S. Macrophage activation in human diseases. *Semin Immunol*. 2015;**27**:249–56.
- 25 Keren-Shaul H, Spinrad A, Weiner A, Matcovitch-Natan O, Dvir-Szternfeld R, Ulland TK, et al. A unique microglia type associated with restricting development of Alzheimer's disease. *Cell* 2017;**169**:1276–90 e17.
- 26 Ramachandran P, Dobie R, Wilson-Kanamori JR, Dora EF, Henderson BEP, Luu NT, et al. Resolving the fibrotic niche of human liver cirrhosis at single-cell level. *Nature* 2019;**575**:512–8.
- 27 Cassetta L, Fragkogianni S, Sims AH, Swierczak A, Forrester LM, Zhang H, et al. Human tumor-associated macrophage and monocyte transcriptional landscapes reveal cancer-specific reprogramming, biomarkers, and therapeutic targets. *Cancer Cell* 2019;**35**:588–602 e10.
- 28 Barreda DR, Hanington PC, Belosevic M. Regulation of myeloid development and function by colony stimulating factors. *Dev Comp Immunol*. 2004;**28**:509–54.
- 29 Hamilton JA. GM-CSF-dependent inflammatory pathways. *Front Immunol*. 2019;**10**:2055.
- 30 Hamilton JA. Colony-stimulating factors in inflammation and autoimmunity. *Nat Rev Immunol*. 2008;**8**:533–44.
- 31 T'Jonck W, Williams M, Bonnardel J. Niche signals and transcription factors involved in tissue-resident macrophage development. *Cell Immunol*. 2018;**330**:43–53.
- 32 Saeed S, Quintin J, Kerstens HH, Rao NA, Aghajani-Refah A, Matarese F, et al. Epigenetic programming of monocyte-to-macrophage differentiation and trained innate immunity. *Science* 2014;**345**:1251086.
- 33 Foster SL, Hargreaves DC, Medzhitov R. Gene-specific control of inflammation by TLR-induced chromatin modifications. *Nature*. 2007;**447**:972–8.
- 34 Netea MG, Dominguez-Andres J, Barreiro LB, Chavakis T, Divangahi M, Fuchs E, et al. Defining trained immunity and its role in health and disease. *Nat Rev Immunol*. 2020;**20**:375–88.
- 35 Wendeln AC, Degenhardt K, Kaurani L, Gertig M, Ulas T, Jain G, et al. Innate immune memory in the brain shapes neurological disease hallmarks. *Nature* 2018;**556**:332–8.
- 36 Fadok VA, Bratton DL, Konowal A, Freed PW, Westcott JY, Henson PM. Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF. *J Clin Invest*. 1998;**101**:890–8.
- 37 Weavers H, Evans IR, Martin P, Wood W. Corpse engulfment generates a molecular memory that primes the macrophage inflammatory response. *Cell* 2016;**165**:1658–71.
- 38 Gordon S, Plueddemann A. Macrophage clearance of apoptotic cells: a critical assessment. *Front Immunol*. 2018;**9**:127.
- 39 Dai H, Lan P, Zhao D, Abou- Daya K, Liu W, Chen W, et al. PIRs mediate innate myeloid cell memory to nonself MHC molecules. *Science* 2020;**368**:1122–7.
- 40 Shemer A, Grozovski J, Tay TL, Tao J, Volaski A, Suss P, et al. Engrafed parenchymal brain macrophages differ from microglia in transcriptome, chromatin landscape and response to challenge. *Nat Commun*. 2018;**9**:5206.
- 41 Bohlen CJ, Bennett FC, Tucker AF, Collins HY, Mulinyawe SB, Barres BA. Diverse requirements for microglial survival, specification, and function revealed by defined-medium cultures. *Neuron* 2017;**94**(4):759–73 e8.
- 42 Mould KJ, Jackson ND, Henson PM, Seibold M, Janssen WJ. Single cell RNA sequencing identifies unique inflammatory airspace macrophage subsets. *JCI Insight*. 2019;**4**:5. <http://dx.doi.org/10.1172/jci.insight.126556>
- 43 Pellizzari G, Hoskin C, Crescioli S, Mele S, Gotovina J, Chiaruttini G, et al. IgE reprograms alternatively-activated human macrophages towards pro-inflammatory anti-tumoural states. *EBioMedicine*. 2019;**43**:67–81.
- 44 Degboe Y, Rauwel B, Baron M, Boyer JF, Ruyssen-Witrand A, Constantin A, et al. Polarization of rheumatoid macrophages by TNF targeting through an IL-10/STAT3 mechanism. *Front Immunol*. 2019;**10**:3.
- 45 Herweijer H, Wolff JA. Progress and prospects: naked DNA gene transfer and therapy. *Gene Ther*. 2003;**10**:453–8.
- 46 Kim SS, Ye C, Kumar P, Chiu I, Subramanya S, Wu H, et al. Targeted delivery of siRNA to macrophages for anti-inflammatory treatment. *Mol Ther*. 2010;**18**:993–1001.
- 47 Collins DE, Reuter JD, Rush HG, Villano JS. Viral vector biosafety in laboratory animal research. *Comp Med*. 2017;**67**:215–21.
- 48 Haccin-Bey-Abina S, Hauer J, Lim A, Picard C, Wang GP, Berry CC, et al. Efficacy of gene therapy for X-linked severe combined immunodeficiency. *N Engl J Med*. 2010;**363**:355–64.
- 49 Gregory SM, Nazir SA, Metcalf JP. Implications of the innate immune response to adenovirus and adenoviral vectors. *Future Virol*. 2011;**6**:357–74.
- 50 Annoni A, Gregori S, Naldini L, Cantore A. Modulation of immune responses in lentiviral vector-mediated gene transfer. *Cell Immunol*. 2019;**342**:103802.
- 51 Anselmo AC, Mitragotri S. Nanoparticles in the clinic: an update. *Bioeng Transl Med*. 2019;**4**:e10143.
- 52 Hu G, Guo M, Xu J, Wu F, Fan J, Huang Q, et al. Nanoparticles targeting macrophages as potential clinical therapeutic agents against cancer and inflammation. *Front Immunol*. 2019;**10**:1998.
- 53 Reichel D, Tripathi M, Perez JM. Biological effects of nanoparticles on macrophage polarization in the tumor microenvironment. *Nanotheranostics*. 2019;**3**:66–88.
- 54 Nakanishi Y, Nakatsuji M, Seno H, Ishizu S, Akitake-Kawano R, Kanda K, et al. COX-2 inhibition alters the phenotype of tumor-associated macrophages from M2 to M1 in ApcMin/+ mouse polyps. *Carcinogenesis* 2011;**32**:1333–9.
- 55 Myers KV, Amend SR, Pienta KJ. Targeting Tyro3, Axl and MerTK (TAM receptors): implications for macrophages in the tumor microenvironment. *Mol Cancer*. 2019;**18**:94.
- 56 Moreira JD, Koch BEV, van Veen S, Walburg KV, Vrieling F, Guimaraes MPD, et al. Functional inhibition of host histone deacetylases (HDACs) enhances in vitro and in vivo anti-mycobacterial activity in human macrophages and in zebrafish. *Front Immunol*. 2020;**11**:36.
- 57 Rodell CB, Arlauckas SP, Cuccarese MF, Garriss CS, Li R, Ahmed MS, et al. TLR7/8-agonist-loaded nanoparticles promote the polarization of tumour-associated macrophages to enhance cancer immunotherapy. *Nat Biomed Eng*. 2018;**2**:578–88.
- 58 Shahbazi MA, Sedighi M, Bauleth-Ramos T, Kant K, Correia A, Poursina N, et al. Targeted reinforcement of macrophage reprogramming toward M2 polarization by IL-4-loaded hyaluronic acid particles. *ACS Omega*. 2018;**3**:18444–55.
- 59 Zidek Z, Anzenbacher P, Kmonickova E. Current status and challenges of cytokine pharmacology. *Br J Pharmacol*. 2009;**157**:342–61.
- 60 Kwon D, Cha BG, Cho Y, Min J, Park EB, Kang SJ, et al. Extra-large pore mesoporous silica nanoparticles for directing in vivo M2 macrophage polarization by delivering IL-4. *Nano Lett*. 2017;**17**:2747–56.
- 61 Thauvin C, Widmer J, Mottas I, Hocesvar S, Allemann E, Bourquin C, et al. Development of resiquimod-loaded modified PLA-based nanoparticles for cancer immunotherapy: a kinetic study. *Eur J Pharm Biopharm*. 2019;**139**:253–61.
- 62 Chenthamara D, Subramaniam S, Ramakrishnan SG, Krishnaswamy S, Essa MM, Lin FH, et al. Therapeutic efficacy of nanoparticles and routes of administration. *Biomater Res*. 2019;**23**:20.
- 63 Parsa R, Andresen P, Gillett A, Mia S, Zhang XM, Mayans S, et al. Adoptive transfer of immunomodulatory M2 macrophages prevents type 1 diabetes in NOD mice. *Diabetes* 2012;**61**:2881–92.
- 64 Shields CWT, Evans MA, Wang LL, Baugh N, Iyer S, Wu D, et al. Cellular backpacks for macrophage immunotherapy. *Sci Adv*. 2020;**6**:eaz6579.
- 65 Moroni F, Dwyer BJ, Graham C, Pass C, Bailey L, Ritchie L, et al. Safety profile of autologous macrophage therapy for liver cirrhosis. *Nat Med*. 2019;**25**:1560–5.
- 66 Epelman S, Lavine KJ, Beaudin AE, Sojka DK, Carrero JA, Calderon B, et al. Embryonic and adult-derived resident cardiac macrophages are maintained through distinct mechanisms at steady state and during inflammation. *Immunity* 2014;**40**:91–104.
- 67 Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, et al. Progressive replacement of embryo-derived cardiac macrophages with age. *J Exp Med*. 2014;**211**:2151–8.
- 68 Hulsmans M, Clauss S, Xiao L, Aguirre AD, King KR, Hanley A, et al. Macrophages facilitate electrical conduction in the heart. *Cell* 2017;**169**:510–22 e20.
- 69 Pinto AR, Paolicelli R, Salimova E, Gospocic J, Slonimsky E, Bilbao-Cortes D, et al. An abundant tissue macrophage population in the adult murine heart with a distinct alternatively-activated macrophage profile. *PLoS One* 2012;**7**:e36814.

- 70 Leid J, Carrelha J, Boukarabila H, Epelman S, Jacobsen SE, Lavine KJ. Primitive embryonic macrophages are required for coronary development and maturation. *Circ Res*. 2016;**118**:1498–511.
- 71 Ruparelina N, Godec J, Lee R, Chai JT, Dall'Armellina E, McAndrew D, et al. Acute myocardial infarction activates distinct inflammation and proliferation pathways in circulating monocytes, prior to recruitment, and identified through conserved transcriptional responses in mice and humans. *Eur Heart J*. 2015;**36**:1923–34.
- 72 Tsujioka H, Imanishi T, Ikejima H, Kuroi A, Takarada S, Tanimoto T, et al. Impact of heterogeneity of human peripheral blood monocyte subsets on myocardial salvage in patients with primary acute myocardial infarction. *J Am Coll Cardiol*. 2009;**54**:130–8.
- 73 Heidt T, Courties G, Dutta P, Sager HB, Sebas M, Iwamoto Y, et al. Differential contribution of monocytes to heart macrophages in steady-state and after myocardial infarction. *Circ Res*. 2014;**115**:284–95.
- 74 Frodermann V, Nahrendorf M. Macrophages and cardiovascular health. *Physiol Rev*. 2018;**98**:2523–69.
- 75 Mouton AJ, DeLeon-Pennell KY, Rivera Gonzalez OJ, Flynn ER, Freeman TC, Saucerman JJ, et al. Mapping macrophage polarization over the myocardial infarction time continuum. *Basic Res Cardiol*. 2018;**113**:26.
- 76 Travers JG, Kamal FA, Robbins J, Yutzey KE, Blaxall BC. Cardiac fibrosis: the fibroblast awakens. *Circ Res*. 2016;**118**:1021–40.
- 77 Dick SA, Macklin JA, Nejat S, Momen A, Clemente-Casares X, Althagafi MG, et al. Self-renewing resident cardiac macrophages limit adverse remodeling following myocardial infarction. *Nat Immunol*. 2019;**20**:29–39.
- 78 Deniset JF, Belke D, Lee WY, Jorch SK, Deppermann C, Hassanabad AF, et al. Gata6 (+) pericardial cavity macrophages relocate to the injured heart and prevent cardiac fibrosis. *Immunity* 2019;**51**:131–40 e5.
- 79 Benjamin EJ, Virani SS, Callaway CW, Chamberlain AM, Chang AR, Cheng S, et al. Heart disease and stroke statistics-2018 update: a report from the American Heart Association. *Circulation* 2018;**137**:e67–e492.
- 80 Giugliano GR, Giugliano RP, Gibson CM, Kuntz RE. Meta-analysis of corticosteroid treatment in acute myocardial infarction. *Am J Cardiol*. 2003;**91**:1055–9.
- 81 Abbate A, Trankle CR, Buckley LF, Lipinski MJ, Appleton D, Kadariya D, et al. Interleukin-1 blockade inhibits the acute inflammatory response in patients with ST-segment-elevation myocardial infarction. *J Am Heart Assoc*. 2020;**9**:e014941.
- 82 Liu J, Wang H, Li J. Inflammation and inflammatory cells in myocardial infarction and reperfusion injury: a double-edged sword. *Clin Med Insights Cardiol*. 2016;**10**:79–84.
- 83 Hoyer FF, Naxerova K, Schloss MJ, Hulsmans M, Nair AV, Dutta P, et al. Tissue-specific macrophage responses to remote injury impact the outcome of subsequent local immune challenge. *Immunity* 2019;**51**:899–914 e7.
- 84 Courties G, Heidt T, Sebas M, Iwamoto Y, Jeon D, Truelove J, et al. *in vivo* silencing of the transcription factor IRF5 reprograms the macrophage phenotype and improves infarct healing. *J Am Coll Cardiol*. 2014;**63**:1556–66.
- 85 Bejerano T, Etzion S, Elyagon S, Etzion Y, Cohen S. Nanoparticle delivery of miRNA-21 mimic to cardiac macrophages improves myocardial remodeling after myocardial infarction. *Nano Lett*. 2018;**18**:5885–91.
- 86 Yang L, Wang B, Zhou Q, Wang Y, Liu X, Liu Z, et al. MicroRNA-21 prevents excessive inflammation and cardiac dysfunction after myocardial infarction through targeting KBTBD7. *Cell Death Dis*. 2018;**9**:769.
- 87 Podaru MN, Fields L, Kainuma S, Ichihara Y, Hussain M, Ito T, et al. Reparative macrophage transplantation for myocardial repair: a refinement of bone marrow mononuclear cell-based therapy. *Basic Res Cardiol*. 2019;**114**:34.
- 88 Bajpai G, Schneider C, Wong N, Bredemeyer A, Hulsmans M, Nahrendorf M, et al. The human heart contains distinct macrophage subsets with divergent origins and functions. *Nat Med*. 2018;**24**:1234–45.
- 89 Lopez-Atalaya JP, Askew KE, Sierra A, Gomez-Nicola D. Development and maintenance of the brain's immune toolkit: Microglia and non-parenchymal brain macrophages. *Dev Neurobiol*. 2018;**78**:561–79.
- 90 Garre JM, Yang G. Contributions of monocytes to nervous system disorders. *J Mol Med (Berl)*. 2018;**96**:873–83.
- 91 Banks WA. Characteristics of compounds that cross the blood-brain barrier. *BMC Neurol*. 2009;**9**(Suppl 1):S3.
- 92 Fuhrmann M, Bittner T, Jung CK, Burgold S, Page RM, Mitteregger G, et al. Microglial Cx3cr1 knockout prevents neuron loss in a mouse model of Alzheimer's disease. *Nat Neurosci*. 2010;**13**:411–3.
- 93 International Genomics of Alzheimer's Disease C. Convergent genetic and expression data implicate immunity in Alzheimer's disease. *Alzheimer's Dementia*. 2015;**11**:658–71.
- 94 Chu F, Shi M, Zheng C, Shen D, Zhu J, Zheng X, et al. The roles of macrophages and microglia in multiple sclerosis and experimental autoimmune encephalomyelitis. *J Neuroimmunol*. 2018;**318**:1–7.
- 95 Beers DR, Appel SH. Immune dysregulation in amyotrophic lateral sclerosis: mechanisms and emerging therapies. *Lancet Neurol*. 2019;**18**:211–20.
- 96 Goldenberg MM. Multiple sclerosis review. *P T*. 2012;**37**:175–84.
- 97 Christophi GP, Panos M, Hudson CA, Christophi RL, Gruber RC, Mersich AT, et al. Macrophages of multiple sclerosis patients display deficient SHP-1 expression and enhanced inflammatory phenotype. *Lab Invest*. 2009;**89**:742–59.
- 98 Zrzavy T, Hametner S, Wimmer I, Butovsky O, Weiner HL, Lassmann H. Loss of 'homeostatic' microglia and patterns of their activation in active multiple sclerosis. *Brain* 2017;**140**:1900–13.
- 99 Locatelli G, Theodorou D, Kendirli A, Jordao MJC, Staszewski O, Phulphagar K, et al. Mononuclear phagocytes locally specify and adapt their phenotype in a multiple sclerosis model. *Nat Neurosci*. 2018;**21**:1196–208.
- 100 Moreno MA, Burns T, Yao P, Miers L, Pleasure D, Soulika AM. Therapeutic depletion of monocyte-derived cells protects from long-term axonal loss in experimental autoimmune encephalomyelitis. *J Neuroimmunol*. 2016;**290**:36–46.
- 101 Heppner FL, Greter M, Marino D, Falsig J, Raivich G, Hovelmeyer N, et al. Experimental autoimmune encephalomyelitis repressed by microglial paralysis. *Nat Med*. 2005;**11**:146–52.
- 102 Rothe T, Ipseiz N, Faas M, Lang S, Perez-Branguli F, Metzger D, et al. The nuclear receptor Nr4a1 acts as a microglia rheostat and serves as a therapeutic target in autoimmune-driven central nervous system inflammation. *J Immunol*. 2017;**198**:3878–85.
- 103 Constantinescu CS, Farooqi N, O'Brien K, Gran B. Experimental autoimmune encephalomyelitis (EAE) as a model for multiple sclerosis (MS). *Br J Pharmacol*. 2011;**164**:1079–106.
- 104 Pul R, Moharreggh-Khiabani D, Skuljec J, Skripuletz T, Garde N, Voss EV, et al. Glatiramer acetate modulates TNF-alpha and IL-10 secretion in microglia and promotes their phagocytic activity. *J Neuroimmune Pharmacol*. 2011;**6**:381–8.
- 105 Liu J, Johnson TV, Lin J, Ramirez SH, Bronich TK, Caplan S, et al. T cell independent mechanism for copolymer-1-induced neuroprotection. *Eur J Immunol*. 2007;**37**:3143–54.
- 106 Wynn DR. Enduring clinical value of copaxone(R) (Glatiramer Acetate) in multiple sclerosis after 20 years of use. *Mult Scler Int*. 2019;**2019**:7151685.
- 107 Boster A, Bartoszek MP, O'Connell C, Pitt D, Racke M. Efficacy, safety, and cost-effectiveness of glatiramer acetate in the treatment of relapsing-remitting multiple sclerosis. *Ther Adv Neurol Disord*. 2011;**4**:319–32.
- 108 Olah M, Amor S, Brouwer N, Vinet J, Eggen B, Biber K, et al. Identification of a microglia phenotype supportive of remyelination. *Glia* 2012;**60**:306–21.
- 109 Getts DR, Terry RL, Getts MT, Deffrasnes C, Muller M, van Vreden C, et al. Therapeutic inflammatory monocyte modulation using immune-modifying microparticles. *Sci Transl Med*. 2014;**6**:219ra7.
- 110 Giladi A, Wagner LK, Li H, Dorr D, Medaglia C, Paul F, et al. Cxcl10(+) monocytes define a pathogenic subset in the central nervous system during autoimmune neuroinflammation. *Nat Immunol*. 2020;**21**:525–34.
- 111 Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogava E, Majounie E, et al. TREM2 variants in Alzheimer's disease. *N Engl J Med*. 2013;**368**:117–27.
- 112 Wang S, Mustafa M, Yuede CM, Salazar SV, Kong P, Long H, et al. Anti-human TREM2 induces microglia proliferation and reduces pathology in an Alzheimer's disease model. *J Exp Med*. 2020;**217**:e20200785.
- 113 Solito E, Sastre M. Microglia function in Alzheimer's disease. *Front Pharmacol*. 2012;**3**:14.
- 114 Olmos-Alonso A, Schettters ST, Sri S, Askew K, Mancuso R, Vargas-Caballero M, et al. Pharmacological targeting of CSF1R inhibits microglial proliferation and prevents the progression of Alzheimer's-like pathology. *Brain* 2016;**139**(Pt 3):891–907.
- 115 Dagher NN, Najafi AR, Kayala KM, Elmore MR, White TE, Medeiros R, et al. Colony-stimulating factor 1 receptor inhibition prevents microglial plaque association and improves cognition in 3xTg-AD mice. *J Neuroinflammation*. 2015;**12**:139.
- 116 Spangenberg EE, Lee RJ, Najafi AR, Rice RA, Elmore MR, Blurton-Jones M, et al. Eliminating microglia in Alzheimer's mice prevents neuronal loss without modulating amyloid-beta pathology. *Brain* 2016;**139**(Pt 4):1265–81.
- 117 Lei F, Cui N, Zhou C, Chodosh J, Vavvas DG, Paschalis EI. CSF1R inhibition by a small-molecule inhibitor is not microglia specific; affecting hematopoiesis and the function of macrophages. *Proc Natl Acad Sci USA*. 2020;**117**:23336–8.
- 118 Quigley EMM. Microbiota-brain-gut axis and neurodegenerative diseases. *Curr Neurol Neurosci Rep*. 2017;**17**:94.
- 119 Erdo F, Bors LA, Farkas D, Bajza A, Gizurarson S. Evaluation of intranasal delivery route of drug administration for brain targeting. *Brain Res Bull*. 2018;**143**:155–70.
- 120 Atkinson AJ Jr. Intracerebroventricular drug administration. *Transl Clin Pharmacol*. 2017;**25**:117–24.
- 121 Geloso MC, Corvino V, Marchese E, Serrano A, Michetti F, D'Ambrosi N. The dual role of microglia in ALS: mechanisms and therapeutic approaches. *Front Aging Neurosci*. 2017;**9**:242.
- 122 Du Y, Zhao W, Thonhoff JR, Wang J, Wen S, Appel SH. Increased activation ability of monocytes from ALS patients. *Exp Neurol*. 2020;**328**:113259.

- 123 Wang H, Wang Y, Li D, Liu Z, Zhao Z, Han D, et al. VEGF inhibits the inflammation in spinal cord injury through activation of autophagy. *Biochem Biophys Res Commun*. 2015;464:453–8.
- 124 Wang Y, Duan W, Wang W, Di W, Liu Y, Liu Y, et al. sAAV9-VEGF prolongs the survival of transgenic ALS mice by promoting activation of M2 microglia and the PI3K/Akt pathway. *Brain Res*. 2016;1648(Pt A):1–10.
- 125 Ramaglia V, Sheikh-Mohamed S, Legg K, Park C, Rojas OL, Zandee S, et al. Multiplexed imaging of immune cells in staged multiple sclerosis lesions by mass cytometry. *Elife*. 2019;8:e48051.
- 126 Yurek D, Hasselrot U, Seseoglu-Laird O, Padeigimas L, Cooper M. Intracerebral injections of DNA nanoparticles encoding for a therapeutic gene provide partial neuroprotection in an animal model of neurodegeneration. *Nanomedicine*. 2017;13:2209–17.
- 127 Hu N, Shi X, Zhang Q, Liu W, Zhu Y, Wang Y, et al. Special interstitial route can transport nanoparticles to the brain bypassing the blood-brain barrier. *Nano Research*. 2019;12:2760–5.
- 128 Ali J, Ali M, Baboota S, Sahani JK, Ramassamy C, Dao L, et al. Potential of nanoparticulate drug delivery systems by intranasal administration. *Curr Pharm Des*. 2010;16:1644–53.
- 129 Sierro F, Evrard M, Rizzetto S, Melino M, Mitchell AJ, Florido M, et al. A liver capsular network of monocyte-derived macrophages restricts hepatic dissemination of intraperitoneal bacteria by neutrophil recruitment. *Immunity* 2017;47:374–88 e6.
- 130 Blieriot C, Ginhoux F. Understanding the heterogeneity of resident liver macrophages. *Front Immunol*. 2019;10:2694.
- 131 Guillot A, Tacke F. Liver macrophages: old dogmas and new insights. *Hepatol Commun*. 2019;3:730–43.
- 132 Wang J, Kubes P. A reservoir of mature cavity macrophages that can rapidly invade visceral organs to affect tissue repair. *Cell* 2016;165:668–78.
- 133 Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity* 2013;38:79–91.
- 134 Hoeffel G, Chen J, Lavin Y, Low D, Almeida FF, See P, et al. C-Myb(+) erythro-myeloid progenitor-derived fetal monocytes give rise to adult tissue-resident macrophages. *Immunity* 2015;42:665–78.
- 135 Scott CL, Zheng F, De Baetselier P, Martens L, Saey Y, De Prijck S, et al. Bone marrow-derived monocytes give rise to self-renewing and fully differentiated Kupffer cells. *Nat Commun*. 2016;7:10321.
- 136 Blieriot C, Dupuis T, Jouvion G, Eberl G, Disson O, Lecuit M. Liver-resident macrophage necroptosis orchestrates type 1 microbicidal inflammation and type-2-mediated tissue repair during bacterial infection. *Immunity* 2015;42:145–58.
- 137 Beattie L, Sawtell A, Mann J, Frame TCM, Teal B, de Labastida RF, et al. Bone marrow-derived and resident liver macrophages display unique transcriptomic signatures but similar biological functions. *J Hepatol*. 2016;65:758–68.
- 138 David BA, Rezende RM, Antunes MM, Santos MM, Freitas Lopes MA, Diniz AB, et al. Combination of mass cytometry and imaging analysis reveals origin, location, and functional repopulation of liver myeloid cells in mice. *Gastroenterology* 2016;151:1176–91.
- 139 Soysa R, Lampert S, Yuen S, Douglass AN, Li W, Pfeffer K, et al. Fetal origin confers radioresistance on liver macrophages via p21(cip1/WAF1). *J Hepatol*. 2019;71:553–62.
- 140 Zigmond E, Samia-Grinberg S, Pasmannik-Chor M, Brazowski E, Shibolet O, Halpern Z, et al. Infiltrating monocyte-derived macrophages and resident kupffer cells display different ontogeny and functions in acute liver injury. *J Immunol*. 2014;193:344–53.
- 141 Mass E, Ballesteros I, Farlik M, Halbritter F, Gunther P, Crozet L, et al. Specification of tissue-resident macrophages during organogenesis. *Science* 2016;353:aaf4238.
- 142 Scott CL, T'Jonck W, Martens L, Todorov H, Sichen D, Soen B, et al. The transcription factor ZEB2 is required to maintain the tissue-specific identities of macrophages. *Immunity* 2018;49(2):312–25 e5.
- 143 Bonnardel J, T'Jonck W, Gaublomme D, Browaeys R, Scott CL, Martens L, et al. Stellate cells, hepatocytes, and endothelial cells imprint the kupffer cell identity on monocytes colonizing the liver macrophage niche. *Immunity* 2019;51:638–54 e9.
- 144 Sakai M, Troutman TD, Seidman JS, Ouyang Z, Spann NJ, Abe Y, et al. Liver-derived signals sequentially reprogram myeloid enhancers to initiate and maintain kupffer cell identity. *Immunity* 2019;51:655–70 e8.
- 145 MacParland SA, Liu JC, Ma XZ, Innes BT, Bartczak AM, Gage BK, et al. Single cell RNA sequencing of human liver reveals distinct intrahepatic macrophage populations. *Nat Commun*. 2018;9:4383.
- 146 Aizarani N, Saviano A, Sagar ML, Durand S, Herman JS, et al. A human liver cell atlas reveals heterogeneity and epithelial progenitors. *Nature* 2019;572:199–204.
- 147 Zhao J, Zhang S, Liu Y, He X, Qu M, Xu G, et al. Single-cell RNA sequencing reveals the heterogeneity of liver-resident immune cells in human. *Cell Discov*. 2020;6:22.
- 148 Duffield JS, Forbes SJ, Constandinou CM, Clay S, Partolina M, Vuthoori S, et al. Selective depletion of macrophages reveals distinct, opposing roles during liver injury and repair. *J Clin Invest*. 2005;115:56–65.
- 149 Xiong X, Kuang H, Ansari S, Liu T, Gong J, Wang S, et al. Landscape of intercellular crosstalk in healthy and NASH liver revealed by single-cell secretome gene analysis. *Mol Cell*. 2019;75:644–60 e5.
- 150 Seidman JS, Troutman TD, Sakai M, Gola A, Spann NJ, Bennett H, et al. Niche-specific reprogramming of epigenetic landscapes drives myeloid cell diversity in nonalcoholic steatohepatitis. *Immunity* 2020;52:1057–74 e7.
- 151 Triantafyllou E, Pop OT, Possamai LA, Wilhelm A, Liaskou E, Singanayagam A, et al. MerTK expressing hepatic macrophages promote the resolution of inflammation in acute liver failure. *Gut* 2018;67:333–47.
- 152 Robert O, Boujedidi H, Bigorgne A, Ferrere G, Voican CS, Vettorazzi S, et al. Decreased expression of the glucocorticoid receptor-GILZ pathway in Kupffer cells promotes liver inflammation in obese mice. *J Hepatol*. 2016;64:916–24.
- 153 Bartneck M, Scheida KM, Warzecha KT, Rizzo LY, Hittatiya K, Luedde T, et al. Fluorescent cell-traceable dexamethasone-loaded liposomes for the treatment of inflammatory liver diseases. *Biomaterials* 2015;37:367–82.
- 154 Svendsen P, Graversen JH, Etzerodt A, Hager H, Roge R, Gronbaek H, et al. Antibody-directed glucocorticoid targeting to CD163 in M2-type macrophages attenuates fructose-induced liver inflammatory changes. *Mol Ther Methods Clin Dev*. 2017;4:50–61.
- 155 He C, Yin L, Tang C, Yin C. Multifunctional polymeric nanoparticles for oral delivery of TNF-alpha siRNA to macrophages. *Biomaterials* 2013;34:2843–54.
- 156 Azzimato V, Jager J, Chen P, Morgantini C, Levi L, Barreby E, et al. Liver macrophages inhibit the endogenous antioxidant response in obesity-associated insulin resistance. *Sci Transl Med*. 2020;12:eaa9709.
- 157 Traber PG, Chou H, Zomer E, Hong F, Klyosov A, Fiel MI, et al. Regression of fibrosis and reversal of cirrhosis in rats by galectin inhibitors in thioacetamide-induced liver disease. *PLoS One* 2013;8:e75361.
- 158 Traber PG, Zomer E. Therapy of experimental NASH and fibrosis with galectin inhibitors. *PLoS One* 2013;8:e83481.
- 159 Volarevic V, Milovanovic M, Ljujic B, Pejnovic N, Arsenijevic N, Nilsson U, et al. Galectin-3 deficiency prevents concanavalin A-induced hepatitis in mice. *Hepatology* 2012;55:1954–64.
- 160 Mossanen JC, Krenkel O, Ergen C, Govaere O, Liepelt A, Puengel T, et al. Chemokine (C-C motif) receptor 2-positive monocytes aggravate the early phase of acetaminophen-induced acute liver injury. *Hepatology* 2016;64:1667–82.
- 161 Krenkel O, Puengel T, Govaere O, Abdallah AT, Mossanen JC, Kohlhepp M, et al. Therapeutic inhibition of inflammatory monocyte recruitment reduces steatohepatitis and liver fibrosis. *Hepatology* 2018;67:1270–83.
- 162 Anstee QM, Neuschwander-Tetri BA, Wong VW, Abdelmalek MF, Younossi ZM, Yuan J, et al. Cenicriviroc for the treatment of liver fibrosis in adults with nonalcoholic steatohepatitis: AURORA Phase 3 study design. *Contemp Clin Trials*. 2020;89:105922.
- 163 Starkey Lewis P, Campana L, Aleksieva N, Cartwright JA, Mackinnon A, O'Duibhir E, et al. Alternatively activated macrophages promote resolution of necrosis following acute liver injury. *J Hepatol*. 2020;73:349–60.
- 164 Thomas JA, Pope C, Wojtacha D, Robson AJ, Gordon-Walker TT, Hartland S, et al. Macrophage therapy for murine liver fibrosis recruits host effector cells improving fibrosis, regeneration, and function. *Hepatology* 2011;53:2003–15.
- 165 Haideri SS, McKinnon AC, Taylor AH, Kirkwood P, Starkey Lewis PJ, O'Duibhir E, et al. Injection of embryonic stem cell derived macrophages ameliorates fibrosis in a murine model of liver injury. *NPJ Regen Med*. 2017;2:14.
- 166 Hutchinson JA, Riquelme P, Sawitzki B, Tomiuk S, Miqueu P, Zuhayra M, et al. Cutting Edge: Immunological consequences and trafficking of human regulatory macrophages administered to renal transplant recipients. *J Immunol*. 2011;187:2072–8.
- 167 Bird TG, Lu WY, Boulter L, Gordon-Keylock S, Ridgway RA, Williams MJ, et al. Bone marrow injection stimulates hepatic ductular reactions in the absence of injury via macrophage-mediated TWEAK signaling. *Proc Natl Acad Sci U S A*. 2013;110:6542–7.
- 168 Sharkey J, Starkey Lewis PJ, Barrow M, Alwahsh SM, Noble J, Livingstone E, et al. Functionalized superparamagnetic iron oxide nanoparticles provide highly efficient iron-labeling in macrophages for magnetic resonance-based detection *in vivo*. *Cytherapy*. 2017;19:555–69.
- 169 Udalova IA, Mantovani A, Feldmann M. Macrophage heterogeneity in the context of rheumatoid arthritis. *Nat Rev Rheumatol*. 2016;12:472–85.
- 170 McInnes IB, Buckley CD, Isaacs JD. Cytokines in rheumatoid arthritis - shaping the immunological landscape. *Nat Rev Rheumatol*. 2016;12(1):63–8.
- 171 Smith MD. The normal synovium. *Open Rheumatol J*. 2011;5:100–6.
- 172 Alivernini S, Tolusso B, Petricca L, Bui L, Di Sante G, Peluso G, et al. Synovial features of patients with rheumatoid arthritis and psoriatic arthritis in clinical and ultrasound remission differ under anti-TNF therapy: a clue to interpret different chances of relapse after clinical remission? *Ann Rheum Dis*. 2017;76:1228–36.
- 173 Misharin AV, Cuda CM, Saber R, Turner JD, Gierut AK, Haines GK 3rd, et al. Non-classical Ly6C(-) monocytes drive the development of inflammatory arthritis in mice. *Cell Rep*. 2014;9:591–604.

- 174 Culemann S, Gruneboom A, Nicolas-Avila JA, Weidner D, Lammle KF, Rothe T, et al. Locally renewing resident synovial macrophages provide a protective barrier for the joint. *Nature* 2019;**572**:670–5.
- 175 Alivernini S, MacDonald L, Elmesari A, Finlay S, Toluoso B, Gigante MR, et al. Distinct synovial tissue macrophage subsets regulate inflammation and remission in rheumatoid arthritis. *Nat Med* 2020;**26**:1295–306.
- 176 Zhang F, Wei K, Slowikowski K, Fonseka CY, Rao DA, Kelly S, et al. Defining inflammatory cell states in rheumatoid arthritis joint synovial tissues by integrating single-cell transcriptomics and mass cytometry. *Nat Immunol*. 2019;**20**:928–42.
- 177 Kuo D, Ding J, Cohn IS, Zhang F, Wei K, Rao DA, et al. HBEGF(+) macrophages in rheumatoid arthritis induce fibroblast invasiveness. *Sci Transl Med*. 2019;**11**:eaau8587.
- 178 Swanson CD, Akama-Garren EH, Stein EA, Petralia JD, Ruiz PJ, Edalati A, et al. Inhibition of epidermal growth factor receptor tyrosine kinase ameliorates collagen-induced arthritis. *J Immunol*. 2012;**188**:3513–21.
- 179 Cook AD, Pobjoy J, Steidl S, Durr M, Braine EL, Turner AL, et al. Granulocyte-macrophage colony-stimulating factor is a key mediator in experimental osteoarthritis pain and disease development. *Arthritis Res Ther*. 2012;**14**:R199.
- 180 Hamilton JA. GM-CSF as a target in inflammatory/autoimmune disease: current evidence and future therapeutic potential. *Expert Rev Clin Immunol*. 2015;**11**:457–65.
- 181 Hamilton JA, Cook AD, Tak PP. Anti-colony-stimulating factor therapies for inflammatory and autoimmune diseases. *Nat Rev Drug Discov*. 2016;**16**:53–70.
- 182 Campbell IK, Rich MJ, Bischof RJ, Dunn AR, Grail D, Hamilton JA. Protection from collagen-induced arthritis in granulocyte-macrophage colony-stimulating factor-deficient mice. *J Immunol*. 1998;**161**:3639–44.
- 183 Cook AD, Braine EL, Campbell IK, Rich MJ, Hamilton JA. Blockade of collagen-induced arthritis post-onset by antibody to granulocyte-macrophage colony-stimulating factor (GM-CSF): requirement for GM-CSF in the effector phase of disease. *Arthritis Res*. 2001;**3**:293–8.
- 184 Hamilton JA. GM-CSF in inflammation. *J Exp Med*. 2020;**217**:1.
- 185 Cook AD, Hamilton JA. Investigational therapies targeting the granulocyte macrophage colony-stimulating factor receptor- α in rheumatoid arthritis: focus on mavrilimumab. *Ther Adv Musculoskelet Dis*. 2018;**10**:29–38.
- 186 Kong JS, Park JH, Yoo SA, Kim KM, Bae YJ, Park YJ, et al. Dynamic transcriptome analysis unveils key proresolving factors of chronic inflammatory arthritis. *J Clin Invest*. 2020;**130**:3974–86.
- 187 Mathivanan S, Simpson RJ. ExoCarta: a compendium of exosomal proteins and RNA. *Proteomics* 2009;**9**:4997–5000.
- 188 Kaplan A, Bueno M, Fournier AE. Extracellular functions of 14-3-3 adaptor proteins. *Cell Signal*. 2017;**31**:26–30.
- 189 Dobson M, Ramakrishnan G, Ma S, Kaplun L, Balan V, Fridman R, et al. Bimodal regulation of FoxO3 by AKT and 14-3-3. *Biochim Biophys Acta*. 2011;**1813**:1453–64.
- 190 Stoelcklin G, Stubbs T, Kedersha N, Wax S, Rigby WF, Blackwell TK, et al. MK2-induced tristetraprolin:14-3-3 complexes prevent stress granule association and ARE-mRNA decay. *EMBO J*. 2004;**23**:1313–24.
- 191 Schaljo B, Kratochvill F, Gratz N, Sadzak I, Sauer I, Hammer M, et al. Tristetraprolin is required for full anti-inflammatory response of murine macrophages to IL-10. *J Immunol*. 2009;**183**:1197–206.
- 192 Jain S, Tran TH, Amiji M. Macrophage repolarization with targeted alginate nanoparticles containing IL-10 plasmid DNA for the treatment of experimental arthritis. *Biomaterials* 2015;**61**:162–77.
- 193 Gonzalez H, Hagerling C, Werb Z. Roles of the immune system in cancer: from tumor initiation to metastatic progression. *Genes Dev*. 2018;**32**:1267–84.
- 194 Helm O, Held-Feindt J, Grage-Griebenow E, Reiling N, Ungefroren H, Vogel I, et al. Tumor-associated macrophages exhibit pro- and anti-inflammatory properties by which they impact on pancreatic tumorigenesis. *Int J Cancer*. 2014;**135**:843–61.
- 195 Bucana C, Hoyer LC, Hobbs B, Breesman S, McDaniel M, Hanna MG Jr. Morphological evidence for the translocation of lysosomal organelles from cytotoxic macrophages into the cytoplasm of tumor target cells. *Cancer Res*. 1976;**36**:4444–58.
- 196 Chen Y, Song Y, Du W, Gong L, Chang H, Zou Z. Tumor-associated macrophages: an accomplice in solid tumor progression. *J Biomed Sci*. 2019;**26**:78.
- 197 Liu Y, Cao X. The origin and function of tumor-associated macrophages. *Cell Mol Immunol*. 2015;**12**:1–4.
- 198 Adams S, Kozhaya L, Martiniuk F, Meng TC, Chiriboga L, Liebes L, et al. Topical TLR7 agonist imiquimod can induce immune-mediated rejection of skin metastases in patients with breast cancer. *Clin Cancer Res*. 2012;**18**:6748–57.
- 199 Salazar LG, Lu H, Reichow JL, Childs JS, Coveler AL, Higgins DM, et al. Topical imiquimod plus nab-paclitaxel for breast cancer cutaneous metastases: a phase 2 clinical trial. *JAMA Oncol*. 2017;**3**:969–73.
- 200 Pachman DR, Barton DL, Clayton AC, McGovern RM, Jefferies JA, Novotny PJ, et al. Randomized clinical trial of imiquimod: an adjunct to treating cervical dysplasia. *Am J Obstet Gynecol*. 2012;**206**:42 e1–7.
- 201 Grimm C, Polterauer S, Natter C, Rahhal J, Hefler L, Tempfer CB, et al. Treatment of cervical intraepithelial neoplasia with topical imiquimod: a randomized controlled trial. *Obstet Gynecol*. 2012;**120**:152–9.
- 202 Rook AH, Gelfand JM, Wysocka M, Troxel AB, Benoit B, Surber C, et al. Topical resiquimod can induce disease regression and enhance T-cell effector functions in cutaneous T-cell lymphoma. *Blood* 2015;**126**:1452–61.
- 203 Beatty GL, Torigan DA, Chiorean EG, Saboury B, Brothers A, Alavi A, et al. A phase I study of an agonist CD40 monoclonal antibody (CP-870,893) in combination with gemcitabine in patients with advanced pancreatic ductal adenocarcinoma. *Clin Cancer Res*. 2013;**19**:6286–95.
- 204 Beatty GL, Chiorean EG, Fishman MP, Saboury B, Teitelbaum UR, Sun W, et al. CD40 agonists alter tumor stroma and show efficacy against pancreatic carcinoma in mice and humans. *Science* 2011;**331**:1612–6.
- 205 Calvo A, Joensuu H, Sebastian M, Naing A, Bang Y-J, Martin M, et al. Phase Ib/II study of lacnotuzumab (MCS110) combined with spartalizumab (PDR001) in patients (pts) with advanced tumors. *J Clin Oncol* 2018;**36**(15_suppl):3014.
- 206 Autio KA, Klebanoff CA, Schaer D, Kauh JS, Slovin SF, Blinder VS, et al. Phase I study of LY3022855, a colony-stimulating factor-1 receptor (CSF-1R) inhibitor, in patients with metastatic breast cancer (MBC) or metastatic castration-resistant prostate cancer (MCRPC). *J Clin Oncol* 2019;**37**(15_suppl):2548.
- 207 Strachan DC, Ruffell B, Oei Y, Bissell MJ, Coussens LM, Pryer N, et al. CSF1R inhibition delays cervical and mammary tumor growth in murine models by attenuating the turnover of tumor-associated macrophages and enhancing infiltration by CD8(+) T cells. *Oncoimmunology*. 2013;**2**:e26968.
- 208 Pyonteck SM, Akkari L, Schuhmacher AJ, Bowman RL, Sevenich L, Quail DF, et al. CSF-1R inhibition alters macrophage polarization and blocks glioma progression. *Nat Med*. 2013;**19**:1264–72.
- 209 Ao JY, Zhu XD, Chai ZT, Cai H, Zhang YY, Zhang KZ, et al. Colony-stimulating factor 1 receptor blockade inhibits tumor growth by altering the polarization of tumor-associated macrophages in hepatocellular carcinoma. *Mol Cancer Ther*. 2017;**16**:1544–54.
- 210 Gomez-Roca CA, Cassier PA, Italiano A, Cannarile M, Ries C, Brillouet A, et al. Phase I study of RG7155, a novel anti-CSF1R antibody, in patients with advanced/metastatic solid tumors. *J Clin Oncol* 2015;**33**(15_suppl):3005.
- 211 Gomez-Roca CA, Italiano A, Le Tourneau C, Cassier PA, Toulmonde M, D'Angelo SP, et al. Phase I study of emactuzumab single agent or in combination with paclitaxel in patients with advanced/metastatic solid tumors reveals depletion of immunosuppressive M2-like macrophages. *Ann Oncol*. 2019;**30**:1381–92.
- 212 Quail DF, Bowman RL, Akkari L, Quick ML, Schuhmacher AJ, Huse JT, et al. The tumor microenvironment underlies acquired resistance to CSF-1R inhibition in gliomas. *Science* 2016;**352**:aad3018.
- 213 Kawai T, Akira S. Signaling to NF- κ B by Toll-like receptors. *Trends Mol Med*. 2007;**13**:460–9.
- 214 Schon MP, Schon M. Imiquimod: mode of action. *Br J Dermatol*. 2007;**157**(Suppl 2):8–13.
- 215 Zhao J, Zhang Z, Xue Y, Wang G, Cheng Y, Pan Y, et al. Anti-tumor macrophages activated by ferumoxytol combined or surface-functionalized with the TLR3 agonist poly (I : C) promote melanoma regression. *Theranostics*. 2018;**8**:6307–21.
- 216 van Kooten C, Banchereau J. CD40-CD40 ligand. *J Leukoc Biol*. 2000;**67**:2–17.
- 217 Piechotta M, Berghoff AS. New emerging targets in cancer immunotherapy: the role of Cluster of Differentiation 40 (CD40/TNFR5). *ESMO Open*. 2019;**4**(Suppl 3):e000510.
- 218 Bubna AK. Vorinostat-an overview. *Indian J Dermatol*. 2015;**60**:419.
- 219 Peng H, Chen B, Huang W, Tang Y, Jiang Y, Zhang W, et al. Reprogramming tumor-associated macrophages to reverse EGFR(T790M) resistance by dual-targeting codelivery of gefitinib/vorinostat. *Nano Lett*. 2017;**17**:7684–90.
- 220 Kaneda MM, Cappello P, Nguyen AV, Ralainirina N, Hardamon CR, Foubert P, et al. Macrophage PI3Kgamma drives pancreatic ductal adenocarcinoma progression. *Cancer Discov*. 2016;**6**:870–85.
- 221 Sullivan RJ, Hong DS, Tolcher AW, Patnaik A, Shapiro G, Chmielowski B, et al. Initial results from first-in-human study of IPI-549, a tumor macrophage-targeting agent, combined with nivolumab in advanced solid tumors. *J Clin Oncol* 2018;**36**(15_suppl):3013.
- 222 Zhang F, Parayath NN, Ene CI, Stephan SB, Koehne AL, Coon ME, et al. Genetic programming of macrophages to perform anti-tumor functions using targeted mRNA nanocarriers. *Nat Commun*. 2019;**10**:3974.
- 223 Song Y, Tang C, Yin C. Combination antitumor immunotherapy with VEGF and PIGF siRNA via systemic delivery of multi-functionalized nanoparticles to tumor-associated macrophages and breast cancer cells. *Biomaterials* 2018;**185**:117–32.
- 224 Shobaki N, Sato Y, Suzuki Y, Okabe N, Harashima H. Manipulating the function of tumor-associated macrophages by siRNA-loaded lipid nanoparticles for cancer immunotherapy. *J Control Release*. 2020;**325**:235–48.

- 225 He XY, Liu BY, Xu C, Zhuo RX, Cheng SX. A multi-functional macrophage and tumor targeting gene delivery system for the regulation of macrophage polarity and reversal of cancer immunoresistance. *Nanoscale*. 2018;**10**:15578–87.
- 226 Cai X, Yin Y, Li N, Zhu D, Zhang J, Zhang CY, et al. Re-polarization of tumor-associated macrophages to pro-inflammatory M1 macrophages by microRNA-155. *J Mol Cell Biol*. 2012;**4**:341–3.
- 227 Ray M, Lee YW, Hardie J, Mout R, Yesilbag Tonga G, Farkas ME, et al. CRISPRed macrophages for cell-based cancer immunotherapy. *Bioconjug Chem*. 2018;**29**:445–50.
- 228 Andreessen R, Hennemann B, Krause SW. Adoptive immunotherapy of cancer using monocyte-derived macrophages: rationale, current status, and perspectives. *J Leukoc Biol*. 1998;**64**:419–26.