



POSTER PRESENTATION

Open Access

# Caspase-1 is active since the early phase of rheumatoid arthritis

R Cascão<sup>1</sup>, J Polido-Pereira<sup>1,2</sup>, H Canhão<sup>1,2</sup>, A M Rodrigues<sup>1,2</sup>, M Navalho<sup>1,3</sup>, H Raquel<sup>1</sup>, A F Mourão<sup>1,4</sup>,  
C Resende<sup>2</sup>, J E Fonseca<sup>1,2\*</sup>, L F Moita<sup>1\*</sup>

From 5th European Workshop on Immune-Mediated Inflammatory Diseases  
Sitges-Barcelona, Spain. 1-3 December 2010

## Introduction

Rheumatoid arthritis (RA) is a chronic autoimmune disease characterized by joint destruction. We have previously reported an increase in interleukin (IL)-1 $\beta$  levels in the serum of very early RA and in the synovial fluid (SF) of RA patients. Therefore, the molecular mechanisms that regulate IL-1 $\beta$  activation, such as caspase-1, might play a relevant role and have potential as therapeutic targets in RA.

## Aim

The main goal of this study was to evaluate the state of activation of caspase-1 in early polyarthritis and established RA patients.

## Patients and methods

Blood samples were collected from 12 untreated early polyarthritis patients with less than 12 months of disease duration, 12 methotrexate (MTX) treated established RA patients and from 10 healthy controls. Caspase-1 activity was assessed using the Caspase Assay kit.

## Results

We found that both early polyarthritis and established RA patients have higher levels of active caspase-1 than healthy controls. Surprisingly, caspase-1 activation levels in patient cells remain unchanged when cells were stimulated with *E. coli* and ATP for 4h and 24h, whereas in healthy control cells caspase-1 levels are significantly increased after stimulation. We observed no significant differences when comparing early polyarthritis with established RA samples. However, in RA samples from patients with higher disease activity (DAS28>5.2) the active caspase-1 ratio is higher.

## Conclusions

Our results suggest that leukocytes in both early polyarthritis and established RA patients have high baseline levels of activated caspase-1, which cannot be significantly increased by bacterial and ATP stimulation. We also show that MTX treatment does not affect caspase-1 activation. Altogether, our data supports the hypothesis that caspase-1 is already activated in the early phase of RA.

## Author details

<sup>1</sup>Instituto de Medicina Molecular, Faculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal. <sup>2</sup>Rheumatology Dept., Centro Hospitalar de Lisboa Norte, EPE, Hospital de Santa Maria, Lisbon, Portugal. <sup>3</sup>Radiology Dept., Hospital da Luz, Lisbon, Portugal. <sup>4</sup>Rheumatology Dept., Centro Hospitalar de Lisboa Ocidental, EPE, Hospital Egas Moniz, Lisbon, Portugal.

Published: 25 November 2010

doi:10.1186/1479-5876-8-S1-P35

**Cite this article as:** Cascão et al.: Caspase-1 is active since the early phase of rheumatoid arthritis. *Journal of Translational Medicine* 2010 **8**(Suppl 1):P35.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Instituto de Medicina Molecular, Faculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal  
Full list of author information is available at the end of the article