

Research

Open Access

Sandwich hybridisation assay for quantitative detection of yeast RNAs in crude cell lysates

Jari Rautio¹, Kim Bundvig Barken², Juhani Lahdenperä¹, Antje Breitenstein¹, Søren Molin³ and Peter Neubauer*³

Address: ¹Bioprocess Engineering Laboratory, Department of Process and Environmental Engineering and Biocenter Oulu, University of Oulu, Finland, ²Novozymes A/S, Smørmosevej 25, DK2880 Bagsvaerd, Denmark and ³BioCentrum-DTU, Building 301, The Technical University of Denmark, DK-2800 Lyngby, Denmark

Email: Jari Rautio - jari.rautio@oulu.fi; Kim Bundvig Barken - kibb@novozymes.com; Juhani Lahdenperä - juhani.lahdenpera@perkinelmer.com; Antje Breitenstein - antje.breitenstein@scanbec.com; Søren Molin - soeren.molin@biocentrum.dtu.dk; Peter Neubauer* - peter.neubauer@oulu.fi

* Corresponding author

Published: 28 April 2003

Received: 20 March 2003

Microbial Cell Factories 2003, 2:4

Accepted: 28 April 2003

This article is available from: <http://www.microbialcellfactories.com/content/2/1/4>

© 2003 Rautio et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: A rapid microtiter plate based sandwich hybridization assay was developed for detection and quantification of single RNA species using magnetic beads. Following solution hybridization target RNA molecules were collected by biotin-streptavidin affinity binding and detected by fluorescence signal generated by alkaline phosphatase. The 18S rRNA and SUC2 mRNA of *Saccharomyces cerevisiae* were used as model RNA target molecules.

Results: The sensitivity of the assay was approximately 1.2×10^9 (2 fmol) molecules of target RNA. The developed method was feasible with crude cell lysates of *S. cerevisiae carlsbergensis* and was evaluated by measuring the levels of 18S rRNA during cell growth and SUC2 mRNA under repressive and inductive conditions. The 18S rRNA expression level followed the changes in the specific growth rate. SUC2 mRNA levels were in good correlation with the measured invertase enzyme activities.

Conclusions: The here presented sandwich hybridisation method was successfully applied for monitoring the amounts of ribosomal RNA and mRNA with high expression level in shake flask cultivation conditions. Sandwich hybridisation method offers a fast and convenient tool for following single key RNA species of interest in the production conditions.

Background

Development of novel methods for fast, sensitive and reliable RNA quantification has recently got increasing attention. Conventional methods for RNA analysis, like Northern and slot blot hybridization are in general time consuming, laborious and allow only relative quantification within a narrow concentration range. More novel methods for RNA analysis e.g. RT-PCR and real time RT-PCR are highly sensitive, but also slightly susceptible to

experimental interferences, like template inhibition due to insufficient purification [2] and lack accuracy for quantification due to biases connected to PCR and reverse transcription reactions, which in general are accepted errors connected to these methods.

The sandwich hybridization method is a suitable alternative for RNA quantification. The method is based on the detection of hybridization events between two specific




Figure 1

Principle of the sandwich hybridization assay. In solution hybridization the biotin labeled capture probe and the digoxigenin labeled detection probe are first hybridized with target RNA (a) followed by hybrid immobilization on magnetic beads and detection with anti-DIG – alkaline phosphatase FAB fragments (b). Alkaline phosphatase cleaves BBTP (2'-[2-benzothiazoyl]-6'-hydroxybenzothiazole phosphate, Atto-Phos[®]) to inorganic phosphate (Pi) and BBT (2'-[2-benzothiazoyl]-6'-hydroxybenzothiazole)

oligonucleotide probes and the target nucleic acids. The capture probe is used to immobilize the target sequence on a solid support and the detection probe is labeled with a detectable marker (see Fig. 1). Sandwich hybridization is relatively sensitive and can be performed with crude biological samples [20].

Sandwich hybridization assays from crude cell samples or in connection to PCR have been extensively used in clinical diagnostics for detection of nucleic acids from bacteria [4,6,7,13,15,21,23], viruses [1,3,9,11,14,17,25] of gene mutations [5] and for cell typing [19]. The aim in our laboratory is to apply this method for monitoring of bioprocesses. The sandwich hybridization method would be ideal for measuring the levels of specific mRNAs in yeast and bacterial cells during the fermentation processes.

Here we describe a sandwich hybridization method for quantification of RNA to be applied for measuring levels of specific RNAs in yeast cells as an informative tool for the control and state analysis of bioprocesses. The developed method is evaluated by measuring the levels of *Saccharomyces cerevisiae carlsbergensis* 18S rRNA and SUC2 mRNA in shake flask experiments.

Results

The aim of the present study was the development of a fast and reliable method for quantification of single RNA species in solution using a sandwich hybridization assay applicable for analysis of bioprocesses. The influence of different parameters like probe concentrations, concentrations of the components of the hybridization solution, reaction times, amount of beads and hybridization temperatures were investigated to optimize the hybridiza-

tion protocol and the sensitivity level. The 18S rRNA and SUC2 mRNA encoding invertase enzyme of *S. cerevisiae* were used as model RNA target molecules within this study.

Detection limit and linear range of the developed sandwich hybridization assay

The sensitivity level and the linear range of the assay were determined with the solution based sandwich hybridization system using $3.1 \times 10^6 - 7.2 \times 10^{11}$ (5 amol – 1.2 pmol) *in vitro* transcribed 18S rRNA molecules of *S. cerevisiae* as target. The biotin labeled oligonucleotide probe 18S rRNA-400 was used as capture probe and the digoxigenin labeled oligonucleotide probe 18S rRNA-1302 was used for detection. The signals of the sandwich hybridization method were compared to those of the slot blot hybridization.

1.2×10^9 (2 fmol) target molecules in the solution hybridization gave a signal that was 4-fold above the noise level. The reaction was linear up to 5.9×10^{11} (980 fmol) target molecules (Fig 2.)

The same amounts of 18S rRNA target molecules were immobilized onto a filter membrane in slot blot hybridization assay. Digoxigenin labeled 18S rRNA-1302 probe was used for chemiluminescence detection of the target molecules. Approximately the same amount of target molecules as in sandwich hybridization assay gave a signal that was 4-fold above the noise level. With the used detection system the linear range of the slot blot method was between 1.2×10^9 and 3×10^{10} (2 – 50 fmol) target molecules.

Real-time PCR experiments were performed with 18S rRNA ranging from $2 \times 10^4 - 6 \times 10^9$ target molecules (0.04 amol – 10 fmol). Detection limit was estimated to be 4×10^5 molecules (0.6 amol) and the linear range was greater than 4 orders of magnitude, making this method most sensitive with widest linear range.

Comparison of hybridization with crude cell lysates and extracted RNA

To create a fast analytical system it was investigated, whether a separate total RNA extraction step is necessary or whether crude cell lysates could be used as sample material. The assay was performed using either crude lysates of $7 \times 10^4 - 1.2 \times 10^9$ exponentially grown *S. cerevisiae carlsbergensis* cells or total RNA extracted from the same amount of cells. 18S rRNA was used as target for method comparison since it is abundant in the cell.

Fig. 3 shows that when less than 3×10^7 cells as crude lysate were added to the hybridization solution the signal of the respective target molecules was up to 60% higher




Figure 2

Comparison of sandwich solution hybridization with slot blot filter hybridization assay and real-time PCR. In **a)** $3.1 \times 10^7 - 7.2 \times 10^{11}$ 18S rRNA target molecules were added to the sandwich hybridization solution (○) or to the membrane for slot blot hybridization (●). Fluorescence (sandwich assay) and chemiluminescence (slot blot assay) signals were both generated by anti-DIG-alkaline phosphatase-FAB fragments. In **b)** $2 \times 10^5 - 6 \times 10^{10}$ 18S rRNA target molecules were amplified by the real-time PCR reaction, using SYBR-green as detection dye (Δ). Ct is number of cycles needed to reach threshold fluorescence signal level. The error bars show the \pm SD of three parallel samples.

when compared to signal measured from total RNA extracted from the same amount of cells. When the amount of crude cell lysate was further increased the signal started to decrease probably due to unspecific reactions of the crude cell material. With total RNA the same inhibition effect was observed when more than 2.5×10^8 yeast cells were used for extraction.

The effect of biological material

The effect of biological material on the signal of the solution hybridization was studied by incubating various amounts ($2 \times 10^5 - 1 \times 10^9$) of lysed *S. cerevisiae* BY4737 cells or total RNA extract of the cells together with 1×10^{11} *in vitro* transcribed SUC2 mRNA target molecules. BY4743 yeast cells contained undetectable levels of SUC2 mRNA when grown on glucose (data not shown). No inhibition was observed when 10^6 or less lysed yeast cells or their RNA content were added to the hybridization (Fig. 4). 40 – 55% decrease in the signal of SUC2 mRNA was observed when 10^7 to 10^9 lysed cells were added to the hybridization solution. With extracted RNA the decrease was respectively 15 – 40%.

The inhibition effect of cell lysate on the hybridization was taken in to consideration, when the signal of target

RNA in sandwich hybridization was quantified in growth experiments (see Fig. 5 and 6).

Monitoring of 18S rRNA and SUC2 mRNA in growth experiments with *S. cerevisiae carlsbergensis*

To evaluate the developed sandwich hybridization assay with respect to its application in monitoring of different RNA species in bioprocesses, 18S rRNA and SUC2 mRNA were analyzed at different growth phases during shake flask cultivations of *S. cerevisiae carlsbergensis*. 18S rRNA was used as target for method evaluation due to its high cellular amount and stability. mRNA molecules represent only approximately 5% of total RNA in yeast [24] and thus are suitable targets for sensitivity evaluation. Expression of SUC2 mRNA expression can additionally be easily controlled by glucose repression [10].

Fig. 5 shows the level of 18S rRNA detected by the solution based sandwich hybridization assay in relation to the optical density and the specific growth rate of the yeast. The amount of 18S rRNA molecules per cell was calculated on the basis of a standard curve created with *in vitro* transcribed 18S rRNA. The hybridisation efficiencies of probes to native 18S rRNA and *in vitro* transcribed 18S rRNA were assumed to be the same. The assay was performed with a low number of $1 \times 10^6 - 3 \times 10^6$ lysed yeast




Figure 3
Comparison of extracted total RNA and crude cell lysate as sample material in the sandwich hybridization assay. 18S rRNA was used as target molecule for method evaluation. 5 ml of $OD_{600} = 2$ yeast culture was lysed (○) or used for extraction of total RNA (●). Different dilutions of lysed yeast cells or total RNA extracted from the same number of cells were added to the sandwich hybridization solution. Fluorescence (FLU) measured with 18S rRNA probes is presented in relation to the amount yeast cells used for lysis or extraction of RNA. The error bars show the \pm SD of three parallel experiments.




Figure 4
Effect of cell lysate and total RNA on the signal of the solution hybridization. 1.5×10^{11} SUC2 *in vitro* transcribed RNA molecules were added to the hybridization solution containing crude cell lysate (○) or RNA extract (●) from $1 \times 10^6 - 1 \times 10^9$ *S. cerevisiae* cells grown on glucose. The results are expressed as a percentage value of the signal compared to that obtained with the same amount of pure *in vitro* transcribed target SUC2 mRNA. Approximately 1.4 μ g total RNA was extracted from 1×10^8 cells. The error bars show the \pm SD of three parallel samples.

cells. Therefore the inhibitory effect of the cell extract on the assay was negligible (see Fig. 4).

During the first 5 hours the 18S rRNA level was slightly increasing and after that a nearly constant decrease was observed until 60 hours of growth. These results are in good correlation with the determined growth rate μ at the examined sampling points. Due to the fact that the preculture used for inoculation was in exponential growth phase, the increase of the 18S rRNA during the initial lag phase is relatively low.

Invertase activity is known to be under glucose repression and the corresponding SUC2 gene expression is induced by sucrose [10].

In Fig. 6 the level of SUC2 mRNA and the invertase enzyme activity were detected in cultures of *S. cerevisiae carlsbergensis* grown in shake flask cultures. In the two experiments the cultivation medium contained either sucrose or glucose as carbon source. For the calculation of the level of SUC2 mRNA the hybridization assay was performed with a cell extract containing $7 \times 10^7 - 1.2 \times 10^8$ lysed cells. In this case the inhibition effect of the




Figure 5
Level of 18S rRNA of *S. cerevisiae carlsbergensis* determined by sandwich hybridization as molecules per cell (●) in different growth states. The growth was followed by measuring the optical density at 600 nm (○) and the specific growth rate μ (□) was determined. The yeast was cultured in medium containing 1% yeast extract, 2% Bacto peptone solution and 2% glucose as carbon source. $1 \times 10^6 - 3 \times 10^6$ yeast cells as crude lysate was added to sandwich hybridization for 18S rRNA analysis. $OD_{600} = 1$ corresponds to 5.5×10^8 yeast cells ml^{-1} . The error bars of show the \pm SD of three parallel samples.




Figure 6

Level of SUC2 mRNA of *S. cerevisiae carlsbergensis* given as molecular amount per cell determined by sandwich hybridization (●) and the activity of invertase (▲) during exponential growth in medium containing 1% yeast extract, 2% Bacto peptone solution and either **a)** 2% sucrose or **b)** 2% glucose as carbon source. Growth was followed by optical density at 600 nm (□). $7 \times 10^7 - 1.2 \times 10^8$ yeast cells as crude lysate were added to the sandwich hybridization for 18S rRNA analysis. The error bars show the \pm SD of three parallel samples.

biological material on the assay had to be considered according to Fig. 4 and the signals were accordingly corrected by 50 – 55% in dependence on the specific amounts of extract used.

During the initial growth phase the invertase activity was about 200 to 300 U μg^{-1} fresh cells in sucrose grown culture but slightly lower (100 – 200 U μg^{-1}) in glucose grown culture. In contrast the level of SUC2 mRNA was 5 – 10 times higher in the sucrose culture than in the glucose containing culture. After 24 h of cultivation the total activity of invertase was significantly higher in the sucrose grown culture than in the glucose grown culture (800 to 1100 U μg^{-1} versus 450 U μg^{-1}). Correspondingly there was an approximately four times higher amount of SUC2 mRNA detected in the sucrose grown culture than in the glucose containing culture (Fig. 6).

Discussion

The developed assay is based on hybridization of RNA target molecules with a biotin labeled capture probe and a digoxigenin labeled detection probe. Alkaline phos-

phatase enzyme is attached to the detection probe after hybridization, which amplifies a fluorescence signal used for quantification. The indirect enzyme labeling system was able to detect 10^9 RNA target molecules. This is comparable to the sensitivities reached by others with similar detection systems used for PCR product quantification [16,23]. With direct probe labeling by thermostable alkaline phosphatase 100-fold higher sensitivities have been reached in other studies [12,26]. Although the direct labeling is more sensitive, the indirect labeling method of the probes is beneficial if the aim is to measure a large number of different RNA targets in one microwell plate. Each detection probe does not have to be labeled separately with an amplifying enzyme, which decreases the storage time of the probes.

The aim here was to apply the developed method for RNA expression monitoring of fermentation processes. Recently, this was performed by Jürgen *et al.* [13] and Schweder *et al.* [17] by slot blot analysis, which is in general time consuming and laborious. Here we present a simpler and faster method, with the possibility to use crude cell lysates instead of extracted RNA as a major improvement. We found up to 60% higher signal intensities with crude lysates when less than 3×10^7 lysed yeast cells were used per hybridization, possibly because a loss of target RNA during extraction procedure can be avoided. Carlson *et al.* [3] have reported approximately 30% loss of RNA with affinity resin based extractions.

Avoiding this by use of lysates provides a great benefit compared to RT-PCR and array based methods, which always require a separate RNA purification step. Use of high concentrations of either cell lysate or total RNA caused a decrease of the fluorescence signal possibly due to unspecific binding. With low cellular levels of target RNA more cell material is required in the assay, and correspondingly, then inhibition has to be considered in signal quantification [20]. Thus the amount of cell lysate used in hybridization assay should be optimized for routine analysis.

The sensitivity of the used sandwich hybridization assay was determined to be approximately 2 fmol, which was similar to the membrane based slot blot hybridization assay. Nevertheless, the linear range of the solution hybridization was approximately 20-fold higher compared to membrane hybridization. The time required for the sandwich assay from sampling to detection was about three hours, which could be further shortened. In contrast, the membrane based hybridization and detection could not be completed within less than 12 hours. The main difference of the two methods is the time it takes for the hybridization to reach equilibrium. Real-time PCR takes about 5 hours to complete and was in our investigation the most sensitive method (0.6 amol), but this

method suffers from the demand of a very pure RNA preparation in order to avoid inhibition of the reverse transcriptase and DNA polymerase.

The described method was evaluated in shake flask experiments, where 18S rRNA and SUC2 mRNA levels were measured during different growth phases. rRNA can be used as an indicator of cellular activity or growth state [9]. Results showed that the changes in 18S rRNA expression level were simultaneous with the changes in the specific growth rate μ during different growth phases of the yeast *S. cerevisiae carlsbergensis*.

SUC2 mRNA levels were in good correlation with the measured invertase enzyme activities. Cultivation of *S. cerevisiae carlsbergensis* in either glucose or sucrose containing medium resulted in different levels of SUC mRNA and invertase activity. Our experiments confirm the knowledge that brewery strains are less sensitive to glucose repression than laboratory strains and are thus often more adapted to disaccharide utilization [8]. Whereas we found expression of SUC2 mRNA in *S. cerevisiae carlsbergensis* glucose-grown cultures, no expression of SUC2 mRNA was found in corresponding cultures of the laboratory strain *S. cerevisiae* BY4737 (data not shown).

The molecular amount of RNA per cell was calculated using *in vitro* transcribed 18S rRNA or SUC2 mRNA as standards. We determined 25 000 – 140 000 18S rRNA molecules per cell in dependence on the different growth phases. These values are in good agreement with results presented by Warner, who calculated about 200 000 ribosomes in exponentially growing yeast cells [24]. For highly expressed genes 200 – 400 mRNA copies per cell have been determined with SAGE analysis [22]. In our current study the SUC2 mRNA level has been calculated in the same range. The number of mRNA molecules varied between 100 – 400 molecules per cell when invertase formation was induced. Quantification of the *in vitro* standard RNA, sample mRNA stability, and the effect of the biological material on the hybridization efficiency can cause inaccuracies to the quantification of the target mRNA species, but our method seems to provide a fair estimation of the molecular levels of RNA.

The sensitivity of the developed method was found to be sufficient for detection of intracellular rRNA molecules and highly expressed mRNA species. For RNA molecules with a low expression level a higher sensitivity would be desirable if crude cell lysates are used as sample material for the analysis, to avoid the use of cell concentrations which inhibit the analysis. Although for physiological studies the number of target RNA molecules of low expressed genes can be increased by increasing the amount

of cell extract, signal inhibition has then to be included into the calculations.

Automation of the sandwich hybridization and technical improvements e.g. detection of targets with electric biochips [21] are making the sandwich hybridization method interesting for many application areas especially for clinical diagnostics, food industry use and even for the detection of bioweapons [18]. One of the challenging applications is the state analysis of bioprocesses based on transcriptional profiling of key genes that reflect the physiological state of the producing organism. The here presented method in view of its automation capabilities offers a valuable tool in this direction.

Conclusions

Alternative and novel methods for RNA analysis are being developed for clinical and industrial diagnostics as well as for studying the physiology of living cells. The aim of the present study was the development of a fast and reliable method for quantification of single RNA species in solution using a sandwich hybridization based assay applicable for physiology analysis of organisms used in bioprocesses. The presented method allows the use of crude cell lysates as sample material and offers a sensitivity level sufficient for measurement of ribosomal RNAs and messenger RNAs with relatively high expression levels. The developed method was evaluated by measuring the levels of *Saccharomyces cerevisiae carlsbergensis* 18S rRNA and SUC2 mRNA in shake flask experiments. The measured 18S rRNA level was consistent with the specific growth rate and thus could be used as an indicator of growth state of a cell. SUC2 mRNA was in accordance with measured invertase activities under glucose repression and sucrose induction.

Methods

Strains and cultivation

S. cerevisiae strain BY4743 and the industrial *S. cerevisiae carlsbergensis* strain LK01 were used for cultivation. Cells were grown in medium containing 1% yeast extract (Difco), 2% Bacto Peptone (Difco) with 2% glucose or sucrose as carbon source at 30°C and 200 rpm in baffled shake flasks. For the culture used in analysis 20 ml of a 24 h grown preculture was inoculated in a 1 l flask containing 200 ml fresh medium. Growth was followed spectrophotometrically by measuring the optical density at 600 nm. The culture was grown for 60 h, during which yeast samples were taken for analysis. For RNA analysis 5–10 ml samples were taken depending on the growth state. The cell number in relation to OD was determined by Thoma cell counting. Cells were centrifuged at 20800 × g for 10 min at 4°C, washed with cold DEPC water and frozen at -70°C before analysis.

Table 1: Sequences of the oligonucleotide probes used in sandwich and slot blot hybridizations. All probes are designed for *S. cerevisiae*.

Target name	Probe name	Probe sequence 5'→3'	Location of probe
	Capture probes		
YLI162W	SUC2-1400	UUGUAGUCUUGUGAUUUGGCAGC- (AAGAG)*	1400–1422
RDN18-2	18S rRNA-400	CUGCUGCCUUCUUGGAUGUGGUAGCCGUU- (GAGAG)	400–429
	Detection probes		
YLI162W	SUC2-581	GACUUGAAUGGUUGGUUGUUGAC- (GUGAC)	581–603
RDN18-2	18S rRNA-1302	UCUCGUUCGUUAUCGCAAUAAGCAGACAA- (GCAGC)	1302–1331

* 5nt long spacer arms in brack

Yeast cell lysis and RNA extraction

Frozen yeast cells were suspended to 250 – 500 mg fresh yeast ml⁻¹ with lysis buffer containing 550 U ml⁻¹ Lyticase (Sigma), 1 × SSC (0.15 M sodium chloride, 0.015 M sodium citrate, pH 7.0) and 0.2% RNA guard RNase inhibitor (Amersham Pharmacia). After addition of 100 µl of acid-washed beads (diameter 425–600 µm, Sigma) cells were disrupted with a FastPrep cell homogenizer (ThermoSavant) using 6 m sec⁻¹ for 45 sec. Lysed cells and beads were rinsed with 100 µl DEPC-treated water. The cell lysate was directly used for RNA analysis.

Total RNA from freshly grown yeast cells was extracted using the TOTAL RNA kit (A&A Biotechnology, Poland) following the protocol provided with the kit. A maximum of 50 mg of cells was used per extraction to not exceed the binding capacity of the silica columns. Extracted RNA was quantified by absorbance measurement at A₂₆₀ (DNA Quant, Pharmacia).

Synthesis of external control RNA

RNA used as an external control for the sandwich hybridization was synthesized *in vitro* by T7-RNA polymerase from the respective PCR products containing a T7 promoter sequence using the DIG RNA labeling Kit (Roche Diagnostics) as recommended by the manufacturer but with unlabeled nucleotides. The following primers were used for synthesis of the appropriate PCR product:

5'CCACATCTCCATCGTTGAAG and 5'CTAATACGACT-CACTATAGGGAGATTCC-TTTTCCTTTTGGCTGG for SUC2

5'TCCTCTAAATGACCAAGTTTG and 5'CTAATACGACT-CACTATAGGGAGATGGA-AGAGATGTATTTATTAGAT for 18S rRNA

Synthesized RNA was quantified by RiboGreen RNA Quantification Kit (Molecular Probes) as recommended by the manufacturer.

Synthesis of oligonucleotide probes

3' end biotin labeled capture probes, were custom-synthesized by Sigma Genosys. Detection probes were labeled with digoxigenin with oligonucleotide 3' end labeling Kit (Roche Diagnostics) (Table 1) as recommended by the manufacturer.

Sandwich hybridization assay

The total RNA extract, crude cell lysates or *in vitro* synthesized control RNA were diluted in buffer containing H₂O, 20 × SSC and deionized formamide in a ratio of 5:3:2 and denatured at 65°C for 3–5 min before addition to hybridization reaction with 3 × 10¹² molecules of biotin labeled capture probe and 2.4 × 10¹² molecules of DIG-labeled detection probe. The hybridization reactions were carried out in 96-well plates (Greiner) at 50°C for 30 min at 600 rpm shaking (Thermomixer Comfort, Eppendorf) in a total volume of 100 µl containing 5 × SSC, 0.5% SDS, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% BSA, 20% deionized formamide, 4% dextran sulfate.

The hybrid collection was performed by addition of 20 µg of streptavidin coated magnetic beads (Promega) to the hybridization reaction followed by incubation at 37°C for 30 min at 600 rpm. The beads were washed three times with 1 × SSC buffer and afterwards 50 µl of conjugation buffer (50 mM Tris pH 7.5, 150 mM NaCl, 0.3% Tween 20, 1% BSA) containing 35 mU anti-DIG-alkaline phosphatase FAB fragments (Roche Diagnostics) were added to the beads and incubated for 30 min at room temperature. After six washing steps with washing buffer (50 mM Tris pH 7.5, 150 mM NaCl, 0.3% Tween 20), 150 µl of AttoPhos substrate (Promega) were added to the beads and incubated for 20 min at 37°C. The magnetic beads were separated between the washing steps with a MagnaBot 96 Magnetic Separation Device (Promega). The clear supernatants were transferred to a black 96-well plate (Greiner) for fluorescence measurement performed with Victor 2 fluorescence reader (Wallac) at excitation wavelength 430 nm and emission wavelength 560 nm.

Slot Blot hybridization

For detection of yeast 18S rRNA by slot blot hybridization the 3' end DIG-labeled probe 18SrRNA-1302 (table 1) was used. After denaturation in buffer containing H₂O, 20 × SSC and formaldehyde in the ratio of 5:3:2 at 65 °C for 10 min dilution series of *in vitro* transcribed 18S rRNA were blotted onto positively charged nylon membranes (Amersham Pharmacia) using Bio Dot vacuum (BioRad). 1 h prehybridization and over night hybridization were carried out at 50 °C using High SDS hybridization buffer (7% SDS, 50% formamide, 5 × SSC, 2% Blocking Reagent (Roche Diagnostics), 50 mM sodium phosphate, 0.1% N-lauroylsarcosine, pH 7.0). Signal detection was performed by chemiluminescence using CDP Star (Roche Diagnostics) as a substrate as recommended by the manufacturer and exposure to chemiluminescence films (Hyperfilm ECL, Pharmacia) for 60 min at room temperature. The film sheets were scanned by use of a flatbed scanner (Epson) and relative amounts of detected 18S rRNA were calculated by volume densitometric analysis using the Quantity One[®] Quantitation Software from BioRad.

Real-time PCR

Real-time PCR was carried out on an ABI Prism 7700 (Perkin-Elmer-Applied Biosystems, Foster City, CA, USA), using SYBR-green as detection dye. cDNA synthesis of *in vitro* transcribed 18S rRNA was performed with Enhanced Avian RT first Strand Synthesis kit (Sigma), using the antisense specific primer: 18S-741R; 5'CTTGGCAAATGCTTTCGCAG. The PCR reaction was done with SYBR Green JumpStart Taq ReadyMix (Sigma), using the primers: 18S-523F; 5'CCTTGTGGCTCTTGGCGAAC and 18S-741R.

Invertase activity measurement

Invertase activity was measured in cell lysates in a two-phase reaction. In the first phase invertase hydrolyses sucrose to glucose and fructose. In the second phase glucose and ATP form glucose-6-phosphate and ADP in a reaction catalyzed by hexokinase. Glucose-6-phosphate and NADP form 6-phosphogluconate and NADPH in a glucose-6-phosphate dehydrogenase catalyzed reaction. NADPH is measured at 340 nm.

Reaction was started by adding lysed cells to a concentration of 0.6 mg ml⁻¹ in 50 mM sodium acetate (pH 4.5) and 20 mM sucrose at 30 °C with a final volume of 1 ml. Reaction was stopped after 10 min by boiling. 50 µl of the reaction solution were added to 950 µl of 50 mM Hepes (pH 8.0), 5 mM MgCl₂, 3 mM ATP and 1 mM NADP. A₃₄₀ was measured spectrophotometrically (Ultrospec 2100 pro, Amersham Pharmacia) before addition of 4.5 U hexokinase (Sigma) and 1.2 U glucose-6-phosphate dehydrogenase (Sigma) and again after 30 min reaction at room temperature.

Author's contribution

JR designed the experiments and carried them out for the method development and growth experiments. KBB performed the experiments with real-time PCR. KBB and JL participated in the method development and optimisation work. SM contributed with ideas and discussions in the writing of the manuscript. AB and PN participated in the design and supervision of the work. All authors read and approved the final manuscript.

Acknowledgements

These studies were supported by the European RTD project QLRT-1999-00533, the SUNARE program of the Academy of Finland (project no. 52796) and personal grants to Jari Rautio by the Graduate School in Chemical Engineering (Finland) and Biocenter Oulu (Finland).

References

- Albretsen C, Kalland KH, Haukanes BI, Havarstein LS and Kleppe K **Applications of magnetic beads with covalently attached oligonucleotides in hybridization: isolation and detection of specific measles virus mRNA from a crude cell lysate.** *Anal Biochem* 1990, **189**:40-50
- Bustin SA **Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays.** *J Mol Endocrinol* 2000, **25**:169-193
- Carlsson B, Jernas M, Lindell K and Carlsson LM **Total RNA and array-based expression monitoring.** *Nat Biotechnol* 2000, **18**:579
- Casademont I, Bizet C, Chevrier D and Guesdon JL **Rapid detection of *Campylobacter fetus* by polymerase chain reaction combined with non-radioactive hybridization using an oligonucleotide covalently bound to microwells.** *Mol Cell Probes* 2000, **14**:233-240
- Casademont I, Chevrier D, Denoyelle F, Petit C and Guesdon JL **A simple and reliable method for the detection of the 30delG mutation of the CX26 gene.** *Mol Cell Probes* 2000, **14**:149-152
- Chevrier D, Oprisan G, Maresca A, Matsiota-Bernard P and Guesdon JL **Isolation of a specific DNA fragment and development of a PCR-based method for the detection of *Mycobacterium genavense*.** *FEMS Immunol Med Microbiol* 1999, **23**:243-252
- Chevrier D, Popoff MY, Dion MP, Hermant D and Guesdon JL **Rapid detection of *Salmonella* subspecies I by PCR combined with non-radioactive hybridisation using covalently immobilised oligonucleotide on a microplate.**
- Crumplén RM, Slaughter JC and Stewart GG **Characteristics of maltose transporter activity in an ale and large strain of the yeast *Saccharomyces cerevisiae*.** *Lett Appl Microbiol* 1996, **23**:448-452
- DeLong EF, Wickham GS and Pace NR **Phylogenetic stains: ribosomal RNA-based probes for the identification of single cells.** *Science* 1989, **243**:1360-1363
- Gancedo JM **Yeast carbon catabolite repression.** *Microbiol Mol Biol Rev* 1998, **62**:334-361
- Iezzoni JC, Kang JH, Montone KT, Reed JA and Brigati DJ **Colorimetric detection of herpes simplex virus by DNA in situ sandwich hybridization: a rapid, formamide-free, random oligomer-enhanced method.** *Nucleic Acids Res* 1992, **20**:1149-1150
- Ishii JK and Ghosh SS **Bead-based sandwich hybridization characteristics of oligonucleotide-alkaline phosphatase conjugates and their potential for quantitating target RNA sequences.** *Bioconjug Chem* 1993, **4**:34-41
- Jurgen B, Lin HY, Riemschneider S, Scharf C, Neubauer P, Schmid R, Hecker M and Schweder T **Monitoring of genes that respond to overproduction of an insoluble recombinant protein in *Escherichia coli* glucose-limited fed-batch fermentations.** *Biotechnol Bioeng* 2000, **70**:217-224
- Kwoh DY, Davis GR, Whitfield KM, Chappelle HL, DiMichele LJ and Gingeras TR **Transcription-based amplification system and detection of amplified human immunodeficiency virus type I with a bead-based sandwich hybridization format.** *Proc Natl Acad Sci U S A* 1989, **86**:1173-1177

15. Patra G, Sylvestre P, Ramisse V, Therasse J and Guesdon JL **Isolation of a specific chromosomal DNA sequence of *Bacillus anthracis* and its possible use in diagnosis.** *FEMS Immunol Med Microbiol* 1996, **15**:223-231
16. Satokari R, Juvonen R, Mallison K, von Wright A and Haikara A **Detection of beer spoilage bacteria *Megasphaera* and *Pectinatus* by polymerase chain reaction and colorimetric microplate hybridization.** *Int J Food Microbiol* 1998, **45**:119-127
17. Schweder T, Lin HY, Jurgen B, Breitenstein A, Riemschneider S, Khalameyzer V, Gupta A, Buttner K and Neubauer P **Role of the general stress response during strong overexpression of a heterologous gene in *Escherichia coli*.** *Appl Microbiol Biotechnol* 2002, **58**:330-337
18. Service RF **Analytical chemistry. New test could speed bio-weapon detection.** *Science* 2002, **295**:1447
19. Sjoroos M, Ilonen J, Reijonen H and Lovgren T **Time-resolved fluorometry based sandwich hybridisation assay for HLA-DQA1 typing.** *Dis Markers* 1998, **14**:9-19
20. Tenhunen J, Eloranta J, Kallio A and Soderlund H **A solution hybridization method for quantification of mRNAs: determining the amount and stability of oncogene mRNA.** *Genet Anal Tech Appl* 1990, **7**:228-233
21. Umek RM, Lin SW, Vielmetter J, Terbruggen RH, Irvine B, Yu CJ, Kayyem JF, Yowanto H, Blackburn GF, Farkas DH and Chen YP **Electronic detection of nucleic acids: a versatile platform for molecular diagnostics.** *J Mol Diagn* 2001, **3**:74-84
22. Velculescu VE, Zhang L, Zhou W, Vogelstein J, Basrai MA, Bassett DE Jr, Hieter P, Vogelstein B and Kinzler KW **Characterization of the yeast transcriptome.** *Cell* 1997, **88**:243-251
23. Vesanen M, Piiparinen H, Kallio A and Vaheeri A **Detection of herpes simplex virus DNA in cerebrospinal fluid samples using the polymerase chain reaction and microplate hybridization.** *J Virol Methods* 1996, **59**:1-11
24. Warner JR **The economics of ribosome biosynthesis in yeast.** *Trends Biochem Sci* 1999, **24**:437-440
25. Whetsell AJ, Drew JB, Milman G, Hoff R, Dragon EA, Adler K, Hui J, Otto P, Gupta P and Farzadegan H **Comparison of three nonradioisotopic polymerase chain reaction-based methods for detection of human immunodeficiency virus type 1.** *J Clin Microbiol* 1992, **30**:845-853
26. Wicks B, Cook DB, Barer MR, O'Donnell AG and Self CH **A sandwich hybridization assay employing enzyme amplification for termination of specific ribosomal RNA from unpurified cell lysates.** *Anal Biochem* 1998, **259**:258-264

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

