GENOME SEQUENCES



AMERICAN SOCIETY FOR MICROBIOLOGY

Draft Genome Sequence of *Psychrobacter okhotskensis* Strain 5179-1A, Isolated from a Raw Cured Ham Storage Crate

🖻 Joseph Wambui,ª Marina Morach,ª Nicole Cernela,ª Marc J. A. Stevens,ª Giovanni Ghielmetti,ª 🖻 Roger Stephanª

alnstitute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland

ABSTRACT We present the draft genome sequence of *Psychrobacter okhotskensis* strain 5179-1A, which was isolated from a raw cured ham storage crate. Its size and GC content are 3.4 Mb and 43.4%, respectively. The 16S rRNA sequences of strain 5179-1A and *P. okhotskensis* MD17^T are 100% identical.

Psychrobacter okhotskensis is a facultatively psychrophilic Gram-negative, catalaseand oxidase-positive, nonmotile, aerobic coccobacillus (1). To the best of our knowledge, *P. okhotskensis* has so far been isolated only from permafrost and the marine environment (1–3), and there is no publicly available genome. Here, we have determined the draft genome sequence of *P. okhotskensis* 5179-1A.

An inoculum, prepared from a swab sample taken from a crate that was used for storing raw cured ham in Zurich, Switzerland, was streaked onto Columbia blood agar supplemented with 5% defibrinated sheep blood and incubated aerobically at 30°C for 3 days. This strain was purified by subculturing on the same medium under similar growth conditions. Genomic DNA was isolated from the pure culture using the DNA blood and tissue kit (Qiagen, Hombrechtikon, Switzerland). The DNA was prepared using a Nextera DNA Flex sample preparation kit (Illumina, San Diego, CA, USA), and the resulting transposome-based libraries were sequenced on a MiniSeq sequencer (Illumina). The sequencing output was 254 Mb of 150-bp paired-end reads. Reads were checked for quality using the software package FastQC v0.11.7 (4) and then assembled using the SPAdes v3.0 (5)-based software Shovill v1.0.4 (https://github.com/tseemann/ shovill). The assembly was filtered, retaining contigs of >500 bp. The genome was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (6) and the RAST pipeline (7). Default parameters were used for all software and Web servers. The sequenced genome was assembled into 72 contigs and comprises 3.4 Mb (GC content, 43.4%) with 2,982 predicted protein-coding sequences. The genome coverage, contig N_{50} , and contig L_{50} were 80.0×, 111,840 bp, and 10, respectively.

RASTtk (7) and tRNAscan-SE (8) identified 47 RNAs and 43 tRNAs, respectively. Species identification was carried out using the 16S rRNA gene-based identification server (9). The strain's 1,539-bp 16S rRNA sequence was 100% similar, with a coverage of 100%, to that of *P. okhotskensis* MD17^T. The GC content of *P. okhotskensis* 5179-1A is lower than that of *P. okhotskensis* MD17^T, which is 46.7% (1). RASTtk (7) identified 24 features for fluoroquinolone, β -lactam, cadmium, copper, zinc, and cobalt resistance or tolerance. VFDB 2019 (10) predicted 22 potential virulence factors, including hemolysin and a siderophore orthologous to Baumannii acinetobactin utilization A protein (BauA). The probability of *P. okhotskensis* being a human pathogen was predicted to be 0.54 by PathogenFinder (11); 10 pathogenic families were matched from *Psychrobacter arcticus* 273-4, which corresponds with previous reports that *Psychrobacter* spp. can be considered opportunistic pathogens in humans and livestock (12–15).

Data availability. This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. JABRVQ000000000. The version described

Citation Wambui J, Morach M, Cernela N, Stevens MJA, Ghielmetti G, Stephan R. 2020. Draft genome sequence of *Psychrobacter okhotskensis* strain 5179-1A, isolated from a raw cured ham storage crate. Microbiol Resour Announc 9:e00682-20. https://doi.org/10.1128/ MRA.00682-20.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2020 Wambui et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Joseph Wambui, joseph.wambui@uzh.ch.

Received 12 June 2020 Accepted 24 June 2020 Published 16 July 2020 in this paper is version JABRVQ010000000. The raw sequencing reads have been deposited in the SRA under the accession no. SRP265202.

REFERENCES

- Yumoto I, Hirota K, Sogabe Y, Nodasaka Y, Yokota Y, Hoshino T. 2003. *Psychrobacter okhotskensis* sp. nov., a lipase-producing facultative psychrophile isolated from the coast of the Okhotsk Sea. Int J Syst Evol Microbiol 53:1985–1989. https://doi.org/10.1099/ijs.0.02686-0.
- Yu Y, Li H, Zeng Y, Chen B. 2010. Phylogenetic diversity of culturable bacteria from Antarctic sandy intertidal sediments. Polar Biol 33: 869–875. https://doi.org/10.1007/s00300-009-0758-3.
- Pacova Z, Urbanova E, Durnova E. 2001. *Psychrobacter immobilis* isolated from foods: characteristics and identification. Vet Med (Praha) 46: 95–100. https://doi.org/10.17221/7866-VETMED.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. https://www.bioinformatics.babraham.ac.uk/projects/ fastqc.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. https://doi.org/10.1089/cmb.2012.0021.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity GM, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. Omics 12:137–141. https://doi.org/10.1089/omi.2008.0017.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. Sci Rep 5:8365. https:// doi.org/10.1038/srep08365.

- Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. Nucleic Acids Res 44: W54–W57. https://doi.org/10.1093/nar/gkw413.
- Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. Int J Syst Evol Microbiol 67: 1613–1617. https://doi.org/10.1099/ijsem.0.001755.
- Liu B, Zheng D, Jin Q, Chen L, Yang J. 2019. VFDB 2019: a comparative pathogenomic platform with an interactive Web interface. Nucleic Acids Res 47:D687–D692. https://doi.org/10.1093/nar/gky1080.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother 67:2640–2644. https://doi .org/10.1093/jac/dks261.
- Stepanović S, Vuković D, Bedora-Faure M, K'ouas G, Djukić S, Svabić-Vlahović M, Carlier J-P. 2007. Surgical wound infection associated with *Psychrobacter phenylpyruvicus*-like organism. Diagn Microbiol Infect Dis 57:217–219. https://doi.org/10.1016/j.diagmicrobio.2006.08.002.
- Le Guern R, Wallet F, Vega E, Courcol RJ, Loïez C. 2014. Psychrobacter sanguinis: an unusual bacterium for nosocomial meningitis. J Clin Microbiol 52:3475–3477. https://doi.org/10.1128/JCM.01197-14.
- Caspar Y, Recule C, Pouzol P, Lafeuillade B, Mallaret MR, Maurin M, Croize J. 2013. *Psychrobacter arenosus* bacteremia after blood transfusion, France. Emerg Infect Dis 19:1118–1120. https://doi.org/10.3201/eid1907 .121599.
- Vela AI, Collins MD, Latre MV, Mateos A, Moreno MA, Hutson R, Domínguez L, Fernández-Garayzábal JF. 2003. *Psychrobacter pulmonis* sp. nov., isolated from the lungs of lambs. Int J Syst Evol Microbiol 53:415–419. https://doi.org/10.1099/ijs.0.02413-0.