

Research article

Open Access

PPAR α L162V underlies variation in serum triglycerides and subcutaneous fat volume in young males

Julieta Uthurralt¹, Heather Gordish-Dressman¹, Meg Bradbury¹, Carolina Tesi-Rocha¹, Joseph Devaney¹, Brennan Harmon¹, Erica K Reeves¹, Cinzia Brandoli¹, Barbara C Hansen², Richard L Seip³, Paul D Thompson³, Thomas B Price^{3,4}, Theodore J Angelopoulos⁵, Priscilla M Clarkson⁶, Niall M Moyna⁷, Linda S Pescatello⁸, Paul S Visich⁹, Robert F Zoeller¹⁰, Paul M Gordon¹¹ and Eric P Hoffman*¹

Address: ¹Research Center for Genetic Medicine, Children's National Medical Center, Washington DC, 20010, USA, ²Obesity, Diabetes and Aging Research Center, College of Medicine, University of South Florida, 12901 Bruce B. Downs Blvd, Tampa, Florida 33612, USA, ³Division of Cardiology, Henry Low Heart Center, Hartford Hospital, Hartford, CT, 06102, USA, ⁴Department of Diagnostic Radiology, Yale University School of Medicine, New Haven, CT 06520, USA, ⁵Center for Lifestyle Medicine and Department of Health Professions, University of Central Florida, Orlando FL 32816, USA, ⁶Department of Exercise Science, Totman Building, University of Massachusetts, Amherst, MA, 01003, USA, ⁷Department of Sport Science and Health, Dublin City University, Dublin 9, Ireland, ⁸School of Allied Health, University of Connecticut, Storrs, CT 06269, USA, ⁹Human Performance Laboratory, Central Michigan University, Mount Pleasant, Mich. 48859, USA, ¹⁰Department of Exercise Science and Health Promotion, Florida Atlantic University, Davie, FL 33314, USA and ¹¹Division of Exercise Physiology, School of Medicine, West Virginia University, Morgantown WV 26506, USA

Email: Julieta Uthurralt - juthurralt@cnmcresearch.org; Heather Gordish-Dressman - hgordish@cnmcresearch.org; Meg Bradbury - mbradbury@cnmcresearch.org; Carolina Tesi-Rocha - ctesirocha@cnmcresearch.org; Joseph Devaney - jdevaney@cnmcresearch.org; Brennan Harmon - bharmon@cnmcresearch.org; Erica K Reeves - ereeves@cnmcresearch.org; Cinzia Brandoli - cbrandoli@cnmcresearch.org; Barbara C Hansen - bchansen@aol.com; Richard L Seip - rseip@harthosp.org; Paul D Thompson - pthomps@harthosp.org; Thomas B Price - Thomas.price@yale.edu; Theodore J Angelopoulos - tangelop@mail.ucf.edu; Priscilla M Clarkson - clarkson@excsci.umass.edu; Niall M Moyna - niall.moyna@dcu.ie; Linda S Pescatello - linda.pescatello@uconn.edu; Paul S Visich - paul.visich@cmich.edu; Robert F Zoeller - rzoeller@fau.edu; Paul M Gordon - pgordon@wvu.edu; Eric P Hoffman* - ehoffman@cnmcresearch.org

* Corresponding author

Published: 16 August 2007

Received: 14 February 2007

BMC Medical Genetics 2007, 8:55 doi:10.1186/1471-2350-8-55

Accepted: 16 August 2007

This article is available from: <http://www.biomedcentral.com/1471-2350/8/55>

© 2007 Uthurralt et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Of the five sub-phenotypes defining metabolic syndrome, all are known to have strong genetic components (typically 50–80% of population variation). Studies defining genetic predispositions have typically focused on older populations with metabolic syndrome and/or type 2 diabetes. We hypothesized that the study of younger populations would mitigate many confounding variables, and allow us to better define genetic predisposition loci for metabolic syndrome.

Methods: We studied 610 young adult volunteers (average age 24 yrs) for metabolic syndrome markers, and volumetric MRI of upper arm muscle, bone, and fat pre- and post-unilateral resistance training.

Results: We found the PPAR α L162V polymorphism to be a strong determinant of serum triglyceride levels in young White males, where carriers of the V allele showed 78% increase in

triglycerides relative to L homozygotes (LL = 116 ± 11 mg/dL, LV = 208 ± 30 mg/dL; $p = 0.004$). Men with the V allele showed lower HDL (LL = 42 ± 1 mg/dL, LV = 34 ± 2 mg/dL; $p = 0.001$), but women did not. Subcutaneous fat volume was higher in males carrying the V allele, however, exercise training increased fat volume of the untrained arm in V carriers, while LL genotypes significantly decreased in fat volume (LL = $-1,707 \pm 21$ mm³, LV = $17,617 \pm 58$ mm³; $p = 0.002$), indicating a systemic effect of the V allele on adiposity after unilateral training. Our study suggests that the primary effect of PPAR α L162V is on serum triglycerides, with downstream effects on adiposity and response to training.

Conclusion: Our results on association of PPAR α and triglycerides in males showed a much larger effect of the V allele than previously reported in older and less healthy populations. Specifically, we showed the V allele to increase triglycerides by 78% ($p = 0.004$), and this single polymorphism accounted for 3.8% of all variation in serum triglycerides in males ($p = 0.0037$).

Background

Metabolic syndrome is considered to be a pre-diabetic state, with abnormal values for triglycerides, HDL, adiposity, blood pressure, and insulin resistance [1]. Twin and family studies have demonstrated the prominent influence of genetic factors on metabolic syndrome sub-phenotypes [2,3]. To date, most research aimed to identify these genetic underpinnings have been done in elderly or diseased populations. Less progress has been made on the identification of initial risk factors such as lipid levels and insulin resistance in young individuals. With regards to these sub-phenotypes, it is known that the genetic factors in an aged population may be different than those in a younger population. For example, a longitudinal twin study has reported that only 40% of the genetic factors affecting BMI are shared at the age of 20 y and 40 y [4].

We have previously described a genetic association study of university-age young adults, with measurement of baseline strength and tissue volumes of the upper arm, then response to a 12 wk unilateral resistance training of the non-dominant arm [5-8]. We chose the non-dominant arm to minimize environmental influences (use), and thus maximize sensitivity for detecting genetic influences. We recently expanded the study to include baseline metabolic syndrome measures. In this study, we used upper arm subcutaneous fat as a surrogate marker for adiposity, although visceral fat is more highly studied as a measure of obesity. Here, our goal was to test the effect of the PPAR α gene polymorphism (L162V) on a young adult volunteer population, and to begin to better dissect the effects of PPAR α genotype on the earliest phenotypes.

Peroxisome-proliferator-receptor-alpha (PPAR α) is involved in adipocyte differentiation, lipid and lipoprotein metabolism. Studies in mice have shown that PPAR α -deficient animals were unable to metabolize lipids and develop late onset obesity even when kept on a stable diet [9,10]. PPAR α is activated by circulating fatty acids, and peroxisome proliferator receptor agonists such as hypoli-

pidemic drugs (fibrates), commercially used plasticizers, synthetic fatty acids, steroid hormones, herbicides and pesticides [11]. Activation of the receptor results in the increased expression of genes involved in lipid oxidation, lipoprotein metabolism, inhibition of vascular inflammation and adipocyte differentiation through binding to peroxisome proliferator response elements (PPRE) in the DNA sequence of target genes [12]. In summary the activation of PPAR α stimulates fatty acid transportation and oxidation primarily in liver and muscle reducing the storage of fat in adipocytes. Thus, PPAR α is a strong candidate gene for the genetic component of metabolic syndrome.

The PPAR α gene has been screened for common polymorphic variations. The most studied variant is a missense mutation in exon 5 that results in a nonconservative amino acid substitution, leucine 162 valine (L162V). The V allele is Leucine 162 is located within the DNA binding domain and is highly conserved among species [13,14]. Functional studies of the PPAR α L162V polymorphism have shown differences in the activity of the receptor depending on the ligand concentration. At low ligand concentrations the activity of the receptor was lower for the V isoform than the L isoform, whereas at high ligand concentrations the opposite occurred (V activity decreased relative to L) [15]. The L162V polymorphism has been previously associated with dyslipidemia [16-21], and some measures of adiposity [22,23]. However, some of these studies showed opposite findings or negative associations. Many factors may contribute to an inability to replicate genetic association studies. First, differences in the population studied with most published studies focusing on older or unhealthy populations. Second, different methods of measuring both regional and total adiposity are used, and each of these methods has well-documented issues concerning reliability of measure, sensitivity, and specificity for measuring fat.

The aim of this study was to identify the role that PPAR α L162V polymorphism may play in the variability of serum

lipid concentration and variations in subcutaneous fat volume and BMI in 610 young healthy volunteers (average age 24 years). Fat volume was determined using semi-automated three-dimensional reconstruction analysis software. We tested the PPAR α L162V polymorphism against our volumetric MRI measures and variations in serum lipid profiles. We found strong statistical support for sex-specific effects of PPAR α genotype with plasma triglyceride, HDL, and BMI. The effect of PPAR α genotype was much stronger in our young adult population than previously reported for older populations, suggesting that younger populations may afford greater sensitivity in identification of the genetic underpinnings of metabolic syndrome sub-phenotypes.

Methods

Study overview and subjects

The study population was derived from a multicenter, NIH funded study designed to identify genetic factors that dictate baseline bone, muscle and fat volume and the variability in response to exercise training. The study design protocol has been described in detail elsewhere [24], and preliminary reports of genetic associations with muscle strength and size have been reported [5-8]. Briefly, 1,300 men and women, average age 24 (range 18 – 40 yrs) were recruited by one of the 8 centers (University of Massachusetts Amherst, University of Connecticut, Dublin University (Ireland), Florida Atlantic University, Hartford Hospital, University of Central Florida, West Virginia University, Central Michigan University). Participants were excluded if they: were <18 y or >40 y; used medications known to affect skeletal muscle such as corticosteroids; had any restriction of activity; had chronic medical conditions such as diabetes; had metal implants in arms, eyes, head, brain, neck, or heart which would prohibit MRI testing; had performed strength training or employment requiring repetitive use of the arms within the prior 12 months; consumed on average >2 alcoholic drinks daily; or had used dietary supplements reported to build muscle size/strength or to cause weight gain such as protein supplements, creatine, or androgenic precursors. Subjects were asked to maintain their normal dietary intake for the duration of the study. Written informed consent was obtained from each volunteer. The study was approved by the Children's National medical Center Institutional Review Board (protocol #2449), and was in compliance with the Helsinki Declaration.

We limited this current study to 610 Caucasians that had values for entry metabolic syndrome markers, completed pre- and post-MRI measurements, or both (table 1).

Exercise Training Program

Resistant training was performed with the non-dominant arm. The protocol was described elsewhere [24]. Briefly it

consisted of two 45 – 60 minute sessions per week for 12 weeks. Each session was supervised by an exercise physiologist professional or a trained student. Before each session, participants warmed-up with 2 sets of 12 repetitions of the biceps preacher curl and the seated overhead triceps extension. Each session included dumbbell biceps curls, dumbbell biceps preacher curls, and incline dumbbell biceps curls, overhead dumbbell triceps extension, and dumbbell triceps kickbacks. The amount of weight was aggressively increased during the 12 wks.

Subject Phenotyping

Anthropometric assessment

Body weight was recorded before and after the exercise training using a balance beam scale. Height was measured using a tape mounted on a wall and recorded in inches. BMI was calculated from weight (kg) and height (m).

MRI assessment

Subjects were scanned in the supine position with arms at their sides and their palms up on the scanner bed surface. The arm maximum circumference was determined with the subject standing, with the shoulder abducted at 90 degrees, the hand supinated, and the biceps flexed. A vitamin E bead was placed on the front of the biceps portion of the arm with the largest circumference to standardized MRI measurements by comparing the bead's measured cross sectional area with that of the MRI determined cross sectional area.

Entry MRI was done 24 to 48 h before the isometric or 1RM (1 repetition maximum) test. Scans post-training MRI was performed 48 – 96 h after the last training session. Fifteen 16 mm contiguous axial slices from each arm were taken from each arm independently. The top of the bead in a sagittal scout view was used to locate the 8th slice going from the top of the arm toward the elbow. Scans for both arms were taken by Fast Spoiled Gradient Recalled (FSPPGR) and Fast Spin Echo (FSE) with TE 1.8/TR 200 msec. All 8 centers submitted the MRI data to the Research Center for Genetic Medicine at Children's National Medical Center (CNMC) in Washington, D.C, via e-mail or DAT disc, and all scans were integrated into the study SQL database.

For both cross-sectional and volumetric analysis of the MRI images, we used Rapidia (INFINITT Inc, Seoul Korea), a PC based software that allows the semi-automatic quantification of muscle, bone and subcutaneous fat (figure 1). The software was optimized to provide automated position adjustment, and to distinguish muscle, fat, and bone, with automatic edge detection, user modification of ambiguous edges, and automated propagation of defined tissue boundaries when possible. Volume measures were taken using an anatomical landmark (met-

Table 1: Demographic characteristics of Caucasian FAMuSS cohort

| Characteristic | Females | | Males | |
|---|---------|-----------------|-------|--------------------|
| | N | Mean ± SD | N | Mean ± SD |
| Age (years) | 378 | 24.12 ± 7.75 | 223 | 26.48 ± 11.70* |
| Baseline body mass (lbs.) | 378 | 144.35 ± 28.17 | 223 | 180.19 ± 37.39 ** |
| Baseline height (in.) | 378 | 64.96 ± 2.63 | 223 | 69.94 ± 2.76 ** |
| Baseline BMI | 378 | 24.06 ± 4.66 | 223 | 25.88 ± 5.12 ** |
| Post-exercise body mass (lbs.) | 378 | 145.40 ± 28.03 | 223 | 181.04 ± 37.15 ** |
| Post-exercise BMI | 378 | 24.22 ± 4.62 | 223 | 25.98 ± 5.06 ** |
| Fasting glucose (mg/dL) | 378 | 85.59 ± 7.60 | 223 | 90.47 ± 11.67 ** |
| Cholesterol (mg/dL) | 359 | 167.48 ± 32.79 | 218 | 168.61 ± 30.70 |
| HDL (mg/dL) | 359 | 52.02 ± 11.60 | 218 | 40.95 ± 11.14 ** |
| LDL (mg/dL) | 359 | 96.02 ± 28.61 | 218 | 103.75 ± 27.71 *** |
| Fasting insulin (uIU/mL) | 378 | 5.37 ± 5.26 | 223 | 6.09 ± 5.76 |
| Mean BP ^ | 372 | 85.52 ± 8.75 | 218 | 90.17 ± 9.21 ** |
| HOMA ^^ | 378 | 1.16 ± 1.49 | 223 | 1.40 ± 1.48 |
| Subcutaneous fat volume of the trained arm (mm ³) | 273 | 260697 ± 119542 | 146 | 181908 ± 97743 ** |
| Subcutaneous fat volume of the untrained arm (mm ³) | 273 | 261558 ± 122587 | 146 | 184542 ± 105259 ** |

*p = 0.003

**p < 0.0001

***p = 0.0015

^Mean BP calculated as (SBP + DBP*2)/3

^^ HOMA calculated as ((Fasting glucose (mg/dL) * 0.0551 * Fasting insulin (uIU/mL))/22.5)

aphyseal-diaphyseal junction of the humerus) as our starting point and assayed the six 1 cm slices proximal to it. Reliability of measures was determined using the untrained arm, with fat volume measured at a 12 wk interval, showing that our quantitation of subcutaneous fat was highly reliable and sensitive ($R^2 = 0.943$) (figure 2). Measurement values were automatically written and saved in a SQL database together with anthropomorphic and genotyping data.

Serum biochemistry

Venous blood samples were collected from fasting subjects and serum was separated from blood cells by centrifugation at $1100 \times g$ for 10 min and frozen for further analysis. Serum samples were analyzed by QuestDiagnostic.

DNA extraction and genotyping

DNA was extracted from blood samples obtained by phlebotomy before starting the exercise training. Genotyping was done using the TaqMan allele discrimination assay that employs the 5' nuclease activity of Taq polymerase to detect a fluorescent reporter signal generated during PCR reactions. Oligonucleotide primers used for the L163V PPAR α assay are LVF: CAGAAACAAATGCCAGTATTGTCGAT and LVR: CTTACCTACCGTTGTGTGAC ATC. LVV2 VIC: ACAAGTGCCTTTCTG (G allele) and LVM2 AM: CAAGTGCCTTTCTG (C allele)

Statistical analyses

The Hardy-Weinberg equilibrium was determined for each SNP using a χ^2 test to compare the observed genotype frequencies to those expected under H-W equilibrium.

Three volumetric measurements were analyzed as continuous quantitative traits (baseline and post-exercise subcutaneous fat volumes and difference in subcutaneous fat volume from baseline to post exercise) for each arm. Normality of each quantitative trait was confirmed using the Shapiro-Wilk normality test.

Bivariate correlation analyses of each quantitative measurement showed several significant correlations with age and baseline mass, therefore, associations between each SNP and volumetric measurements were assessed using analysis of covariance (ANCOVA) methods. Due to large gender differences in baseline values and the response to training, all analyses were performed separately for men and women. All significant associations from the main ANCOVA model were subjected to pair-wise statistical tests among each of the three genotype groups for each SNP. Linear tests were performed between each of the genotype groups to determine which genotype groups were significantly different from one another. The resulting p-values from these linear tests were adjusted for multiple comparisons using the Sidak post-hoc multiple comparison




Figure 1

3D determination of fat volume. Rapidia determination of subcutaneous fat volume. Panel A: the pink selected area shows the subcutaneous fat of one of the six slices measured in the right arm. The red arrow shows the epiphyseal flare (anatomical landmark used as the starting point of fat determination). Panel B: Left arm, the pink area shows the subcutaneous fat volume for other subject. Panel C and D represent the 3D image of panel A and B.

son test. Linear regression analysis, including likelihood ratio tests between full (containing genotype and covariates) and constrained (containing covariates only) models, were performed to estimate the proportion of variance in volumetric measurements attributable to each SNP's genotype.

Results

Genotype association of PPAR α L162V polymorphism with lipid concentrations

610 Caucasian subjects were selected for this study, based upon availability of serum samples for lipid and glucose determination (n = 578), and/or complete MRI data (n =

526) (Table 1). All subjects were genotyped for the PPAR α L162V polymorphism using automated TaqMan assays. Genotype distribution was in Hardy-Weinberg equilibrium. The allele frequencies of the PPAR α L162V showed the rare Valine allele at 7.5% in Caucasians, with 18% of Caucasians heterozygous for the allele. This is similar to previous reports in other Caucasian populations.

The PPAR α 162 valine allele was associated with significantly greater serum triglyceride and lower HDL-chol levels in Caucasian men. Heterozygotes for L162V showed 78% higher triglyceride levels than homozygotes for the common ancestral allele (n = 578; LL genotype = 116.15




Figure 2
Pearson correlation between baseline and post-training subcutaneous fat in the untrained arm of the entire cohort. The correlation coefficient between baseline and post exercise subcutaneous fat volume in the untrained-arm was $R^2 = 0.943$ ($P = 0.001$).

± 10.82 mg/dL LV genotype = 207.49 ± 29.49 mg/dL; $p = 0.004$) (Table 2). As 18% of the population carried the rare allele, the genotype effect of the SNP for all variation in triglyceride levels was approximately 3.8%. This was a larger effect size than previously reported for any genetic modifier of serum triglyceride levels.

LDL cholesterol levels tended to be higher among carriers of the V162 allele, although these results did not reach statistical significance (LL genotype = 96 mg/dL LV genotype = 98 mg/dL).

Heterozygous men showed 20% lower HDL levels than homozygotes (LL genotype = 41.77 ± 0.78 mg/dL LV genotype = 34.54 ± 2.13 mg/dL $p = 0.001$).

No significant association between PPAR α genotype and lipids in women, for any subtype of lipid (Table 2).

Genotype association of PPAR α L162V polymorphism with insulin resistance

We calculated insulin resistant using homeostatic model assessment (HOMA-IR), where the product of fasting glucose (mmol/L) and fasting insulin (μ U/mL) was divided by 22.5. No significant differences were seen for insulin resistance and the 162V polymorphism in our cohort.

Genotype association of PPAR α L162V polymorphism with baseline subcutaneous fat volume

Fat volume was determined using semi-automated three-dimensional reconstruction analysis software (Figure 1), with anatomical location specified by the epiphyseal flare of the humerus. Fat volume was measured for both upper arms, before and after a 12 week supervised resistance training program implemented only on one arm. Reliability of measures was determined using the untrained arm, with fat volume measured at a 12 wk interval, showing that our quantitation of subcutaneous fat was highly reliable and sensitive ($R^2 = 0.943$) (figure 2). We considered

Table 2: Analysis of Serum measurements and PPAR alpha (L162V) in Caucasians

| Serum Measurement | Group | Covariates | P-value | N; Adjusted mean \pm SEM | p-value for significantly different means | % variation explained by genotype; LRT p-value |
|-------------------|--------------|------------|---------|---|---|--|
| Triglycerides | All subjects | Age | 0.0026 | CC (N = 500; 102.92 ± 4.49)* CG (N = 71; 146.80 ± 11.84)* GG (N = 7; 114.99 ± 37.98) | * $p = 0.0018$ | 2.0%; 0.0025 |
| | Females | Age | 0.0474 | CC (N = 307; 94.76 ± 2.59)* CG (N = 45; 111.11 ± 6.77)* GG (N = 7; 114.72 ± 17.17) | NONE | 1.7%; 0.0458 |
| | Males | Age | 0.0040 | CC (N = 193; 116.15 ± 10.82)* CG (N = 26; 207.49 ± 29.49)* | * $p = 0.0040$ | 3.8%; 0.0037 |
| HDL | All subjects | Age | 0.0076 | CC (N = 500; 48.13 ± 0.56) CG (N = 70; 44.67 ± 1.48)* GG (N = 7; 58.43 ± 4.74)* | * $p = 0.0173$ | 1.7%; 0.0073 |
| | Females | Age | NS | | | |
| | Males | Age | 0.0017 | CC (N = 193; 41.77 ± 0.78)* CG (N = 26; 34.54 ± 2.13)* | * $p = 0.0017$ | 4.4%; 0.0015 |
| Cholesterol | All subjects | Age | NS | | | |
| | Females | Age | NS | | | |
| | Males | Age | NS | | | |
| VLDL-TG | All subjects | Age | NS | | | |
| | Females | Age | 0.0441 | CC (N = 307; 18.94 ± 0.52)* CG (N = 45; 22.29 ± 1.35)* GG (N = 7; 22.77 ± 3.43) | NONE | 1.7%; 0.0426 |
| | Males | Age | NS | | | |
| LDL | All subjects | Age | NS | | | |
| | Females | Age | NS | | | |
| | Males | Age | NS | | | |
| Fasting glucose | All subjects | Age | NS | | | |
| | Females | Age | 0.0440 | CC (N = 307; 85.78 ± 0.41) CG (N = 45; 83.65 ± 1.03) GG (N = 7; 89.71 ± 2.58) | NONE | 1.5%;0.0426 |
| | Males | Age | NS | | | |
| Fasting insulin | All subjects | Age | NS | | | |
| | Females | Age | NS | | | |
| | Males | Age | NS | | | |
| | Males | Age | NS | | | |

six phenotypes for association with PPARα L162V: absolute fat volume baseline (dominant and non-dominant arms), absolute fat volume after unilateral strength training intervention (both dominant, and non-dominant [trained] arms), and absolute change in fat volume (both dominant and non-dominant arms). All data were stratified for sex (Table 3).

The PPARα 162 valine allele was associated with significantly greater baseline subcutaneous fat volume in Caucasian (Table 3). Four different phenotypes all showed significant genotype effect for absolute measures of fat volume (trained, un-trained, entry and exit; Table 3). Male heterozygotes for L162V showed 24% to 38% higher subcutaneous fat of the arm than homozygotes for the common ancestral allele (LL genotype = 162,106 mm³; LV genotype = 256,388 mm³). As 18% of the population carried the rare allele, the genotype effect of the SNP for all variation in regional baseline subcutaneous fat volume in men was approximately 2% to 4.2% ; Table 3).

We also examined the association between PPARα and BMI in Caucasian men (as an independent variable). The Valine allele was associated with higher BMI in Caucasian

men (P = 0.0031) (LL genotype = 25.10, LV genotype = 27.54) (n = 202).

For women, our analysis did not show any statistically significant genotype effect for subcutaneous fat.

Genotype effect on exercise-induced changes in subcutaneous fat volume

The PPARα 162 valine was associated with increased fat volume of the untrained arm in male carriers, while LL genotypes significantly decreased in fat volume (LL = -1,707 ± 21 mm³, LV = 17,617 ± 58 mm³; p = 0.002) in men (figure 3; table 3). This suggests that exercise results systemic loss of subcutaneous fat in LL males, while the LV carriers showed a large gain in subcutaneous fat.

As with baseline genotype associations, we found no association in women for change of subcutaneous fat as a function of exercise.

Discussion

The major goal of this study was to identify the role that PPARα L162V polymorphism may play in the variability of serum lipid levels, BMI, subcutaneous fat volume, and

Table 3: Analysis of PPAR alpha (L162V) in larger Caucasian only MRI cohort:

| Subcutaneous fat measure | Gender | SNP | N: adjusted mean ± SEM | P-value for significantly different means | Variability attributable to genotype | Likelihood-ratio test p-value * |
|--|--------|--------------------|---|---|--------------------------------------|---------------------------------|
| Full model – with age only adjustment | | | | | | |
| Trained arm – baseline fat volume | Male | PPAR alpha (L162V) | CC (N = 179; 167221 ± 6710)* CG (N = 24; 207418 ± 18325) * | *p = 0.041 | 2.0% | 0.039 |
| Trained arm – post-exercise fat volume | Male | PPAR alpha (L162V) | CC (N = 179; 170608 ± 6952)* CG (N = 24; 214926 ± 18932) * | *p = 0.029 | 2.3% | 0.028 |
| Untrained arm – baseline fat volume | Male | PPAR alpha (L162V) | CC (N = 179; 169890 ± 7106)* CG (N = 24; 224144 ± 19407) * | *p = 0.033 | 2.2% | 0.031 |
| Untrained arm – post-exercise fat volume | Male | PPAR alpha (L162V) | CC (N = 179; 168183 ± 7241)* CG (N = 24; 231960 ± 19776) * | *p = 0.003 | 4.2% | 0.003 |
| Untrained arm – difference in fat volume | Male | PPAR alpha (L162V) | CC (N = 179; -1707 ± 2133)* CG (N = 24; 17617 ± 5826) * | *p = 0.002 | 4.6% | 0.002 |
| Full model – with age and weight adjustment | | | | | | |
| Untrained arm – difference in fat volume | Male | PPAR alpha (L162V) | CC (N = 179; -1921 ± 2134)* CG (N = 24; 17612 ± 5825) * | *p = 0.002 | 4.9% | 0.001 |

Figure 6.




Figure 3
PPAR α L162V is associated with changes in subcutaneous fat volume in response to exercise in men. In men heterozygote for the V162 exercise training increased fat volume of the untrained arm. LL genotypes significantly decreased in fat volume. No significant differences were seen for women. Aged adjusted model.

change in fat volume in response to unilateral upper arm resistance training in 610 young (24 yrs) healthy volunteers.

We found the V allele to increase triglycerides by 78% ($p = 0.004$), and this single polymorphism accounted for 3.8% of all variation in serum triglycerides in males ($p = 0.0037$). PPAR α L162V has been previously shown to be associated with cholesterol [19] LDL-cholesterol [16-18], HDL-cholesterol [18] or triglycerides levels [20,21] depending on the population studied. However, the majority of these association studies have not been replicated. For example, the effect of the PPAR α L162V polymorphism on lipid profiles has been studied in middle age subjects with and without endo-phenotypes of metabolic syndrome in three different reports. Vohl et al. [17] reported 10% higher LDL cholesterol levels in carriers of the V allele in non-diabetic men ($n = 193$), but not in a diabetic population of men and women ($n = 120$) ($p = 0.02$). In contrast, Flavell et al. have found higher levels of LDL cholesterol in carriers of the V allele in subjects with type 2 diabetes ($n = 129$) but not in healthy subjects ($n = 2508$) [18]. Moreover, Lacquemant et al. reported higher cholesterol levels in subjects with metabolic syndrome only in the presence of concomitant coronary heart disease (CHD) [19]. The dis-

crepancies among these studies may be the result of effect modifiers in different populations, with disease state, advanced age, and sex each potentially modulating genotype/phenotype associations. Our underlying rationale in this current study was that analysis of young pre-morbid normal populations would provide the most sensitive and specific window into PPAR α associations with lipid profiles. Consistent with our hypothesis, the association of PPAR α and triglycerides in males in this current study showed a much larger effect of the V allele than previously reported in older and less healthy populations, with the rare V allele nearly doubling circulating triglycerides.

Our data showed that the V allele resulted in a 24–38% increase in subcutaneous fat, and a 20% increase in BMI ($p = 0.0031$) (CC genotype = 25.10 kg/m², CG genotype = 29.54 kg/m²) in White males. We also found that the PPAR α polymorphism had a strong effect on change in subcutaneous fat volume after 12 wks of unilateral resistance training in males. Specifically, males homozygous for the common L allele showed a decrease in fat volume following unilateral resistance training, while V allele heterozygotes showed an increase in subcutaneous fat ($p = 0.002$). The effect was on both the trained and untrained arms, suggesting that the PPAR α genotype had a strong systemic effect on subcutaneous fat metabolism, where the L allele homozygotes showed a beneficial effect of exercise on adiposity, while the V allele carriers put on weight with exercise. To our knowledge, this is the first demonstration of the effect of PPAR α genotype on the outcome of an intervention.

While PPAR α Valine allele showed strong statistical significance with higher subcutaneous fat and BMI in young Caucasian men in our study, the two previous reports showed association of the Valine allele with lower percentage of body fat, and lower BMI, in overweight and diabetic women [22,23]. A possible explanation for this apparent discrepancy is as follows. Functional studies on PPAR α L162V have reported a higher transactivation of the receptor in the presence of the variant allele (162V) in a ligand dependant way [15]. At low ligand levels, the activity of 162V receptor was lower than the wild type. However, at high ligand levels, the activity of the polymorphic receptor was higher. Since the activity of the receptor is in part dictated by the concentration of the receptor agonists and PPAR α is activated by fatty acids among others, we would expect to see higher activity of the polymorphic receptor in obesity, metabolic syndrome and type 2 diabetes (in which the levels of fatty acids are higher). Levels of endogenous and exogenous ligands (such as dietary fatty acids) may dictate the final phenotype observed in the presence of the polymorphism. Our data in young pre-morbid subjects on lipid levels, BMI

and subcutaneous fat volume is consistent with the 162V allele showing lower PPAR α activity.

One finding of our study is that women do not show the same PPAR α L162V associations as do men. It is clear that gender greatly influences both PPAR α expression, and also use of alternative lipid metabolism pathways, and that both are highly hormone responsive. Djouadi et al. [25] reported that in PPAR α null mice the pharmacological inhibition of fatty acid cellular flux caused massive hepatic and cardiac lipid accumulation, hypoglycemia (secondary to reduced gluconeogenesis due to the inhibition of lipid β oxidation), and death in 100% males but only 25% females. The pretreatment of males with estrogens abolished the difference between the two sexes. In other studies, Jalouli et al. [26] found higher PPAR α mRNA and protein levels in male rats than in female rats. Gonadectomy of male rats reduced the expression of PPAR α to similar levels as intact female rats. This suggested that not only estrogens may have a role in the regulation of lipid metabolism but also androgens may be involved through the increased expression of PPAR α . Taking these results together, young females with circulating estrogen levels appear to have alternative lipid metabolism pathways that would serve to mask the effect of the PPAR α polymorphism, as we have found in our young adult populations.

It is important to note that the relatively rare frequency of the V allele leads to small group sizes of V carriers. Future studies using specific interventions on pre-genotyped groups of V allele carriers will be necessary to more fully characterize the effect of the PPAR V allele on metabolism and measures of obesity.

Conclusion

We found statistical support for a well characterized PPAR α polymorphism (L162V) with plasma triglycerides, HDL, subcutaneous fat volume, and change in fat following an exercise intervention in 610 Caucasian young healthy volunteers. Those Caucasian men carrying the 162V allele have nearly double serum triglyceride levels, lower HDL levels, greater regional subcutaneous fat volume, higher BMI, and tend to gain subcutaneous fat volume with an exercise intervention.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

JU performed the genotyping and draft the manuscript. HGD carried out statistical analysis. MB participated in genetic counselling. BM helped to genotype. CTR, JD, BH, EKR and CB carried out the quantification of fat volume.

BCH helped to draft the manuscript. RLS, PDT, TBP, TJA, PMC, NMM, LSP, PSV, RFZ and PMG participated in the design of the study. EPH conceptualized, designed the study and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Supported by a grant from the National Institutes of Health (NINDS/NIAMS/NIA ROI NS40606-01).

References

1. Langefeld CD, Wagenknecht LE, Rotter JJ, Williams AH, Hokanson JE, Saad MF, Bowden DW, Haffner S, Norris JM, Rich SS, Mitchell BD: **Linkage of the metabolic syndrome to 1q23-q31 in Hispanic families: the insulin Resistance Atherosclerosis Study Family Study.** *Diabetes* 2004, **53(4)**:1170-4.
2. Hebebrand J, Sommerlad C, Geller F, Gorg T, Hinney A: **The genetics of obesity: practical implications.** *Int J Obes Relat Metab Disord* 2001, **25(Suppl 1)**:S10-8.
3. Middelberg RP, Martin NG, Whitfield JB: **Longitudinal genetic analysis of plasma lipids.** *Twin Res Hum Genet* 2006, **9(4)**:550-7.
4. Fabsitz RR, Carmelli D, Hewitt JK: **Evidence for independent genetic influences on obesity in middle age.** *Int J Obes Relat MetabDisord* 1992, **16(9)**:657-66.
5. Pistilli EE, Gordish-Dressman H, Angelopoulos TJ, Clarkson PM, Moyna NM, Pescatello LS, Visich PS, Zoeller RF, Hoffman EP, Gordon PM: **Resistin polymorphisms are associated with muscle, bone, and fat phenotypes in white men and women.** *Obesity (Silver Spring)* 2007, **15(2)**:392-402.
6. Pescatello LS, Kostek MA, Gordish-Dressman H, Thompson PD, Seip RL, Price TB, Angelopoulos TJ, Clarkson PM, Gordon PM, Moyna NM, Visich PS, Zoeller RF, Devaney JM, Hoffman EP: **ACE ID genotype and the muscle strength and size response to unilateral resistance training.** *Med Sci Sports Exerc* 2006, **38(6)**:1074-81.
7. Hubal MJ, Gordish-Dressman H, Thompson PD, Price TB, Hoffman EP, Angelopoulos TJ, Gordon PM, Moyna NM, Pescatello LS, Visich PS, Zoeller RF, Seip RL, Clarkson PM: **Variability in muscle size and strength gain after unilateral resistance training.** *Med Sci Sports Exerc* 2005, **37(6)**:964-72.
8. Clarkson PM, Devaney JM, Gordish-Dressman H, Thompson PD, Hubal MJ, Urso M, Price TB, Angelopoulos TJ, Gordon PM, Moyna NM, Pescatello LS, Visich PS, Zoeller RF, Seip RL, Hoffman EP: **ACTN3 genotype is associated with increases in muscle strength in response to resistance training in women.** *J Appl Physiol* 2005, **99(1)**:154-63.
9. Costet P, Legendre C, More J, Edgar A, Galtier P, Pineau T: **Peroxisome proliferator-activated receptor alpha-isoform deficiency leads to progressive dyslipidemia with sexually dimorphic obesity and steatosis.** *J Biol Chem* **273(45)**:29577-85. Nov 6; 1998
10. Djouadi F, Weinheimer CJ, Saffitz JE, Pitchford C, Bastin J, Gonzalez FJ, Kelly DP: **A gender-related defect in lipid metabolism and glucose homeostasis in peroxisome proliferator-activated receptor alpha- deficient mice.** *J Clin Invest* **102(6)**:1083-91. Sep 15; 1998
11. Forman BM, Chen J, Evans RM: **Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta.** *Proc Natl Acad Sci USA* **94(9)**:4312-7. Apr 29; 1997
12. Fruchart JC, Duriez P, Staels B: **Peroxisome proliferator-activated receptor-alpha activators regulate genes governing lipoprotein metabolism, vascular inflammation and atherosclerosis.** *Curr Opin Lipidol* 1999, **10(3)**:245-57.
13. Vohl MC, Lepage P, Gaudet D, Brewer CG, Betard C, Perron P, Houde G, Cellier C, Faith JM, Despres JP, Morgan K, Hudson TJ: **Molecular scanning of the human PPAR α gene: association of the L162v mutation with hyperapobetalipoproteinemia.** *J Lipid Res* 2000, **41(6)**:945-52.
14. Tugwood JD, Aldridge TC, Lambe KG, Macdonald N, Woodyatt NJ: **Peroxisome proliferator-activated receptors: structures and function.** *Ann N Y Acad Sci* **804**:252-65. Dec 27; 1996
15. Sapone A, Peters JM, Sakai S, Tomita S, Papiha SS, Dai R, Friedman FK, Gonzalez FJ: **The human peroxisome proliferator-activated**

- receptor alpha gene: identification and functional characterization of two natural allelic variants.** *Pharmacogenetics* 2000, **10**:321-33.
16. Tai Es, Demissie S, Cupples LA, Corella D, Wilson PW, Schaefer EJ, Ordovas JM: **Association between PPARA L162V polymorphism and plasma lipid levels: the Framingham Offspring Study.** *Arterioscler Thromb Vasc Biol* **22(5)**:805-10. May 1: 2002
 17. Vohl Marie-Claude, Lepage Pierre, Gaudet Daniel, Brewer Carl G, Bétard Christine, Perron Patrice, Houde Ghislaine, Cellier Christine, Faith Janet M, Després Jean-Pierre, Morgan Kenneth, Hudson Thomas J: **Molecular scanning of the human PPARa gene: association of the L162V mutation with hyperapobetalipoproteinemia.** *J Lipid Res* 2000, **41(6)**:945-52.
 18. Flavell DM, Pineda Torra I, Jamshidi Y, Evans D, Diamond JR, Elkeles RS, Bujac SR, Miller G, Talmud PJ, Staels B, Humphries SE: **Variation in the PPAR gene is associated with altered function in vitro and plasma lipid concentrations in type II diabetic subjects.** *Diabetologia* 2000, **43**:673-680.
 19. Lacquemant C, Lepretre F, Pineda Torra I, Manraj M, Charpentier G, Ruiz J, Staels B, Froguel P: **Mutation screening of the PPAR. % Warning: EPS-printer was not specified – Figure omitted! gene in type 2 diabetes associated with coronary heart disease.** *Diabetes Metab* 2000, **26**:393-401.
 20. Robitaille J, Brouillette Cm Houde A, Lemieux S, Perusse L, Tchernof A, Gaudet D, Vohl MC: **Association between the PPARalpha-L162V polymorphism and components of the metabolic syndrome.** *J Hum Genet* 2004, **4;49(9)**:482-489.
 21. Shyong Tai E, Corella Dolores, Demissie Serkalem, Adrienne Cupples L, Coltell Oscar, Schaefer Ernst J, Tucker Katherine L, Ordovas Jose M: **Polyunsaturated Fatty Acids Interact with the PPARA-L162V Polymorphism to Affect Plasma Triglyceride and Apolipoprotein C-III Concentrations in the Framingham Heart Study.** *The American Society for Nutritional Sciences J Nutr* 2005, **135(3)**:397-403.
 22. Evans D, Aberle J, Wendt D, Wolf A, Beisiegel U, Mann WA: **A polymorphism, L162V, in the peroxisome proliferator-activated receptor alpha (PPARalpha) gene is associated with lower body mass index in patients with non-insulin-dependent diabetes mellitus.** *J Mol Med* 2001, **79(4)**:198-204.
 23. Bosse Y, Despres JP, Bouchard C, Perusse L, Vohl MC: **The peroxisome proliferator-activated receptor alpha L162V mutation is associated with reduced adiposity.** *Obes Res* 2003, **11(7)**:809-16.
 24. Thompson PD, Moyna N, Seip R, et al.: **Functional Polymorphisms Associated with Human Muscle Size and Strength.** *Med Sci Sports Exerc* 2004, **36(7)**:1132-9.
 25. Djouadi F, Weinheimer CJ, Saffitz JE, Pitchford C, Bastin J, Gonzalez FJ, Kelly DP: **A gender-related defect in lipid metabolism and glucose homeostasis in peroxisome proliferator- activated receptor alp alpha- deficient mice.** *J Clin Invest* **102(6)**:1083-91. Sep 15; 1998
 26. Jalouli M, Carlsson L, Ameen C, Linden D, Ljungberg A, Michalik L, Eden S, Wahli W, Oscarsson J: **Sex difference in hepatic peroxisome proliferator-activated receptor alpha expression: influence of pituitary and gonadal hormones.** *Endocrinology* 2003, **144(1)**:101-9.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2350/8/55/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

