

Migraine Strikes as Neuronal Excitability Reaches a Tipping Point

Marten Scheffer^{1*}, Albert van den Berg², Michel D. Ferrari³

1 Department of Aquatic Ecology & Water Quality Management, Wageningen University, Wageningen, the Netherlands, **2** MESA+ Institute for Nanotechnology, University of Twente, Enschede, the Netherlands, **3** Department of Neurology, Leiden University Medical Centre, Leiden, the Netherlands

Abstract

Self-propagating waves of cerebral neuronal firing, known as spreading depolarisations, are believed to be at the roots of migraine attacks. We propose that the start of spreading depolarisations corresponds to a critical transition that occurs when dynamic brain networks approach a tipping point. We show that this hypothesis is consistent with current pathogenetic insights and observed dynamics. Our view implies that migraine strikes when modulating factors further raise the neuronal excitability in genetically predisposed subjects to a level where even minor perturbations can trigger spreading depolarisations. A corollary is that recently discovered generic early warning indicators for critical transitions may be used to predict the onset of migraine attacks even before patients are clinically aware. This opens up new avenues for dissecting the mechanisms for the onset of migraine attacks and for identifying novel prophylactic treatment targets for the prevention of attacks.

Citation: Scheffer M, van den Berg A, Ferrari MD (2013) Migraine Strikes as Neuronal Excitability Reaches a Tipping Point. PLoS ONE 8(8): e72514. doi:10.1371/journal.pone.0072514

Editor: Claudia Sommer, University of Würzburg, Germany

Received: March 17, 2013; **Accepted:** July 14, 2013; **Published:** August 29, 2013

Copyright: © 2013 Scheffer et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The research of each of the authors is supported by a Dutch national science prize, the Spinoza Award. M.S. and A.v.d.B. are also supported by a European Research Council Advanced grant. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: marten.scheffer@wur.nl

Introduction

About 12% of humanity is plagued by recurring migraine attacks that are characterized by disabling headaches and associated autonomic symptoms [1]. In up to one third of patients, attacks may be associated with neurological (usually visual) aura symptoms, which are caused by the electrophysiological phenomenon of “cortical spreading depression” [2]. This is a pattern of autonomous neuronal firing activity that starts at a focal point within the brain, and subsequently propagates as a slow wave in all directions across the cortex and into subcortical brain regions, until it hits upon less sensitive regions of the brain [2,3]. Although convincing evidence in humans is still lacking, there is increasing evidence from animal experiments that spreading depression not only is the underlying mechanism for migraine aura but also may activate the trigeminovascular “headache generating” system and thus may trigger the whole attack in at least a proportion of patients [4–7].

While much research has focused on the electrophysiological basis of the spreading process across the cortex [8], surprisingly little is known about the mechanisms causing spreading depression to start. When, why and how does such an outbreak of spontaneous firing strike? Several factors have been shown to play a role, ranging from genetic variants [3,9–12] to hormonal fluctuations [13,14], all leading to neuronal hyperexcitability [12,15]. The hypothesis we put forward is that increasing excitability bring the brain to a tipping point for spreading depression, and that as this tipping point is approached, resilience of the brain decreases in the sense that increasingly small perturbations are sufficient to trigger the onset.

Results

We illustrate the rationale behind our hypothesis with a minimal model of the mechanism we have in mind. The model is ‘minimal’ in the sense that we only focus on the ingredients that are essential for understanding the potential for a tipping point. There is a large literature starting in the 1970’s linking mathematics of dynamical systems to diseases with complex and cyclic dynamics such as Cheyne Stokes respiration, epilepsy, bipolar disorder, and cardiac arrhythmias [16]. Also, there have been sophisticated models in computational neuroscience that specifically address spreading depression, its onset and spatial dynamics [8,17]. Here we do not go into such complex temporal and spatial dynamics. Rather, we focus on the simplest set of mechanisms needed to understand how the brain can arrive at a tipping point for spreading depression. We do not do explicitly consider the dynamics of individual neurons and their interactions in the complex network. Instead, we analyze the total activity and excitability of a local ‘population of neurons’ (a so-called mean field- or mass- model). This allows us to explain the essence based on only three assumptions (Fig. 1):

1. Activity of a group of neurons (A) increases with the level of incoming stimuli (S) and the overall neuronal excitability (ϵ).
2. Excitability is modulated by overall physiological conditions and genetic factors, and promoted by local neural activity, implying positive feedback mechanisms.
3. When excitability surpasses a critical level, neural activity may be triggered merely by the activity of neighboring neurons, thus turning into a locally contagious process.




Figure 1. Causal structure that may lead to a tipping point for autonomous firing, as illustrated by the minimal model.
doi:10.1371/journal.pone.0072514.g001

While the first assumption is straightforward, the background for the second assumption is that neural activity promotes intercellular concentrations of potassium and glutamate, which in turn promote the excitability of neurons in the area [18]. The third assumption follows from the observation that cortical spreading depression spreads centrifugally as a traveling wave over the cortex. Unless orchestrated from another brain area (for which there is no indication) this behavior implies a local contagious component to the firing of neurons.

Figure 2 explains graphically how a tipping point can arise from the above assumptions (for the mathematical model, and a more specific definition of the parameters and variables see the *material and methods* section). Imagine a dynamic equilibrium of neural activity in a small brain region, resulting from the balance between generation and decay of pulses. If we plot the generation and decay of pulses together, it can be seen that multiple equilibria may arise at their intersections if the generation of activity rises relatively sharply when the contagious local neural firing kicks in (positive feedback) around a critical excitability level (ϵ_{crit}). The unstable middle equilibrium marks the boundary between the two ‘basins of attraction’ around the stable states.

To see how the brain can be brought to a tipping point, imagine what happens if the base-line excitability ϵ_0 increases (Fig. 2b). The unstable point moves closer to the normal stable state causing the basin of attraction for the normal state to shrink. Therefore the resilience becomes smaller, in the sense that a smaller perturbation is needed to let the system fall into the attraction basin of the hyperactive (aura/migraine) state. A further increase in base-line excitability can cause the unstable point to collide with the normal stable state, causing it to lose stability and disappear. This is the real ‘tipping point’, or in mathematical terms a fold-bifurcation point. The way in which the equilibria are affected by base-line excitability (and/or stimulus strength) can be summarized in one graph (Fig. 2c) known as a catastrophe fold, where the dashed middle section represents the repelling unstable equilibria that mark the border of the attraction basins of the normal state and the aura state. The essence of the interpretation of changing resilience can be captured in an intuitive way by a stability landscape representation (fig. 3). Each of the landscapes corresponds to a different physiological condition (captured in excitability ϵ_0 in our minimal model). Note that the aura/migraine state in practice is a transient rather than a true stable state. Hyperactivity stops at the rear end of the travelling wave of cortical spreading depression as a result of exhaustion of ATP or oxygen. We are not further addressing the mechanisms of how the process spreads and ultimately halts [17], as we focus on the triggering mechanisms.




Figure 2. Graphical model showing how a tipping point for cortical spreading depression can arise. a) Three equilibria may occur at intersection points where the rate of generation of new pulses (sigmoidal curve) equals the rate of decay (dashed line) of neural pulses. Activity increases when the generation of new pulses exceeds the decay of pulses (sections I and III) and decreases in the other sections (sections II and IV). It can be seen from the arrows representing this direction of change that the middle intersection point is a repeller that marks the border between the basins of attraction of the two alternative stable states. b) Increasing base-line excitability promotes the generation of new pulses causing the unstable equilibrium (open dot) and the stable normal state (left hand solid dot) to move closer together. This reduces resilience of the normal state in the sense that a smaller perturbation is needed to invoke a shift to the Aura state (horizontal dashed arrows in panel). c) Plotting how the intersection points representing equilibria move as a function of base-line excitability, a catastrophe fold arises. The fold bifurcation point (F) marks the loss of stability of the normal state.
doi:10.1371/journal.pone.0072514.g002




Figure 3. Stability landscape interpretation of how resilience of the normal mode of brain activity can be lost at high levels of base-line excitability as determined by genetically coded or other physiological conditions. The catastrophe fold at the base plane corresponds to the one depicted in figure 2c. doi:10.1371/journal.pone.0072514.g003

Discussion

The presence of a tipping point implies four well-defined predictions that can be checked against empirical observations [19,20]:

1. The system occasionally goes through a sudden transition.
2. A system at any given moment tends to be in either of the two contrasting states rather than somewhere in between.
3. A short perturbation can trigger a shift to a lasting contrasting regime.
4. Change in conditions can reduce the resilience of the system in the sense that a smaller perturbation is needed to invoke a shift.

Clearly the character of cortical spreading depression is well in line with these predictions: The onset of cortical spreading depression sharply breaks the normal regime of the brain (1); It is an all-or-none phenomenon with no evidence for stable intermediate states (2) and; Perturbations such as light flashes can invoke a shift (3) [18]. The fact that the latter happens only rarely in non-migraineurs and that an attack may sometimes be provoked by even a small stimulus (e.g. a light flash) implies indirect evidence for prediction (4). In addition, the observation that the amplitude of electroencephalographic responses to light flashes tends to be systematically larger in the days before a

migraine attack [21,22] is well in line with the view that increased excitability may periodically bring patients closer to a tipping point.

The view of cortical spreading depression-induced migraine as a critical transition at a tipping point opens up entirely new angles when it comes to perspectives for attack prediction and treatment. Most importantly, it has recently been shown that the distance to a tipping point in complex systems may be estimated based on generic early warning signals [19]. For instance, the recovery time to the normal regime upon a small perturbation is predicted to become longer as a system comes closer to the critical point [19,23]. This has just been demonstrated for a living system [24], and it would be a logical step to study the decay rate of brain responses to small perturbations such as visual stimuli. As the brain continuously generates activity, development of a robust protocol to detect such slowing down is not trivial. However, if we find ways to objectively measure resilience this way, this opens up novel possibilities. Firstly, we may give patients a simple home test to warn for impending attacks. They could then start pre-emptive short term prophylaxis to reduce excitability and enhance resilience, e.g. with antiepileptic agents known to be also effective in migraine [25], and in addition avoid strong stimuli or activities which are known to specifically trigger their migraine attacks. Secondly, from a research perspective, objective measures of resilience would help to effectively search for mechanisms involved. Now we typically classify subjects into groups such as migraineurs versus healthy persons. However, in our view resilience within these groups may vary strongly, and also within subjects resilience will vary over time. Objective measurements of resilience would greatly enhance the power for instance of statistical tools that can put us on the track of factors related to resilience, including time-varying aspects such as blood or cerebrospinal levels of substances. Some factors contributing to resilience are now well understood, but much could be gained in terms of treatment design if we could develop a more comprehensive insight in the mechanisms that can strengthen or undermine resilience against the onset of cortical spreading depression.

Finally, other episodic brain disorders such as epilepsy may benefit as well from this type of research. Migraine and epilepsy are comorbid episodic brain disorders that have common pathophysiologic mechanisms and treatments. Migraine attacks, like epileptic seizures, may be triggered by excessive neocortical cellular excitability; in migraine, however, the hyperexcitability is believed to transition to cortical spreading depression rather than to the hyper-synchronous activity that characterizes seizures [25].

Materials and Methods

To translate our three assumptions into a mathematical model, imagine a dynamic equilibrium of neural activity (A) in a small region, resulting from the generation and decay of pulses. A simple way to formulate that consistent with our assumptions could be:

$$\frac{dA}{dt} = (\varepsilon \cdot S + q \cdot A \frac{\varepsilon^p}{\varepsilon^p + \varepsilon_{crit}^p}) (1-A) - d \cdot A. \quad (1)$$

The last term in this equation reflects a simple proportional decay (at rate d) of pulses. The rest of the equation describes the generation of pulses. The multiplier $(1-A)$ goes to zero as A approaches unity, causing activity to be limited to values between zero and one. The term $\varepsilon \cdot S$ represents the generation of pulses resulting from external stimuli (S) depending on the excitability (ε)

of the neurons. The remaining term reflects the generation of pulses caused by the firing activity (A) of the other neurons in the area. The maximum intensity of this ‘contagion effect’ is scaled by a factor (q). The effect really kicks in when a critical excitability (ε_{crit}) is reached. This threshold response is formulated as the so-called Hill function ($\varepsilon^p / (\varepsilon^p + \varepsilon_{crit}^p)$) that for sufficiently high values of p increases with ε in a steep sigmoidal way from zero to one around the critical excitability level (ε_{crit}).

To include the feedback effect of activity on excitability we assume a linear relationship for simplicity:

$$\varepsilon = \varepsilon_0 + c \cdot A. \quad (2)$$

Where ε_0 is the base-line excitability as determined by predisposition through genetic factors and general physiological conditions, and c is the steepness of increase in excitability with neural activity. The full model is thus defined by substituting equation 2 for each ε in equation 1:

$$\frac{dA}{dt} = ((\varepsilon_0 + cA) \cdot S + q \cdot A \frac{(\varepsilon_0 + cA)^p}{(\varepsilon_0 + cA)^p + \varepsilon_{crit}^p}) (1-A) - d \cdot A. \quad (3)$$

For suitable parameter settings (e.g. our defaults: $S = 0.1$; $d = 0.1$; $e = 1.5$; $f = 1$; $h = 5$; $p = 4$) this model has a tipping point where the system may switch between a normal activity regime to a state of self-propelled hyperactivity (the spreading depression). To illustrate in an intuitive way why such behavior can arise from our assumptions we turn to a simple graphical representation.

Acknowledgments

The work greatly benefitted from discussions with Ronald Zielman, Gert van Dijk, Egbert H. Van Nes, Vasilis Dakos, and Ingrid van de Leemput.

Author Contributions

Contributed reagents/materials/analysis tools: MS. Wrote the paper: MS AvdB MF. Conceived the idea: MS MF.

References

- Goadsby PJ, Lipton RB, Ferrari MD (2002) Migraine - Current understanding and treatment. *New England Journal of Medicine* 346: 257–270.
- Lauritzen M (1994) Pathophysiology of the migraine aura: The spreading depression theory. *Brain* 117: 199–210.
- Eikermann-Haerter K, Yuzawa I, Qin T, Wang Y, Baek K, et al. (2011) Enhanced subcortical spreading depression in familial hemiplegic migraine type 1 mutant mice. *Journal of Neuroscience* 31: 5755–5763.
- Dalkara T, Zervas NT, Moskowitz MA (2006) From spreading depression to the trigeminovascular system. *Neurological Sciences* 27: s86–s90.
- Zhang X, Levy D, Kainz V, Noseda R, Jakubowski M, et al. (2011) Activation of central trigeminovascular neurons by cortical spreading depression. *Annals of Neurology* 69: 855–865.
- Levy D, Moskowitz MA, Noseda R, Burstein R (2012) Activation of the migraine pain pathway by cortical spreading depression: Do we need more evidence? *Cephalalgia* 32: 581–582.
- Karatas H, Erdener SE, Gursoy-Ozdemir Y, Lule S, Eren-Kocak E, et al. (2013) Spreading Depression Triggers Headache by Activating Neuronal Panx1 Channels. *Science* 339: 1092–1095.
- Dahlem MA, Hadjikhani N (2009) Migraine aura: Retracting particle-like waves in weakly susceptible cortex. *PLoS ONE* 4.
- Van Den Maagdenberg AMJM, Pizzorusso T, Kaja S, Terpolilli N, Shapovalova M, et al. (2010) High cortical spreading depression susceptibility and migraine-associated symptoms in Cav2.1 S218L mice. *Annals of Neurology* 67: 85–98.
- Eikermann-Haerter K, Moskowitz MA (2008) Animal models of migraine headache and aura. *Current Opinion in Neurology* 21: 294–300.
- Ayata C, Jin H, Kudo C, Dalkara T, Moskowitz MA (2006) Suppression of cortical spreading depression in migraine prophylaxis. *Annals of Neurology* 59: 652–661.
- Tottene A, Conti R, Fabbro A, Vecchia D, Shapovalova M, et al. (2009) Enhanced Excitatory Transmission at Cortical Synapses as the Basis for Facilitated Spreading Depression in Cav2.1 Knockin Migraine Mice. *Neuron* 61: 762–773.
- Eikermann-Haerter K, Baum MJ, Ferrari MD, Van Den Maagdenberg AMJM, Moskowitz MA, et al. (2009) Androgenic suppression of spreading depression in familial hemiplegic migraine type 1 mutant mice. *Annals of Neurology* 66: 564–568.
- Eikermann-Haerter K, Dileköz E, Kudo C, Savitz SI, Waerber C, et al. (2009) Genetic and hormonal factors modulate spreading depression and transient hemiparesis in mouse models of familial hemiplegic migraine type 1. *Journal of Clinical Investigation* 119: 99–109.
- Moskowitz MA, Bolay H, Dalkara T (2004) Deciphering Migraine Mechanisms: Clues from Familial Hemiplegic Migraine Genotypes. *Annals of Neurology* 55: 276–280.
- Glass L, Mackey MC (1988) From clocks to chaos: the rhythms of life. Princeton: Princeton University Press. 248 p.
- Dahlem MA, Graf R, Strong AJ, Dreier JP, Dahlem YA, et al. (2010) Two-dimensional wave patterns of spreading depolarization: Retracting, re-entrant, and stationary waves. *Physica D: Nonlinear Phenomena* in press.
- Bowyer SM, Aurora SK, Moran JE, Tepley N, Welch KMA (2001) Magnetoencephalographic fields from patients with spontaneous and induced migraine aura. *Annals of Neurology* 50: 582–587.
- Scheffer M, Bascompte J, Brock WA, Brovkin V, Carpenter SR, et al. (2009) Early-warning signals for critical transitions. *Nature* 461: 53–59.
- Scheffer M, Carpenter SR (2003) Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends in Ecology & Evolution* 18: 648–656.
- Kropp P, Gerber WD (1993) Is increased amplitude of contingent negative variation in migraine due to cortical hyperactivity or to reduced habituation? *Cephalalgia* 13: 37–41.
- Shibata K, Yamane K, Otuka K, Iwata M (2008) Abnormal visual processing in migraine with aura: A study of steady-state visual evoked potentials. *Journal of the Neurological Sciences* 271: 119–126.
- Van Nes EH, Scheffer M (2007) Slow recovery from perturbations as a generic indicator of a nearby catastrophic shift. *American Naturalist* 169: 738–747.
- Veraart AJ, Faassen EJ, Dakos V, van Nes EH, Lürling M, et al. (2012) Recovery rates reflect distance to a tipping point in a living system. *Nature* 481: 357–359.
- Rogawski MA (2008) Common pathophysiological mechanisms in migraine and epilepsy. *Archives of Neurology* 65: 709–714.