

REVIEW**Open Access**

Immunoregulatory properties of rapamycin-conditioned monocyte-derived dendritic cells and their role in transplantation

Camila Macedo¹, Héth Turquist^{1,2}, Diana Metes^{1,2} and Angus W Thomson^{1,2*}

Abstract

In efforts to minimize the chronic administration of immunosuppression (IS) drugs in transplantation and autoimmune disease, various cell-based tolerogenic therapies, including the use of regulatory or tolerogenic dendritic cells (tolDC) have been developed. These DC-based therapies aim to harness the inherent immunoregulatory potential of these professional antigen-presenting cells. In this short review, we describe both the demonstrated tolerogenic properties, and current limitations of rapamycin-conditioned DC (RAPA-DC). RAPA-DC are generated through inhibition of the integrative kinase mammalian target of rapamycin (mTOR) by the immunosuppressive macrolide rapamycin during propagation of monocyte-derived DC. Consistent with the characteristics of tolDC, murine RAPA-DC display resistance to phenotypic maturation induced by pro-inflammatory stimuli; exhibit the ability to migrate to secondary lymphoid tissue (important for 'cross-presentation' of antigen to T cells), and enrich for naturally-occurring CD4⁺ regulatory T cells. In rodent models, delivery of recipient-derived RAPA-DC pulsed with donor antigen prior to organ transplantation can prolong allogeneic heart-graft survival indefinitely, especially when combined with a short course of IS. These encouraging data support ongoing efforts to develop RAPA-DC for clinical testing. When compared to murine RAPA-DC however, human RAPA-DC have proven only partially resistant to maturation triggered by pro-inflammatory cytokines, and display heterogeneity in their impact on effector T-cell expansion and function. In total, the evidence suggests the need for more in-depth studies to better understand the mechanisms by which mTOR controls human DC function. These studies may facilitate the development of RAPA-DC therapy alone or together with agents that preserve/enhance their tolerogenic properties as clinical immunoregulatory vectors.

Keywords: Dendritic cells, Antigen presentation, Rapamycin, T cells, Regulatory T cells, Tolerance, Transplantation

Introduction

Given their capacity to safely prevent and/or reverse acute allograft rejection, immunosuppressive agents have proven crucial to the successful clinical development of organ transplantation. However, there are major limitations associated with drug-based immunosuppression (IS), including lack of antigen (Ag) specificity, failure to support tolerance induction, deficiencies in the prevention of late graft failure (chronic rejection), and significant morbidity. Evaluation of novel, tolerance-promoting

protocols, including cell-based therapies, such as the use of tolerogenic dendritic cells (tolDC), is a dynamic area of investigation and may provide a means to minimize or even replace use of IS drugs [1]. In addition to reducing the toxic burden of chronic IS, it is hoped that these innovative approaches will prevent/reduce chronic rejection, given the strong immunological involvement in its etiology [2,3].

DC play critical roles in Ag presentation to naïve and memory T cells and can either promote T-cell immunity or support the induction of tolerance [4-6]. Experimental protocols are currently being developed with the goal of harnessing the inherent tolerogenicity of DC to act as 'negative cellular vaccines', which can inhibit immune responses in an alloAg-specific manner and promote

* Correspondence: thomsonaw@upmc.edu

¹Thomas E. Starzl Transplantation Institute, Department of Surgery, University of Pittsburgh School of Medicine, 200 Lothrop Street, Pittsburgh, PA 15261, USA

²Department of Immunology, University of Pittsburgh School of Medicine, 200 Lothrop Street, Pittsburgh, PA 15261, USA

tolerance to transplanted cells and organs. TolDC are characteristically immature, express low surface MHC molecules, a low ratio of co-stimulatory to co-inhibitory signals, and an impaired ability to secrete T-helper-1 (Th1) cell-driving or inflammatory cytokines [2].

In particular, administration of tolDC presenting allo-Ag promotes transplant survival through the induction of T-cell hyporesponsiveness to allo-Ag, deletion of alloreactive T cells, or increased Treg incidence or functions [7-10]. This knowledge has driven efforts to identify mechanisms that underline the tolerogenic properties of DC to promote transplant tolerance.

Various anti-inflammatory and immunosuppressive agents have been used to generate tolDC *in vitro*, including interleukin (IL)-10 and transforming growth factor- β (TGF- β), cytotoxic T lymphocyte Ag-4 Ig (CTLA4-Ig), prostaglandin E2 (PGE2), dexamethasone, and vitamin D3 (vitD3), among others [11-13]. Of special interest to our lab has been defining the molecular and functional impact of the immunosuppressive pro-drug rapamycin (RAPA) on DC.

Phenotypic and functional characteristics of RAPA-DC

RAPA is a macrocyclic triene antibiotic with immunosuppressant properties, that was discovered in 1975 as a product of the bacterium *Streptomyces hygroscopicus* in a soil sample from Rapa Nui (Easter Island) [14]. This immunosuppressant inhibits the mammalian target of rapamycin (mTOR), a highly conserved serine/threonine kinase that controls cellular responses to environmental cues [15-17]. In mouse models, RAPA has a profound impact on DC *in vitro*, impairing their maturation following exposure to Toll-like receptor (TLR) ligands and suppressing their T-cell allostimulatory function [11,18-22] (Table 1). RAPA has been reported to have unique tolerance-promoting and Treg facilitating/sparing properties in small animal models of organ transplantation [23,24]. We [10,25] and others [7,21,26,27] have shown that when donor-derived, RAPA-conditioned DC (RAPA-DC) or recipient-derived RAPA-DC pulsed with donor allo-Ag are administered to recipients prior to transplantation, donor graft survival is prolonged indefinitely, especially when combined with a short course of low dose IS, such as RAPA, cyclosporine, or FK506 (Table 2). Taner *et al.* have shown, in the mouse model, prolongation of heart allograft survival when recipient-derived RAPA-DC pulsed with allo-Ag were given i.v. prior transplantation. Such results were improved with short-term administration of subtherapeutic dose FK506, which alone did not prolong graft survival or repeated infusion of RAPA-DC pulsed with allo-Ag (x3; days -10, -3, and 0) [10]. Turnquist *et al.* have also shown long-term heart allograft survival, after a single i.v. dose of

Table 1 Mouse vs. human immature RAPA-DC

	Mouse	Human
Phenotype [19,23,26,27]		
MHC Class II	↓	-
CD86/ CD40	↓↓	↓
B7-H1	↓	↓
Migration [10,23,24]		
CCR7	-	↑
Treg [21-23]		
enrichment/induction/expansion	↑	↑
T cell apoptosis [10]		
	↑	ND
Function [15,19,20,23,27-29]		
Endocytosis	↓↓	ND
MLR	↓↓	↓
IL-12p70	↓	-
IL-10	↓	-

ND, not done; -, not changed.

recipient-derived RAPA-DC pulsed with alloAg (day -7) followed by a short-term course of low-dose RAPA [25].

When rodent and human DC are generated in clinically-relevant concentrations of RAPA, they are phenotypically immature, with low levels of cell surface T-cell co-stimulatory molecules (CD86, CD40); however, only murine RAPA-DC maintain their immature phenotype when exposed to inflammatory stimuli, such as bacterial lipopolysaccharide (LPS) [11,25]. Also, mouse and human RAPA-DC exhibit a paradoxical decrease in cell surface expression of B7-H1 (also known as programmed death ligand-1; PD-L1), a PD-1 ligand, which contributes to the negative regulation of T lymphocyte activation and promotes peripheral tolerance [13,28] (Table 1).

Murine RAPA-DC induce hyporesponsiveness and/or apoptosis of alloreactive T cells [10,11,18,22,25]. Likewise, human RAPA-DC are poorly stimulatory and induce T-cell hyporesponsiveness [11]. Furthermore, murine RAPA-DC retain the capacity to stimulate mouse naturally-occurring Foxp3⁺ Treg, resulting in an overall enrichment of this population relative to T effector cells [25]. A similar capacity for promotion of Foxp3⁺ cells in T-cell cultures has been reported for human RAPA-DC [12] (Table 1).

Another important feature of murine RAPA-DC is their unaltered chemokine receptor (CCR7) expression and capacity for migration to CCL19/CCL21, and thus to secondary lymphoid tissues [10,25,26]. Human RAPA-DC have been reported to upregulate CCR7 expression and to display significantly augmented migration to CCL21 compared to control DC or other *ex vivo*-generated human tolDC, such as those conditioned with IL-10, dexamethasone, TGF- β , or vitamin

Table 2 Prolongation of allograft survival by RAPA-DC

Species	Graft	Source of DC	IS agent <i>in vivo</i>	Reference
Mouse	Heart	Bone marrow (BM) recipient-derived	None	10
			FK506	10
			RAPA	23
Rat	Limb	BM recipient-derived	ALS, CSA	18
Rat	Vascularized skin	BM recipient-derived	ALS, CSA	7
Mouse	Hematopoietic cells	Autologous BM-derived (unpulsed)	None	24
Mouse	Pancreatic islets	BM donor-derived	None	25

ALS, anti-lymphocyte serum; CSA, cyclosporine A; IS, immunosuppression.

D3 [13,30]. The ability of RAPA-DC to retain CCR7 expression/regulation and to migrate *in vivo* to secondary lymphoid tissue, while maintaining low expression of CD86 and diminished T-cell allostimulatory capacity, has important implications for their function as cellular therapy (that is, ‘negative’ vaccines) for prevention of transplant rejection [31] (Table 1).

RAPA-DC are also characterized by their unique cytokine production profile upon LPS or pro-inflammatory cocktail (IL-1 β , tumor necrosis factor (TNF)- α , IL-6, IFN- γ) stimulation. While IL-10 production is consistently reduced in RAPA-DC [12,28], their production of IL-12p70 may be affected differently. DC exposed to RAPA *in vivo* exhibit decreased IL-12p70 production in response to IL-4 stimulation; likewise, when DC are generated in culture with long exposure to RAPA, followed by stimulation with agonistic anti-CD40 mAb, these RAPA-DC display reduced IL-12p40 [18,25]. However, we have described increased IL-12p70 production by human monocyte-derived RAPA-DC after stimulation with LPS [11,29] or pro-inflammatory cytokines (Macedo *et al.*, manuscript in preparation). Increased production of IL-12p70 by RAPA-DC upon maturation (LPS stimulation) has been associated with augmented Th1/Th2-polarization of alloreactive CD4 $^{+}$ T cells [32] and with Th1 responses upon pro-inflammatory cytokine stimulation, towards IFN- γ production (Macedo *et al.*, manuscript in preparation). Further definition of the precise mechanisms by which mTOR controls and coordinates cytokine production and expression of B7-H1 by DC upon exposure to pro-inflammatory stimuli will be important to fundamental understanding of DC immunobiology and aid efforts to harness these promising immunoregulatory vectors in transplant medicine and autoimmune disease.

RAPA-DC from the bench to the clinic

The use of immunogenic or tolDC-based cell therapy in the clinic has been reported by groups working in different medical fields (cancer, HIV infection, and autoimmune diseases) with positive outcomes in terms of its feasibility and safety [33-36]. One aspect of tolDC-based

cell therapy in organ transplantation involves the use of donor-derived tolDC in an effort to improve graft survival; however, such protocols can only be applied in a live-donor setting since the *in-vitro* generation of tolDC takes 5 to 7 days, precluding use of tolDC generated from deceased donors. The generation of recipient-derived DC loaded with donor allo-Ag (donor cell lysate, apoptotic cells, or exosomes) is more advantageous, since the generation of autologous RAPA-DC can be performed at any time before transplantation and host peripheral mononuclear cells (PBMC) can be cryopreserved until time of tolDC generation/infusion. In addition, Ag presentation via the indirect pathway is thought to play an important role in the development of chronic rejection, making recipient-derived DC, if successful in regulating indirectly-alloreactive T cells, a potentially ground-breaking tolerogenic cell therapy in transplantation [37]. Immature DC such as RAPA-DC can also regulate the expansion and differentiation of Treg *in vitro* and *in vivo*, resulting in a ‘feedback’ regulatory loop [38,39]. On the current evidence, we cannot say whether pre- or post-transplant administration of tolDC, or whether autologous or donor-derived tolDCs, will prove to be a superior treatment; however, it is our personal opinion that alloantigen-pulsed recipient-derived DCs represent a pragmatic approach and offer certain theoretical advantages because of their indirect presentation of alloantigen. It is an exciting prospect that The ONE Study consortium will directly compare different approaches to tolerogenic APC therapy in a coherent clinical trial.

A means to obtain large numbers of monocytes is through their enrichment from peripheral blood leukaapheresis products. The ElutraTM cell separation system enriches monocytes untouched by antibodies or microbeads within a closed system on the basis of size and density [35,36]. Although not yet approved for clinical use in many countries, as an alternative, the ClinimACS[®] cell separation system isolates monocytes by positive selection using CD14 microbeads within a closed system with a good purity [40,41]. Monocytes generated using these techniques can be cultured in

medium containing cGMP-grade GM-CSF and IL-4, or using an Aastrom Replicell system [12,35]. After 5 to 7 days of incubation, DC can be loaded with allo-Ag, then evaluated for sterility, viability, recovery, and phenotype; and either aliquoted for cryopreservation or infused into the patient. We envisaged that the addition of RAPA during human DC culture/expansion would promote the tolerogenic features described above for murine RAPA-DC (Table 3). However, exposure of human RAPA-DC to maturation-inducing factors, such as pro-inflammatory cytokines or TLR4 ligands *in vitro* increased their production of IL-12p70, a Th1-inducing cytokine that could augment pathogen-specific CD8⁺ T cell responses and/or promote alloimmunity [11,42] and (Macedo et al., manuscript in preparation). As such, methods to limit IL-12p70 production should be examined as part of any protocol for RAPA-DC generation [11,43-45]. Recently, we have shown that increased IL-12p70 production, by both mouse and human RAPA-DC, following TLR4 ligation results from lost regulation of glycogen synthase kinase 3 (GSK-3) [11]. As treatment of RAPA-DC with GSK-3 inhibitors, such as lithium chloride, ablated IL-12p70 production, RAPA-DC treatment with GSK-3 inhibitors may be useful in limiting any potential danger of increase Th1 immunity following RAPA-DC administration [11].

Another way to maintain RAPA-DC tolerogenicity following infusion of donor-derived or allo-Ag-pulsed recipient-derived tolDC is the concomitant use of costimulation-blocking agents, such as abatacept (CTLA4-Ig) or belatacept (Lea 29Y), a first and second-generation CTLA4-Ag, respectively, that block the B7-CD28 costimulatory pathway [46,47]. Lu et al. [48] showed an increase in experimental organ graft survival when anti-CD40L mAb was administered in conjunction with donor-derived myeloid DC to block the CD40/CD40L pathway, which plays an important role in allogeneic DC-T cell interactions *in vivo*. Later, Kirk et al. [49] and Kenyon et al. [50] showed promising results in renal and pancreatic islet transplantation, respectively, following the administration of humanized CD154-specific monoclonal antibody in rhesus monkeys with acute rejection-

free and prolongation of graft survival. However, in the following year, Kawai et al. [51] documented a high incidence of thromboembolic complications after the use of monoclonal antibody against CD40L in monkeys. Nevertheless, further studies and pre-clinical evaluation of CD40-CD40L pathway blockade in conjunction with tolDC should be explored, including use of anti-CD40 prior to its application in patients.

A further clinically applicable question concerning tolDC therapy is the route of DC administration, since it could promote different outcomes. Giannoukakis et al. [36] have reported injection of autologous NF-κB-inhibited DC intradermally in the abdominal wall overlying the anatomic location of the pancreas in type-1 diabetic patients. As previously described by our group, in the mouse model, systemic (intravenous) administration of RAPA-DC was successful in significantly prolonging alloAg-specific heart graft survival [10]. Macatangay et al. found no difference between subcutaneous and intravenous administration of autologous monocyte-derived DC loaded with HIV-1 peptides delivered to subjects with chronic HIV-1 infection on anti-retroviral therapy [52]. Since CCR7 and CD62L expression on RAPA-DC is not affected by mTOR inhibition, this may allow the cells to traffic normally to secondary lymphoid tissues, where their immunoregulatory function is mediated [10,26,30].

How studies of human RAPA-DC relate to others' work

In recent years, tolDC protocols have offered a potential therapeutic tool in solid organ transplantation [1,2,53]. In order to compare different tolDC protocols, specific characteristics of the tolDC need to be analyzed. These include the phenotype, migration capability, cytokine production (in both immature and mature states), the ability to induce allogeneic T-cell proliferation, and the expansion/induction of Treg. For the purpose of generating tolDC, these can be manipulated *in vitro* with different immune modulators such as RAPA, dexamethasone, IL-10, TGF-β, or vitD3 [1,2,11-13].

In humans, the majority of the tolDC generated using the protocols mentioned above exhibit an immature to

Table 3 Generation of immature RAPA-DC

	Research	Clinical
Monocyte isolation	CD14 microbeads (AutoMACS® or CliniMACS®)	Elutriation or CD14 microbeads (CliniMACS®)
Culture conditions (7 days)	10% fetal calf serum AIM-V media 1000 U/mL rhIL-4 1000 U/mL GM-CSF 10 ng/mL Rapamycin Cell culture plates	10% human serum AIM-V media (c-GMP grade) 1000 U/mL rhIL-4 (c-GMP grade) 1000 U/mL GM-CSF (c-GMP grade) 10 ng/mL Rapamycin (c-GMP grade) Cell culture plates or Aastrom Replicell system

semi-mature cell surface phenotype, with low to intermediate expression of MHC II, CD86, CD83, and B7-H1. RAPA-DC and TGF- β -DC have a higher migration capability in response to CCL19 and CCL21 *in vitro* when compared to IL-10- and VitD3-DC, with higher expression of CCR7. Interestingly, all tolDC (dexamethasone-, IL-10-, RAPA-, TGF- β -, and VitD3-DC) exhibit diminished production of IL-23 when compared to mature untreated DC, whereas IL-10-DC and dexamethasone-DC were the only population to show increased production of IL-10 [11-13]. However, the ability of tolDC to suppress T-cell proliferation in humans is variable. IL-10-, TGF- β -, and VitD3-DC can each suppress T-cell proliferation [11-13]. Contradictory effects of RAPA on DC in culture have been reported. We have shown allo-PBMC hyporesponsiveness in MLR when stimulated with RAPA-DC [11]. Naranjo-Gomez et al. [12] have also shown RAPA-DC suppression of T-cell proliferation in CFSE-MLR on the other hand, Boks et al. [13] did not find RAPA-DC to be suppressive in MLR and Haidinger et al. found an allostimulatory effect of RAPA on DC (augmentation of IL-12, CD86, IL-1 β , and Ag presentation) [28]. Interestingly, Naranjo-Gomez et al. have shown, in humans, as we reported previously in mice, that RAPA-DC are capable of significantly sparing/expanding Treg, which suppress effector T-cell allo-reactivity [11,12].

Other immunosuppressive cell types of myeloid origin, such as macrophages [54], myeloid-derived suppressor cells generated in the presence of PGE2 [55], and mesenchymal stem cells [56], have been introduced recently to the transplant field, with suppressive properties that may be suitable for clinical use. DC treated with different cytokines and/or IS agents, macrophages, and mesenchymal stem cells mentioned above are being studied currently by The ONE Study, a multinational clinical assessment of immunomodulatory cell therapy in renal transplantation [57-59].

Conclusions

In an effort to reduce the adverse side effects of chronic IS after organ transplantation, tolDC protocols have been used to generate ‘negative cellular vaccines’ with potential therapeutic applicability. We have standardized culture conditions to generate human monocyte-derived RAPA-DC that exhibit tolerogenic characteristics, including a more immature phenotype when compared to control untreated DC. However, human RAPA-DC are not fully resistant to maturation, but can induce Treg, and have potential migratory capacity to secondary lymphoid tissue (spleen and lymph nodes). Although RAPA-DC exhibit unique immunoregulatory properties, the immediate clinical implementation of RAPA-DC is complicated by a dysregulation of pro- vs. anti-inflammatory cytokine production, particularly IL-12p70

and IL-10. However, methods to prevent increased IL-12p70 production by RAPA-DC (such as use of lithium chloride [11] or sanglifehrin A [60]) have been identified. Likewise, IL-10 could be delivered with vaccination to offset the reduced capacity of RAPA-DC to make IL-10. Further insights into how mTOR regulates DC cytokine production are critical for development of improved ‘negative’ and ‘positive’ cellular vaccines in general, and to begin to translate these technologies to the bedside.

Abbreviations

Ag: Antigen; CTLA4-Ig: Cytotoxic T lymphocyte Ag-4 immunoglobulin; DC: Dendritic cells; GM-CSF: Granulocyte macrophage colony stimulating factor; GSK3: Glycogen synthase kinase 3; IL: Interleukin; IS: Immunosuppression; LPS: Lipopolysaccharide; MLR: Mixed leukocyte reaction; mTOR: Mammalian target of rapamycin; PBMC: Peripheral blood mononuclear cells; PGE2: Prostaglandin E2; RAPA: Rapamycin; TGF β 1: Transforming growth factor β 1; TLR: Toll-like receptors; TolDC: Tolerogenic dendritic cells; Treg: Regulatory T cells; VEGF: Vascular endothelial growth factor.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CM wrote the manuscript. HT, DM, and AWT participated in editing the final manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by grant NIH-RO1 AI 067541-NAID (PI: AW Thomson).

Received: 6 June 2012 Accepted: 4 September 2012

Published: 28 September 2012

References

1. Ezzelarab M, Thomson AW: Tolerogenic dendritic cells and their role in transplantation. *Semin Immunol* 2011, **23**:252–263.
2. Morelli AE, Thomson AW: Tolerogenic dendritic cells and the quest for transplant tolerance. *Nat Rev Immunol* 2007, **7**:610–621.
3. Libby P, Pober JS: Chronic rejection. *Immunity* 2001, **14**:387–397.
4. Banchereau J, Steinman RM: Dendritic cells and the control of immunity. *Nature* 1998, **392**:245–252.
5. Morelli AE, Thomson AW: Dendritic cells: regulators of alloimmunity and opportunities for tolerance induction. *Immunol Rev* 2003, **196**:125–146.
6. Steinman RM, Hawiger D, Nussenweig MC: Tolerogenic dendritic cells. *Annu Rev Immunol* 2003, **21**:685–711.
7. Horibe EK, Sacks J, Unadkat J, Raimondi G, Wang Z, Ikeguchi R, Marsteller D, Ferreira LM, Thomson AW, Lee WP, Feili-Hariri M: Rapamycin-conditioned, alloantigen-pulsed dendritic cells promote indefinite survival of vascularized skin allografts in association with T regulatory cell expansion. *Transpl Immunol* 2008, **18**:307–318.
8. Mirenda V, Berton I, Read J, Cook T, Smith J, Dorling A, Lechner RI: Modified dendritic cells coexpressing self and allogeneic major histocompatibility complex molecules: an efficient way to induce indirect pathway regulation. *J Am Soc Nephrol* 2004, **15**:987–997.
9. Lutz MB, Suri RM, Niimi M, Ogilvie AL, Kukutsch NA, Rossner S, Schuler G, Austyn JM: Immature dendritic cells generated with low doses of GM-CSF in the absence of IL-4 are maturation resistant and prolong allograft survival *in vivo*. *Eur J Immunol* 2000, **30**:1813–1822.
10. Taner T, Hackstein H, Wang Z, Morelli AE, Thomson AW: Rapamycin-treated, alloantigen-pulsed host dendritic cells induce ag-specific T cell regulation and prolong graft survival. *Am J Transplant* 2005, **5**:228–236.
11. Turnquist HR, Cardinal J, Macedo C, Rosborough BR, Sumpter TL, Geller DA, Metes D, Thomson AW: mTOR and GSK-3 shape the CD4+ T-cell stimulatory and differentiation capacity of myeloid DCs after exposure to LPS. *Blood* 2010, **115**:4758–4769.

12. Naranjo-Gomez M, Raich-Regue D, Onate C, Grau-Lopez L, Ramo-Tello C, Pujol-Borrell R, Martinez-Caceres E, Borras FE: Comparative study of clinical grade human tolerogenic dendritic cells. *J Transl Med* 2011, **9**:89.
13. Boks MA, Kager-Groenland JR, Haasjes MS, Zwaginga JJ, van Ham SM, ten Brinke A: IL-10-generated tolerogenic dendritic cells are optimal for functional regulatory T cell induction—a comparative study of human clinical-applicable DC. *Clin Immunol* 2012, **142**:332–342.
14. Sehgal SN: Rapamune (RAPA, rapamycin, sirolimus): mechanism of action immunosuppressive effect results from blockade of signal transduction and inhibition of cell cycle progression. *Clin Biochem* 1998, **31**:335–340.
15. Hay N, Sonenberg N: Upstream and downstream of mTOR. *Genes Dev* 2004, **18**:1926–1945.
16. Wullschleger S, Loewith R, Hall MN: TOR signaling in growth and metabolism. *Cell* 2006, **124**:471–484.
17. Gao W, Lu Y, El Essawy B, Oukka M, Kuchroo VK, Strom TB: Contrasting effects of cyclosporine and rapamycin in de novo generation of alloantigen-specific regulatory T cells. *Am J Transplant* 2007, **7**:1722–1732.
18. Hackstein H, Taner T, Zahorchak AF, Morelli AE, Logar AJ, Gessner A, Thomson AW: Rapamycin inhibits IL-4-induced dendritic cell maturation in vitro and dendritic cell mobilization and function in vivo. *Blood* 2003, **101**:4457–4463.
19. Monti P, Mercalli A, Leone BE, Valerio DC, Allavena P, Piemonti L: Rapamycin impairs antigen uptake of human dendritic cells. *Transplantation* 2003, **75**:137–145.
20. Woltman AM, van der Kooij SW, Coffer PJ, Offringa R, Daha MR, van Kooten C: Rapamycin specifically interferes with GM-CSF signaling in human dendritic cells, leading to apoptosis via increased p27KIP1 expression. *Blood* 2003, **101**:1439–1445.
21. Ikeguchi R, Sacks JM, Unadkat JV, Solari M, Horibe EK, Thomson AW, Lee AW, Feili-Hariri M: Long-term survival of limb allografts induced by pharmacologically conditioned, donor alloantigen-pulsed dendritic cells without maintenance immunosuppression. *Transplantation* 2008, **85**:237–246.
22. Fischer RT, Turnquist HR, Wang Z, Beer-Stoltz D, Thomson AW: Rapamycin-conditioned, alloantigen-pulsed myeloid dendritic cells present donor MHC class I/peptide via the semi-direct pathway and inhibit survival of antigen-specific CD8(+) T cells in vitro and in vivo. *Transpl Immunol* 2011, **25**:20–26.
23. Battaglia M, Stabilini A, Roncarolo MG: Rapamycin selectively expands CD4 + CD25 + FoxP3+ regulatory T cells. *Blood* 2005, **105**:4743–4748.
24. Coenen JJ, Koenen HJ, van Rijssen E, Hilbrands LB, Joosten I: Rapamycin, and not cyclosporin A, preserves the highly suppressive CD27+ subset of human CD4 + CD25+ regulatory T cells. *Blood* 2006, **107**:1018–1023.
25. Turnquist HR, Raimondi G, Zahorchak AF, Fischer RT, Wang Z, Thomson AW: Rapamycin-conditioned dendritic cells are poor stimulators of allogeneic CD4+ T cells, but enrich for antigen-specific Foxp3+ T regulatory cells and promote organ transplant tolerance. *J Immunol* 2007, **178**:7018–7031.
26. Reichardt W, Durr C, von Elverfeldt D, Juttner E, Gerlach UV, Yamada M, Smith B, Negrin RS, Zeiser R: Impact of mammalian target of rapamycin inhibition on lymphoid homing and tolerogenic function of nanoparticle-labeled dendritic cells following allogeneic hematopoietic cell transplantation. *J Immunol* 2008, **181**:4770–4779.
27. Pothoven KL, Khermandan T, Yang Q, Houlihan JL, Zhang H, Degutes M, Miller SD, Luo X: Rapamycin-conditioned donor dendritic cells differentiate CD4CD25Foxp3 T cells in vitro with TGF-beta1 for islet transplantation. *Am J Transplant* 2010, **10**:1774–1784.
28. Haidinger M, Poglitsch M, Geyreger R, Kasturi S, Zeyda M, Zlabinger GJ, Pulendran B, Horl WH, Saemann MD, Weichhart T: A versatile role of mammalian target of rapamycin in human dendritic cell function and differentiation. *J Immunol* 2010, **185**:3919–3931.
29. Ohtani M, Nagai S, Kondo S, Mizuno S, Nakamura K, Tanabe M, Takeuchi T, Matsuda S, Koyasu S: Mammalian target of rapamycin and glycogen synthase kinase 3 differentially regulate lipopolysaccharide-induced interleukin-12 production in dendritic cells. *Blood* 2008, **112**:635–643.
30. Sordi V, Bianchi G, Buracchi C, Mercalli A, Marchesi F, D'Amico G, Yang CH, Luini W, Vecchi A, Mantovani A, Allavena P, Piemonti L: Differential effects of immunosuppressive drugs on chemokine receptor CCR7 in human monocyte-derived dendritic cells: selective upregulation by rapamycin. *Transplantation* 2006, **82**:826–834.
31. Chauvabel D, Banchereau J: Dendritic cells, therapeutic vectors of immunity and tolerance. *Am J Transplant* 2005, **5**:205–206.
32. Weichhart T, Saemann MD: T helper cell differentiation: understanding the needs of hierarchy. *Immunity* 2010, **32**:727–729.
33. Cranmer LD, Trevor KT, Hersh EM: Clinical applications of dendritic cell vaccination in the treatment of cancer. *Cancer Immunol Immunother* 2004, **53**:275–306.
34. Lesterhuis WJ, de Vries IJ, Adema GJ, Punt CJ: Dendritic cell-based vaccines in cancer immunotherapy: an update on clinical and immunological results. *Ann Oncol* 2004, Suppl 4:iv145–iv151.
35. Whiteside TL, Piazza P, Reiter A, Stanton J, Connolly NC, Rinaldo CR Jr, Riddler SA: Production of a dendritic cell-based vaccine containing inactivated autologous virus for therapy of patients with chronic human immunodeficiency virus type 1 infection. *Clin Vaccine Immunol* 2009, **16**:233–240.
36. Giannoukakis N, Phillips B, Finegold D, Harnaha J, Trucco M: Phase I (safety) study of autologous tolerogenic dendritic cells in type 1 diabetic patients. *Diabetes Care* 2011, **34**:2026–2032.
37. Tanriver Y, Ratnasothy K, Bucy RP, Lombardi G, Lechner R: Targeting MHC class I monomers to dendritic cells inhibits the indirect pathway of allorecognition and the production of IgG alloantibodies leading to long-term allograft survival. *J Immunol* 2010, **184**:1757–1764.
38. Fehervari Z, Sakaguchi S: Control of Foxp3+ CD25 + CD4+ regulatory cell activation and function by dendritic cells. *Int Immunopharmacol* 2004, **16**:1769–1780.
39. Darrasse-Jeze G, Deroubaix S, Mouquet H, Victora GD, Eisenreich T, Yao KH, Masilamani RF, Dustin ML, Rudensky A, Liu K, Nussenzweig MC: Feedback control of regulatory T cell homeostasis by dendritic cells in vivo. *J Exp Med* 2009, **206**:1853–1862.
40. Macke L, Garritsen HS, Meyring W, Hannig H, Pagelow U, Wormann B, Piechaczek C, Geffers R, Rohde M, Lindenmaier W, Dittmar KE: Evaluating maturation and genetic modification of human dendritic cells in a new polyolefin cell culture bag system. *Transfusion* 2010, **50**:843–855.
41. Van Diessche A, Van de Velde AL, Nijs G, Braeckman T, Stein B, De Vries JM, Berneman ZN, Van Tendeloo VF: Clinical-grade manufacturing of autologous mature mRNA-electroporated dendritic cells and safety testing in acute myeloid leukemia patients in a phase I dose-escalation clinical trial. *Cytotherapy* 2009, **11**:653–668.
42. Ferrer IR, Wagener ME, Robertson JM, Turner AP, Araki K, Ahmed R, Kirk AD, Larsen CP, Ford ML: Cutting edge: Rapamycin augments pathogen-specific but not graft-reactive CD8+ T cell responses. *J Immunol* 2010, **185**:2004–2008.
43. Gilboa E: DC-based cancer vaccines. *J Clin Invest* 2007, **117**:1195–1203.
44. Han TH, Jin P, Ren J, Slezak S, Marincola FM, Sronecek DF: Evaluation of 3 clinical dendritic cell maturation protocols containing lipopolysaccharide and interferon-gamma. *J Immunother* 2009, **32**:399–407.
45. Mailliard RB, Wankowicz-Kalinska A, Cai Q, Wesa A, Hilkens CM, Kapsenberg ML, Kirkwood JM, Storkus WJ, Kalinski P: alpha-type-1 polarized dendritic cells: a novel immunization tool with optimized CTL-inducing activity. *Cancer Res* 2004, **64**:5934–5937.
46. Davies JK, Barbon CM, Voskerchian A, Nadler LM, Guinan EC: Ex vivo alloantigenization with belatacept: a strategy to selectively modulate alloresponses after transplantation. *Cell Transplant* 2012. doi:10.3727/096368912X637479. Apr 10 [Epub ahead of print].
47. Archeacon P, Dixon C, Belen O, Albrecht R, Meyer J: Summary of the US FDA approval of belatacept. *Am J Transplant* 2012, **12**:554–562.
48. Lu L, Li W, Fu F, Chambers FG, Qian S, Fung JJ, Thomson AW: Blockade of the CD40-CD40 ligand pathway potentiates the capacity of donor-derived dendritic cell progenitors to induce long-term cardiac allograft survival. *Transplantation* 1997, **64**:1808–1815.
49. Kirk AD, Burkly LC, Batty DS, Baumgartner RE, Berning JD, Buchanan K, Fechner JH Jr, Germond RL, Kampen RL, Patterson NB, Swanson SJ, Tadaki DK, TenHoor CN, White L, Knechtle SJ, Harlan DM: Treatment with humanized monoclonal antibody against CD154 prevents acute renal allograft rejection in nonhuman primates. *Nat Med* 1999, **5**:686–693.
50. Kenyon NS, Chatzipetrou M, Masetti M, Ranuncoli A, Oliveira M, Wagner JL, Kirk AD, Harlan DM, Burkly LC, Ricordi C: Long-term survival and function of intrahepatic islet allografts in rhesus monkeys treated with humanized anti-CD154. *Proc Natl Acad Sci USA* 1999, **96**:8132–8137.
51. Kawai T, Andrews D, Colvin RB, Sachs DH, Cosimi AB: Thromboembolic complications after treatment with monoclonal antibody against CD40 ligand. *Nat Med* 2000, **6**:114.

52. Macatangay BJ, Szajnik ME, Whiteside TL, Riddler SA, Rinaldo CR: Regulatory T cell suppression of Gag-specific CD8 T cell polyfunctional response after therapeutic vaccination of HIV-1-infected patients on ART. *PLoS One* 2010, **5**:e9852.
53. Bériou G, Moreau A, Cuturi MC: Tolerogenic dendritic cells: applications for solid organ transplantation. *Curr Opin Organ Transplant* 2012, **17**:42–47.
54. Hutchinson JA, Riquelme P, Sawitzki B, Tomiuk S, Miqueu P, Zuharya M, Oberg HH, Pascher A, Lutzen U, Janssen U, Broichhausen C, Renders L, Thaiss F, Scheuermann E, Henze E, Volk HD, Chatenoud L, Lechner RI, Wood KJ, Kabelitz D, Schlitt HJ, Geissler EK, Fandrich F: Cutting Edge: Immunological consequences and trafficking of human regulatory macrophages administered to renal transplant recipients. *J Immunol* 2011, **187**:2072–2078.
55. Obermajer N, Muthuswamy R, Lesnock J, Edwards RP, Kalinski P: Positive feedback between PGE2 and COX2 redirects the differentiation of human dendritic cells toward stable myeloid-derived suppressor cells. *Blood* 2011, **118**:5498–5505.
56. Eggenhofer E, Steinmann JF, Renner P, Slowik P, Piso P, Geissler EK, Schlitt HJ, Dahlke MH, Popp FC: Mesenchymal stem cells together with mycophenolate mofetil inhibit antigen presenting cell and T cell infiltration into allogeneic heart grafts. *Transpl Immunol* 2011, **24**:157–163.
57. Amodo G, Gregori S: Human tolerogenic DC-10: perspectives for clinical Applications. *Transplantation Res* 2012, **1**:14.
58. Obermajer N, Kalinski P: Generation of myeloid-derived suppressor cells using prostaglandin E₂. *Transplantation Res* 2012, **1**:15.
59. Riquelme P, Geissler EK, Hutchinson JA: Alternative Approaches to Myeloid Suppressor Cell Therapy in Transplantation: Comparing Regulatory Macrophages to Tolerogenic DCs and MDSCs. *Transplantation Res* 2012, **1**:17.
60. Hackstein H, Steinschulte C, Fiedel S, Eisele A, Rathke V, Stadlbauer T, Taner T, Thomson AW, Tillmanns H, Bein G, Holschermann H: Sanglifehrin a blocks key dendritic cell functions in vivo and promotes long-term allograft survival together with low-dose CsA. *Am J Transplant* 2007, **7**:789–798.

doi:10.1186/2047-1440-1-16

Cite this article as: Macedo et al.: Immunoregulatory properties of rapamycin-conditioned monocyte-derived dendritic cells and their role in transplantation. *Transplantation Research* 2012 1:16.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

