

CORRECTION

Correction: Channel Properties of Na_x Expressed in Neurons

The *PLOS ONE* Staff

[S4 Fig](#) is incorrect. Please view the correct [S4 Fig](#) below. The publisher apologizes for the error.

Supporting Information

S4 Fig. Immunohistochemical staining of rat and mouse brains with anti- Na_x antibodies.

Immunohistochemical staining of the coronal sections of rat and mouse brains, containing the median preoptic nucleus (MnPO) (A) and median eminence (B) with anti-rat Na_x [12] and anti-m Na_x antibodies. Immunohistochemical staining was performed as described in S5 File. Neither rat nor mouse MnPO was negative for Na_x (A). On the other hand, the median eminence was clearly stained with both antibodies (B). AC, anterior commissure. Scale bars, 200 μm .

(TIF)

Reference

1. Matsumoto M, Hiyama TY, Kuboyama K, Suzuki R, Fujikawa A, Noda M (2015) Channel Properties of Na_x Expressed in Neurons. *PLoS ONE* 10(5): e0126109. doi:[10.1371/journal.pone.0126109](https://doi.org/10.1371/journal.pone.0126109) PMID: [25961826](https://pubmed.ncbi.nlm.nih.gov/25961826/)



OPEN ACCESS

Citation: The *PLOS ONE* Staff (2015) Correction: Channel Properties of Na_x Expressed in Neurons. *PLoS ONE* 10(6): e0130107. doi:[10.1371/journal.pone.0130107](https://doi.org/10.1371/journal.pone.0130107)

Published: June 4, 2015

Copyright: © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.