



Research Article

Expression of hsv1-miR-H18 and hsv2-miR-H9 as a field defect marker for detecting prostate cancer



Young Joon Byun ^{a,☆}, Ho Won Kang ^{a,b,☆}, Xuan-Mei Piao ^a, Chuang-Ming Zheng ^a,
Sung-Kwon Moon ^c, Yung Hyun Choi ^d, Won Tae Kim ^{a,b}, Sang-Cheol Lee ^{a,b},
Seok Joong Yun ^{a,b,*}, Wun-Jae Kim ^{a,e,**}

^a Department of Urology, College of Medicine, Chungbuk National University, Cheongju, Korea

^b Department of Urology, Chungbuk National University Hospital, Cheongju, Korea

^c Department of Food Science and Technology, Chung-Ang University, Ansong, 456-756, Korea

^d Department of Biochemistry, Donggwi University College of Oriental Medicine, Busan, Korea

^e Institute of Urotech, Cheongju, 28644, Korea

ARTICLE INFO

Article history:

Received 27 October 2021

Received in revised form

24 November 2021

Accepted 29 November 2021

Available online 3 December 2021

Keywords:

Biomarkers

Diagnosis

microRNAs

Prostate neoplasms

Prostate-specific antigen

ABSTRACT

Background: Prostate-specific antigen (PSA) is a marker of prostate cancer (PCa), although its efficacy as a diagnostic marker remains controversial. A high false-positive rate leads to repeat biopsy in approximately 70% of patients, which may not be necessary. Epigenetic biomarkers of field cancerization have been investigated widely as promising tools for the diagnosis of patients with suspected tumors. In the current study, we examined the diagnostic value of two microRNA (miRNA) candidates, hsv1-miR-H18 and hsv2-miR-H9, using formalin-fixed paraffin-embedded (FFPE) tissues from patients with PCa or benign prostate hyperplasia (BPH) (as controls) to determine the usefulness of these markers for detecting the presence of cancer.

Methods: Expression of hsv1-miR-H18 and hsv2-miR-H9 in 201 FFPE tissues, including 52 primary tumors, 73 surrounding noncancerous tissues, and 90 BPH nontumor controls was examined by real-time PCR.

Results: Expression of hsv1-miR-H18 and hsv2-miR-H9 was significantly higher in primary tumors from PCa patients than in BPH controls ($P < 0.0001$). In patients within the PSA gray zone, the two viral miRNAs could distinguish PCa from controls with appropriate sensitivity and specificity. Expression of the two miRNAs did not differ between primary tumors and noncancerous surrounding tissues.

Conclusions: The viral miRNAs hsv1-miR-H18 and hsv2-miR-H9 may be associated with field cancerization of PCa and could be promising supplemental biomarkers to the PSA assay to decrease the rate of unnecessary biopsy, particularly in patients within the PSA gray zone.

© 2022 Asian Pacific Prostate Society. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Prostate-specific antigen (PSA) screening is commonly used to detect early prostate cancer (PCa). However, this assay remains controversial because of its limited diagnostic performance.^{1–3} The false-positive rate is particularly high in patients with serum PSA levels of 3.0–10.0 ng/mL, which is known as the 'PSA gray zone'.³ Approximately 70% of those patients undergo repeat biopsy and are often overdiagnosed and overtreated.⁴ Thus, identifying a more accurate biomarker to PSA is an urgent need.

The concept of field cancerization (also known as field defect) was first described by Slaughter et al. in 1953 to explain the development of precancerous changes in histologically normal

* Corresponding author. Department of Urology, College of Medicine, Chungbuk National University, Chungdae-ro 1, Seowon-Gu, Cheongju, Chungbuk 28644, Korea.

** Corresponding author. Department of Urology, College of Medicine, Chungbuk National University, Institute of Urotech, Chungdae-ro 1, Seowon-Gu, Cheongju, Chungbuk 28644, Korea.

E-mail addresses: sjyun@chungbuk.ac.kr (S.J. Yun), wjkim@chungbuk.ac.kr (W.-J. Kim).

☆ Contributed equally to this work.

Table 1
Clinical characteristics of the study subjects

Variable	BPH	PCa	p-value ^a
No. of patients	90	125	
Age (y)	70.00 ± 6.71	69.00 ± 5.41	0.163
PSA (ng/mL)	2.94 ± 8.36	11.42 ± 599.14	<0.001
Operation			
TURP	90 (100.0)	23 (18.4)	
Radical prostatectomy	-	102 (81.6)	
Gleason score			
6	-	2 (1.6)	
7 (3 + 4)	-	43 (34.4)	
7 (4 + 3)	-	35 (28.0)	
≥8	-	45 (36.0)	
TNM stage		TURP	RP
Unknown	-	3 (2)	-
T2≤	-	1 (1)	65 (52)
T3	-	19 (15)	25 (20)
T4 or metastasis	-	-	12 (10)

Values are presented as numbers only, mean ± standard deviation, or number (%).

BPH, benign prostate hyperplasia; PCa, prostate cancer; PSA, prostate-specific antigen; TURP, transurethral resection of the prostate.

^a) All p-values were obtained by the Mann–Whitney *U*-test for comparisons between formalin-fixed paraffin-embedded from non-tumor controls and PCa patients in the study.

tissues.⁵ It has been reviewed in various types of cancer, including PCa, as paramount evidence of the multifocality of tumors.⁶ Transrectal prostate biopsy is prone to sampling error, which can lead to spurious negative findings, particularly in cases within the PSA gray zone. Therefore, examining preneoplastic molecular alterations has become a promising strategy for the selection of appropriate procedures.^{7,8}

MicroRNAs (miRNAs) are small, nonprotein-encoding RNAs of approximately 20 nucleotides in length that regulate gene expression. Many miRNAs have been suggested as biomarkers because of their involvement in biological processes, including field cancerization.^{9–11} miRNA profiling can be used to evaluate the likelihood of tumor development and to decrease the rate of histological false-negative results.^{12,13}

In the current study, we measured the expression of two miRNAs, hsv1-miR-H18 and hsv2-miR-H9, as relevant epigenetic markers for PCa in formalin-fixed paraffin-embedded (FFPE) tissues to determine their ability to distinguish tumors from nontumor tissues.

2. Materials and methods

2.1. Cases

A total of 215 FFPE tissues from 125 PCa patients and 90 benign prostate hyperplasia (BPH) controls were obtained from Chungbuk National University Hospital between May 2003 and June 2015 (approval number: 2012-02-017-001) (Table 1). Primary tumor tissues and surrounding noncancerous tissues were collected from PCa patients who underwent radical prostatectomy (RP) or palliative transurethral resection of the prostate (TURP). Control tissues were harvested from patients with BPH who underwent TURP. Fresh tissues were formalin-fixed for >24 h, embedded in paraffin blocks, and stored at room temperature until use. Cases with a serum PSA concentration of 3–10 ng/mL were considered to be in the PSA gray zone. Gleason grades were assigned to samples obtained from TURP or RP. Tumor stage was estimated from biospecimens harvested by RP or from computed tomography, magnetic resonance imaging, or bone scans. The study methodologies conformed to the standards set by the Declaration of Helsinki.

2.2. miRNA microarray analysis

The RecoverAll Total Nucleic Acid Isolation Kit (Life Technologies, Carlsbad, CA, USA) was used to extract total RNA from each specimen. The amount and integrity of extracted RNA were examined using the RNA 6000 Pico Chip Kit (Agilent Technologies, Santa Clara, CA, USA) and the Agilent 2100 Bioanalyzer. The Agilent Human miRNA Microarray Release 16.0 platform, which contains 1,205 human and 144 viral miRNAs, was used for miRNA profiling.¹⁴ The protocol used for generating the microarray gene expression datasets have been described previously.¹⁵

2.3. Purification of miRNAs from FFPE tissues

FFPE tissues were cut into 5 µm thick sections with 10–15 slides and placed in 1.5 mL Eppendorf Safe-lock micro-centrifuge tubes and stored at –70°C until use. For extracting miRNA from FFPE sections, a NucleoSpin Total RNA FFPE XS kit (Macherey-Nagel, Düren, Germany) was used according to the manufacturer's protocol.

2.4. Synthesis of complementary (c)DNA from miRNA

The concentration of RNA isolated from FFPE tissues was measured using the Quant-IT RiboGreen RNA Reagent and Kit (Invitrogen, Grand Island, NY, USA). The Mir-X™ miRNA First Strand cDNA Synthesis Kit (TAKARA BIO, Otsu, Japan) was used to synthesize miRNA-specific cDNA according to the manufacturer's protocol.

2.5. Amplification of miRNAs by real-time PCR (RT-PCR)

For quantifying miRNA expression, RT-PCR amplification was performed using a Rotor-Gene Q (Qiagen, Valencia, CA, USA) apparatus and SYBR Premix EX Taq (TAKARA BIO, Otsu, Japan). Reactions were performed in micro-reaction tubes (Corbett Research Mortlake, Australia) in a final volume of 10 µL. Standard curves were generated using chemically synthesized RNA oligonucleotides (Cosmogenetech, Seoul, Korea) corresponding to the target miRNAs. The standard curves ranged from 2.25×10^5 to 2.25×10^8 copies. All samples were run in triplicate, and RT-PCR

conditions followed the manufacturer's protocol. Rotor-Gene Q software 2.3.1.49 was used to capture and analyze spectral data.

2.6. Statistical analysis

Receiver operating characteristic (ROC) curves were generated to estimate the optimal cut-off point yielding the highest combined sensitivity and specificity of the two viral miRNAs in FFPE tissues from patients with PCa and from BPH controls. The expression levels of candidate miRNAs in PCa, in surrounding noncancerous tissues, and in BPH controls were examined using the Kruskal–Wallis H and Mann–Whitney U tests. Statistical analysis was performed using IBM SPSS version 24.0 (IBM, Armonk, NY, USA) and GraphPad Prism 7 (GraphPad Software, San Diego, CA, USA). $P < 0.05$ was considered statistically significant.

3. Results

3.1. Expression levels of the two miRNAs

The expression levels of hsv1-miR-H18 and hsv2-miR-H9 were similar in PCa and surrounding noncancerous tissues, and

significantly higher than in BPH controls ($P < 0.0001$) (Fig. 1A and C). For cases within the 'PSA gray zone', expression of the two viral miRNAs was higher in PCa and surrounding noncancerous tissues than in BPH controls ($P < 0.0001$ and $P = 0.005$, respectively; Fig. 1B and D). These findings indicate that hsv1-miR-H18 and hsv2-miR-H9 are associated with field cancerization of cells surrounding tumors.

3.2. Diagnostic performance of the two miRNAs in FFPE samples from PCa, surrounding noncancerous tissue, and BPH controls

ROC analysis was performed to determine whether hsv1-miR-H18 and hsv2-miR-H9 could distinguish patients with tumors from those with BPH. The areas under the curve (AUCs) for the two candidates were 0.800 and 0.706, with a sensitivity of 76.9% and 80.8%, and specificity of 81.1% and 55.6%, respectively (Figs. 2A and 3A). To further examine the diagnostic power of the markers, we compared surrounding noncancerous tissues with BPH tissues and found that the AUCs of the target miRNAs were 0.840 and 0.724, respectively, with a sensitivity of 89.0% and 76.7% and specificity of 75.6% and 60.0%, respectively (Figs. 2B and 3B). In patients within the PSA gray zone, the miRNAs showed improved diagnostic




Fig. 1. Expression of hsv1-miR-H18 and hsv2-miR-H9 in FFPE tissues from BPH controls, surrounding noncancerous tissues, and primary tumors of PCa patients. Levels of the two viral miRNAs were higher in FFPE tissues from PCa patients than from controls and showed similar expression patterns between surrounding noncancerous tissues and primary PCa tumors. BPH, benign prostate hyperplasia; FFPE, formalin-fixed paraffin-embedded; PCa, prostate cancer; PSA, prostate-specific antigen. P-values were determined by the Mann-Whitney U-test. * $P < 0.05$, ** $P < 0.01$, and **** $P < 0.0001$.




Fig. 2. Receiver operating characteristic curves of hsv1-miR-H18 for distinguishing PCa patients from non-tumor controls. (A) and (B) for all patients. (C) and (D) for patients within the PSA gray zone (3.0–10.0 ng/mL). AUC, area under the curve; BPH, benign prostate hyperplasia; PCa prostate cancer; PSA, prostate-specific antigen. *P-value was determined by the Z-score.




Fig. 3. Receiver operating characteristic curves of hsv2-miR-H9 for distinguishing PCa patients from non-tumor controls. (A) and (B) for all patients. (C) and (D) for patients within the PSA gray zone (3.0–10.0 ng/mL). AUC, area under the curve; BPH, benign prostate hyperplasia; PCa prostate cancer; PSA, prostate-specific antigen. *P-value was determined by the Z-score.

performance. When comparing PCa with BPH controls, the sensitivity and specificity of hsv1-miR-H18/hsv2-miR-H9 were 92.3%/84.6% and 86.7%/53.3%, respectively (AUC = 0.915 and 0.738; and $P < 0.0001$ and $P < 0.05$; respectively) (Figs. 2C and 3C). Similar results were obtained when comparing surrounding noncancerous tissues with BPH controls, which showed sensitivity and specificity for hsv1-miR-H18/hsv2-miR-H9 of 90.6%/87.5% and 80.0%/56.7%, respectively (AUC = 0.886 and 0.704; and $P < 0.0001$ and < 0.01 ; respectively) (Figs. 2D and 3D). These results suggest that hsv1-miR-H18 and hsv2-miR-H9 can distinguish patients with tumors from those with BPH, thereby preventing miss-diagnosis by biopsy sampling error.

4. Discussion

PSA is broadly used as a standard biomarker for the diagnosis of PCa. However, PSA-based PCa screening is limited by poor specificity and sensitivity, especially in patients within the PSA gray zone (3–10 ng/mL).^{1,16,17} Patients within the PSA gray zone often undergo repeat biopsy, although the detection rate after this procedure is $< 30\%$.^{1–4,18,19} Therefore, approximately 70% of these patients are overdiagnosed that resulted in meaningless re-biopsies with no clinical benefits, underscoring the need to identify novel methods for early detection of PCa that can reduce the rate of unnecessary re-biopsy.

Recent studies identified biomarkers that increase the detection rate of PCa; however, there are several limitations. Prostate cancer antigen 3 (PCA3) is a common PCa biomarker, and increased expression of the *PCA3* gene is detected in the urine of PCa patients. However, the threshold value of PCA3 required to distinguish patients at risk of PCa from healthy controls remains unclear, and the cost-effectiveness of this assay was not determined yet, indicating that the performance of PCA3 for clinical practice applications is debatable.^{20,21}

The field defect or field cancerization is considered a source of valuable biomarkers for PCa.^{9,22} The basic concept of field cancerization, suggested by Slaughter, describes the development of cancer in multifocal areas.⁵ There are many studies analyzing field cancerization in head and neck, colorectal, bladder, and other cancers.^{23,24} In PCa, methylated *GSTP1* has been studied extensively as a tumor marker.²⁵ The ConfirmMDx is a quantitative methylation assay used to diagnose PCa. This assay quantifies the DNA methylation rates of three genes, *GSTP1*, *APC*, and *RASSF1*, to diagnose PCa.²⁶ However it has several limitations; for example, *APC* methylation can be affected by infection or inflammation of tissues, and the performance of this assay in the recent DOCUMENT study was not satisfactory.²⁷ Many genes have been studied and reported as promising diagnostic biomarkers for PCa; however, most are in the initial stages and require complex protocols that are impractical for clinical application, whereas others failed to show reproducible results in various tests.^{28,29}

Recently, many researchers have focused on bacteria and viruses that may associate with infection and inflammation causing PCa.^{30,31} Studies suggested that several viruses, including herpes types, encoded their own miRNAs, and they may affect carcinogenesis of the prostate and lead to the transformation of epithelial cells of prostate.^{32,33} Although, the exact function of viral miRNA is still unknown, diverse expression levels of viral miRNAs were identified in PCa when compared with BPH controls, indicating that viral miRNA could be a respectable risk factor for PCa.³⁴

Previously, we have identified the increased values of hsv1-miR-H18 and hsv2-miR-H9 that may be associated with PCa and suggested that these two viral miRNAs could be relevant diagnostic biomarkers for PCa to decrease biopsy burdens.¹⁶

Currently, we examined the expression values of hsv1-miR-H18 and hsv2-miR-H9 using FFPE tissues from PCa patients and BPH controls. For further investigation of two viral miRNAs in the biopsy, samples should be prepared as FFPE type, as FFPE biospecimen can be repeatedly used, which could provide molecular and pathological diagnosis at the same time. The hsv1-miR-H18 and hsv2-miR-H9 showed appropriate sensitivity and specificity, independently of histological outcomes, and showed advanced diagnostic performance in cases within the PSA gray zone. These results suggest that hsv1-miR-H18 and hsv2-miR-H9 are upregulated in prostate tumors and show potential as field defect markers for PCa diagnosis. However, a correlation between the two miRNAs and clinicopathological parameters of PCa was not observed (data not shown), which indicates that their ability to predict the aggressiveness of PCa is limited.

The results of this study indicate that hsv1-miR-H18 and hsv2-miR-H9 could serve as diagnostic biomarkers for PCa and may help to decrease the rate of unnecessary biopsy, especially in patients within the PSA gray zone. To apply this diagnosing strategy in clinical practice, validation of these miRNAs in biopsy specimens should be performed as the step for setting precise 'cut-off point' or 'cut-off ranges' to distinguish patients with high potential of PCa presents, then investigated in blinded cohort to determine whether they actually could reduce biopsy sampling errors.

Funding

This research was supported by the Osong Medical Innovation Foundation, funded by Chungcheongbuk-do (No.: 2020017001).

Author contribution statement

Research conception and design: Young Joon Byun, Seok Joong Yun, and Wun-Jae Kim.

Data acquisition: Young Joon Byun, Ho Won Kang, Xuan-Mei Piao, Chang-Ming Zheng.

Statistical analysis: Young Joon Byun.

Data analysis and interpretation: Young Joon Byun.

Drafting of the manuscript: Young Joon Byun.

Critical revision of manuscript: Ho Won Kang, Xuan-Mei Piao, Sung-Kwon Moon, Yung Hyun Choi, Won Tae Kim, Sang-Cheol Lee, and Seok Joong Yun.

Administrative, technical, or material support: Seok Joong Yun and Wun-Jae Kim.

Supervision: Seok Joong Yun and Wun-Jae Kim.

Approval of the final manuscript: All authors.

Ethics statement

The Ethics Committee of Chungbuk National University approved the protocol, and written informed consent was obtained from each patient. The collection and analysis of all samples were approved by the Institutional Review Board of Chungbuk National University (IRB approval number 2010-02-017-001), and informed consent was obtained from each patient.

Conflicts of interest

All authors have no conflict of interest to declare.

Acknowledgments

The biospecimens for this study were provided by Chungbuk National University Hospital and Kyungpook National University Hospital, members of the National Biobank of Korea, which is

supported by the Ministry of Health, Welfare, and Family Affairs. All samples derived from the National Biobank of Korea were obtained with informed consent under Institutional Review Board-approved protocols.

References

- Smith DS, Humphrey PA, Catalona WJ. The early detection of prostate carcinoma with prostate specific antigen: the Washington University experience. *Cancer* 1997;80:1852–6.
- Kim Y-H, Byun YJ, Kim WT, Jeong P, Yan C, Kang HW, et al. CDC6 mRNA expression is associated with the aggressiveness of prostate cancer. *J Korean Med Sci* 2018;33.
- Zuo W, Hiraoka Y. Prostate-specific antigen levels and density in the internal and external glands of the prostate in benign prostatic hyperplasia patients with normal or gray-zone PSA levels. *Urol Int* 2003;71:154–9.
- Byun YJ, Piao X-M, Jeong P, Kang HW, Seo SP, Moon S-K, et al. Urinary microRNA-1913 to microRNA-3659 expression ratio as a non-invasive diagnostic biomarker for prostate cancer. *Investig Clin Urol* 2021;62:340.
- Slaughter DP, Southwick HW, Smejkal W. "Field cancerization" in oral stratified squamous epithelium. Clinical implications of multicentric origin. *Cancer* 1953;6:963–8.
- Mehrotra J, Varde S, Wang H, Chiu H, Vargo J, Gray K, et al. Quantitative, spatial resolution of the epigenetic field effect in prostate cancer. *Prostate* 2008;68:152–60.
- Van Neste L, Bigley J, Toll A, Otto G, Clark J, Delrée P, et al. A tissue biopsy-based epigenetic multiplex PCR assay for prostate cancer detection. *BMC Urol* 2012;12:1–8.
- Babaian RJ. Extended field prostate biopsy enhances cancer detection. *Urology* 2000;55:453–6.
- Fabbri M, Calin GA. Epigenetics and miRNAs in human cancer. *Adv Genet* 2010;70:87–99.
- Nonn L, Ananthanarayanan V, Gann PH. Evidence for field cancerization of the prostate. *Prostate* 2009;69:1470–9.
- Ganci F, Sacconi A, Manciooco V, Sperduti I, Battaglia P, Covelto R, et al. MicroRNA expression as predictor of local recurrence risk in oral squamous cell carcinoma. *Head Neck* 2016;38:E189–97.
- Pereira AL, Magalhães L, Moreira FC, Reis-das-Mercês L, Vidal AF, Ribeiro-dos-Santos AM, et al. Epigenetic field cancerization in gastric cancer: microRNAs as promising biomarkers. *J Cancer* 2019;10:1560.
- Kanwal R, Gupta S. Epigenetic modifications in cancer. *Clin Genet* 2012;81:303–11.
- Wang H, Ach RA, Curry B. Direct and sensitive miRNA profiling from low-input total RNA. *RNA* 2007;13:151–9.
- Takeshita N, Hoshino I, Mori M, Akutsu Y, Hanari N, Yoneyama Y, et al. Serum microRNA expression profile: miR-1246 as a novel diagnostic and prognostic biomarker for oesophageal squamous cell carcinoma. *Br J Cancer* 2013;108:644–52.
- Yun SJ, Jeong P, Kang HW, Shinn HK, Kim Y-H, Yan C, et al. Increased expression of herpes virus-encoded hsv1-miR-H18 and hsv2-miR-H9-5p in cancer-containing prostate tissue compared to that in benign prostate hyperplasia tissue. *Int Neurourol J* 2016;20:122.
- Roobol MJ, Steyerberg EW, Kranse R, Wolters T, van den Bergh RC, Bangma CH, et al. A risk-based strategy improves prostate-specific antigen-driven detection of prostate cancer. *Eur Urol* 2010;57:79–85.
- Djavan B, Zlotta A, Remzi M, Ghawidel K, Basharkhah A, Schulman CC, et al. Optimal predictors of prostate cancer on repeat prostate biopsy: a prospective study of 1,051 men. *J Urol* 2000;163:1144–9.
- Partin AW, Brawer MK, Bartsch G, Horninger W, Taneja SS, Lepor H, et al. Complexed prostate specific antigen improves specificity for prostate cancer detection: results of a prospective multicenter clinical trial. *J Urol* 2003;170:1787–91.
- Nicholson A, Mahon J, Boland A, Beale S, Dwan K, Fleeman N, et al. The clinical effectiveness and cost-effectiveness of the PROGENSA® prostate cancer antigen 3 assay and the Prostate Health Index in the diagnosis of prostate cancer: a systematic review and economic evaluation. *Health Technol Assess (Winchester, England)* 2015;19. i-xxxi, 1.
- Ploussard G, de la Taille A. The role of prostate cancer antigen 3 (PCA3) in prostate cancer detection. *Expert Rev Anticancer Ther* 2018;18:1013–20.
- Truong M, Yang B, Livermore A, Wagner J, Weeratunga P, Huang W, et al. Using the epigenetic field defect to detect prostate cancer in biopsy negative patients. *J Urol* 2013;189:2335–41.
- Suzuki H, Watkins DN, Jair K-W, Schuebel KE, Markowitz SD, Chen WD, et al. Epigenetic inactivation of SFRP genes allows constitutive WNT signaling in colorectal cancer. *Nat Genet* 2004;36:417–22.
- Shen L, Kondo Y, Rosner GL, Xiao L, Hernandez NS, Vilaythong J, et al. MGMT promoter methylation and field defect in sporadic colorectal cancer. *J Natl Cancer Inst* 2005;97:1330–8.
- Stewart GD, Van Neste L, Delvenne P, Delrée P, Delga A, McNeill SA, et al. Clinical utility of an epigenetic assay to detect occult prostate cancer in histopathologically negative biopsies: results of the MATLOC study. *J Urol* 2013;189:1110–6.
- Narayan VM, Konety BR, Warlick C. Novel biomarkers for prostate cancer: An evidence-based review for use in clinical practice. *Int J Urol* 2017;24:352–60.
- Partin AW, Van Neste L, Klein EA, Marks LS, Gee JR, Troyer DA, et al. Clinical validation of an epigenetic assay to predict negative histopathological results in repeat prostate biopsies. *J Urol* 2014;192:1081–7.
- Blute Jr ML, Damaschke NA, Jarrard DF. The epigenetics of PCA diagnosis and prognosis: update on clinical applications. *Curr Opin Urol* 2015;25:83.
- Krop I, Player A, Tablante A, Taylor-Parker M, Lahti-Domenici J, Fukuoka J, et al. Frequent HIN-1 Promoter Methylation and Lack of Expression in Multiple Human Tumor Types11National Cancer Institute SPORE in Breast Cancer at Dana-Farber/Harvard Cancer Center grant CA89393, NIH RO1 grant CA94074-01A1, NIH National Research Service Award CA94787-01, Dunkin'Donuts "Rising Stars" Award, and V Foundation. *Mol Cancer Res* 2004;2:489–94.
- De Marzo AM, Platz EA, Sutcliffe S, Xu J, Grönberg H, Drake CG, et al. Inflammation in prostate carcinogenesis. *Nat Rev Cancer* 2007;7:256–69.
- Kwon O-J, Zhang L, Ittmann MM, Xin L. Prostatic inflammation enhances basal-to-luminal differentiation and accelerates initiation of prostate cancer with a basal cell origin. *Proc Natl Acad Sci USA* 2014;111:E592–600.
- Sutcliffe S. Sexually transmitted infections and risk of prostate cancer: review of historical and emerging hypotheses. *Future Oncol* 2010;6:1289–311.
- Ge X, Wang X, Shen P. Herpes simplex virus type 2 or human herpesvirus 8 infection and prostate cancer risk: A meta-analysis. *Biomed Rep* 2013;1:433–9.
- Golais F, Mrázová V. Human alpha and beta herpesviruses and cancer: passengers or foes? *Folia Microbiol* 2020;65:439–49.