

Allostatic Load, Income, and Race Among Black and White Men in the United States

American Journal of Men's Health
March -April 1–11
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15579883221092290
journals.sagepub.com/home/jmh


Carlos D. Tavares¹ , Caryn N. Bell², Hossein Zare^{3,4} ,
Darrell Hudson⁵ , and Roland J. Thorpe Jr.⁶ 

Abstract

Research indicates that income is significantly associated with allostatic load (AL) and that this association may differ between White and Black Americans. Most existing income–AL link work focuses on women and less is known about this association among men. Using data from the National Health and Nutrition Examination Survey (NHANES), we examined whether race moderates the association between income and AL among Black and White men in the United States ($n = 5,685$). We find that, regardless of income levels, Black men have significantly higher prevalence of being in the high-AL group compared with high-income White men. Our findings suggest that Black men do not receive the same health benefits for increased income relative to their White counterparts.

Keywords

black men, special populations, allostatic load, income, unequal returns

Received August 11, 2021; revised February 4, 2022; accepted March 17, 2022

Introduction

Despite significant social advantages, men in the United States have shorter life spans and greater levels of life-threatening chronic diseases relative to women (Griffith et al., 2019; Rieker & Bird, 2005). Evidence from previous research demonstrates that Black men's health profiles are worse compared with their White counterparts (Bell & Thorpe, 2019; Wilson-Frederick et al., 2014). A number of explanations have been explored for men's health disparities including masculinity, income, and stress (Bell et al., 2018; Bell & Thorpe, 2019; Gilbert et al., 2016; Griffith et al., 2016; Nuru-Jeter et al., 2018).

Stress exposure is an important mechanism through which social and environmental experiences become embodied (Upchurch et al., 2015). Allostatic load (AL) is one of the primary measures scholars use to quantify the cumulative burden of exposure to stress. AL is theorized to capture repeated adaptations to stressors that can lead to dysregulation in multiple physiological human systems (Beckie, 2012; Geronimus et al., 2006). This dysregulation has been identified to be detrimental to health and well-being, with a number of studies demonstrating

that AL is associated with higher levels of mortality and poor physical and mental health outcomes (Beckie, 2012; Borrell et al., 2020; Thorpe et al., 2020).

Previous research suggests that lower income levels significantly predict AL levels (Hickson et al., 2012; Upchurch et al., 2015). Consistent with fundamental

¹Department of Anthropology and Sociology, Lafayette College, Easton, PA, USA

²Department of Social, Behavioral, and Population Sciences, Tulane University, New Orleans, LA, USA

³Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

⁴Global Health Services and Administration, University of Maryland Global Campus, Baltimore, MD, USA

⁵Brown School, Washington University in St. Louis, St. Louis, MO, USA

⁶Department of Health, Behavior, and Society, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Corresponding Author:

Roland J. Thorpe Jr., Professor, Department of Health, Behavior, and Society, Johns Hopkins Bloomberg School of Public Health, 624 N. Broadway, Ste 708, Baltimore, MD 21205, USA.
Email: rthorpe@jhu.edu



cause theory, which posits that resources, especially related to income, significantly shape health disparities (Phelan et al., 2010), there is a well-established link between income and AL, such that most previous studies indicate that higher income is generally associated with lower AL scores (Merkin et al., 2020; Rainisch & Upchurch, 2013). These findings further align with the conservation of resources (COR) theory, which proposes that socioeconomic resources, such as income, are significantly linked with the availability of resources and the ability to gain and protect resources, which in turn are associated with well-being (Hobfoll, 1989, 2011). There is evidence documenting significant racial and ethnic disparities in AL (Gaskin et al., 2020; Geronimus et al., 2006; Upchurch et al., 2015). These disparities are not fully explained by income differences across groups (Gaskin et al., 2020; Geronimus et al., 2006). Research suggests that, relative to White Americans, biological dysregulation among Black Americans is not as significantly patterned by income and other SES measures (Hickson et al., 2012). Most of the existing research focuses specifically on women contrasting the income–AL link between different racial groups. This research suggests that income differences may explain most of the AL differences between White and Hispanic women, but does not account for as much variation between White and Black American women (Gaskin et al., 2020). The little work that includes men focuses on comparing the income–AL association among Black men and women (Hickson et al., 2012). These studies provide some preliminary evidence that the income–AL relationship may work differently among Black American men, but it is not well understood whether race moderates the income–AL link among men in the United States.

Minority Stress Theory

Minority stress theory is a relevant framework for exploring the role race may play in the income–AL association. It posits that members of oppressed social groups experience additional stressors as a result of their social position with the social structure (Meyer, 2003; Williams et al., 2019). This theory further suggests that the nature of minority stress is chronic and tied to social processes. Important here, Black Americans experience added stressors (Sternthall et al., 2011) and that these stressors are chronic (Lewis et al., 2015). This racialized stress has been linked to poor health outcomes (Cobb et al., 2019; Colen et al., 2018). Other research indicates that resources, such as income, which are generally health protective, may not have the same effect for Black Americans (Turner et al., 2017) in large part because of racism-related stress (Colen et al., 2018). In the context of our study, it is important to understand whether race

moderates the relationship between income and AL as unique racialized stressors among Black men may contribute to differential outcomes relative to White men.

Although there is strong evidence that both income and race have important implications for AL, it is less clear whether the income–AL link varies by race among American men. Given the increasing evidence of relatively poor health among Black men amid significant exposure to minority stressors, it is important to understand if these patterns extend to the association between income and AL among Black and White men. In this study, we examined whether race moderated the association between income and AL among U.S. men. Relative to White men, we expected that higher income will not be linked with lower risk of high AL for Black men. Understanding race differences in the association between income and AL among men will contribute important knowledge to the study of men's health disparities in AL.

Method

Data for this study were drawn from the National Health and Nutrition Examination Survey (NHANES), which focuses on the health, nutritional, and functional status of the noninstitutionalized civilian population in the United States. Conducted as a continuous annual survey since 1999 (Chen et al., 2018), NHANES has released public use data in 2-year increments (e.g., NHANES 1999–2000, NHANES 2001–2002, and NHANES 2003–2004). Data were collected from respondents in two steps. First, information regarding the participant's health history, health behaviors, and risk factors were obtained during the home interview. Second, participants were invited to participate in a medical examination where they received a detailed physical examination at a mobile examination center (Chen et al., 2018). Each of these surveys used a version of a stratified, multistage probability sampling design (Fakhouri et al., 2020). Additional details regarding NHANES data and methodology is described elsewhere (NHANES, 2018). The National Center for Health Statistics ethics review board approved protocols and written informed consent was obtained from all NHANES participants.

In this analysis, we combined nine waves of the NHANES data between 1999 and 2016 ($n = 92,062$). We limited our analytical sample to all men aged ≥ 20 years who had valid data on biomarkers used to create AL score and identified themselves as non-Hispanic White (NHW) and non-Hispanic Black (NHB). Important covariates including education and marital status are only asked of respondents aged ≥ 20 years. The NHANES family income variable ranges from 1 ($\$0$ – $\$4,999$) to 11 ($\$75,000$ and above). We dichotomize this variable into less than




Figure 1. Race \times Income Marginal Effects.

Source. National Health and Nutrition Examination Study, 1999–2016.

\$75,000 and equal or more than \$75,000. This yielded an analytic sample of 5,685 men, which included 1,633 men who reported family income greater than or equal to \$75,000 and 4,529 men who reported family income less than \$75,000.

Our dependent variable, AL, was derived from eight biomarkers, including systolic blood pressure (mm Hg), diastolic blood pressure (mm Hg), pulse rate (beats/min), body mass index (BMI; kg/m²), glycohemoglobin (%), direct high-density lipoprotein (HDL)-cholesterol (mg/dL), total cholesterol (mg/dL), and serum Albumin (g/dL). Following the approach developed by Chyu and Upchurch (2011), we considered values above the 75th percentile as high risk for all the biomarkers, except for HDL and serum albumin, for which values below the 25th percentile was defined as high risk (Chyu & Upchurch, 2011; Geronimus et al., 2006). Each biomarker that was considered to be high risk was assigned a 1 and those considered to be low risk were assigned a 0. After summing across the biomarker indicators, we obtained a score ranging from 0 to 8 for each respondent. Finally, based on prior literature (Geronimus et al., 2006; Seeman et al., 1997), we created a dummy variable to identify men with a high AL versus low AL (1, if AL \geq 4; 0, if AL < 4).

Covariates included demographic and health-related characteristics. Demographic variables included age (years), marital status (1 = *married*, 0 = *not married*), and education categories (less than high school graduate, high school graduate or General Educational Development equivalent recipient, or more than high school education). Health-related characteristics included health insurance (1 = *yes*, 0 = *no*), self-reported health (fair/poor vs. excellent/very good/good), and household size (total number of people in the household).

Analytic Approach

The mean and proportional differences between NHB and NHW men with income \geq \$75,000 and income <\$75,000 for demographic, health-related characteristics and biomarkers were evaluated using Student's *t* tests and chi-square test. In our analyses, the prevalence of AL \geq 4 was greater than 10%; therefore, we used a weighted modified Poisson regression analysis (McNutt et al., 2003; Thorpe et al., 2017; Zou, 2004) that produced prevalence ratios (PRs) and corresponding 95% confidence intervals (CIs; McNutt et al., 2003; Thorpe et al., 2017; Zou, 2004). We ran three models for our analyses. Model 1 was an unadjusted model examining the relationship between income–race and AL score. We created a categorical variable by interacting income more than \$75,000 and race/ethnicity (NHW and NHB) and included that variable as the main independent variable. Model 2 was an adjusted Model 1, including all control variables except self-reported health, and Model 3 is the full model including all control variables. Because the interaction between income and race/ethnicity was significant ($p = .0005$), we followed the interaction approach used in Gaskin et al. (2014) and created four dummy variables using race (NHB and NHW) and income (if >\$75K) as interaction between race/ethnicity and income. The interaction marginal effects are plotted in Figure 1. All descriptive statistics and Poisson regression estimates were weighted using the NHANES individual-level sampling weights for 1999 to 2016 (nine waves of data). The p values < .05 were considered statistically significant; *t* tests were two-sided. All statistical procedures were performed using STATA statistical software, Version 15.

Results

The distribution of sociodemographic characteristics and AL among NHB and NHW men for the total sample and by income is presented in Table 1. Of the 5,685 men, 12.9% were NHB and the average age was 47.3 ± 12.9 years. Overall, the majority of men reported their education as some college and above, family income between \$35,000 and \$74,999, being married, and having health insurance. A small portion of the men reported their health as fair or poor. Men who reported family income greater than or equal to \$75,000 were more likely to be younger, have some college education or above, be married, and have health insurance than men who reported family income less than \$75,000. Men who reported income greater than or equal to \$75,000 were less likely to rate their health as fair or poor relative to men who reported income less than \$75,000.

Table 1. Distribution of Sociodemographic Characteristics and Mean Allostatic Load Score Among Adult Men, NHANES 1999–2016.

Variables	All (N = 5,685)	Family income ≥\$75K (n = 1,528)	Family income <\$75K (n = 4,665)	p value ^a
Age—years (M, SD)	47.3 (12.9)	46.5 (9.5)	47.7 (14.7)	.0455
Non-Hispanic Black (%)	1,778 (12.1)	347 (6.5)	1,431 (15.2)	.000
Education (%)				
Less than high school	1,337 (14.2)	117 (4.9)	1,220 (19.5)	.000
High school graduate/GED	1,676 (25.4)	278 (16.8)	1,398 (30.4)	
Some college and above	3,319 (60.2)	1,279 (78.1)	2,040 (49.9)	
Family income (%)				
\$0–\$34,999	2,416 (25.7)	—	2,416 (40.5)	
\$35,000–\$74,999	2,249 (37.7)	—	2,249 (59.4)	
≥\$75,000	1,675 (36.5)	1,675 (100)	—	
Missing	106 (1.1)			
Married (%)	4,154 (68.6)	1,327 (78.9)	2,827 (62.7)	.000
Self-rated fair/poor (%)	277 (3.1)	25 (1.1)	252 (4.3)	.000
Covered by health insurance (%)	5,196 (84.4)	1,572 (94.0)	3,624 (79.0)	.000
Household size	2.9 (1.1)	3.1 (1.0)	2.7 (1.2)	.000

Note. GED = General Educational Development.

^ap value reports adjusted Wald test results for age and chi-square for all other variables.

Table 2. Distribution of Individual Biomarkers and AL Score Among Income in Adult Men, NHANES 1999–2016.

Variables	All (N = 5,685), M/SD	Family income ≥\$75K (n = 1,633), M/SD	Family income <\$75K (n = 4,529), M/SD	p value ^a	Quartiles		
					25%	50%	75%
Cardiovascular markers (%)							
Systolic blood pressure (mm Hg)	2,251 (29.6)	476 (24.0)	1,775 (32.9)	<.001	108	116.7	128
Diastolic blood pressure (mm Hg)	2,215 (36.8)	647 (39.1)	1,568 (35.4)	.030	61.3	68.7	76
Pulse rate (beats /min)	1,237 (19.9)	288 (18.4)	949 (20.8)	.089	64	72	80
Metabolic markers (%)							
Body mass index (kg/m ²)	2,429 (40.0)	687 (40.7)	1,742 (39.8)	.636	22.7	26.6	31.2
Glycohemoglobin (%)	2,167 (26.8)	459 (22.1)	1,708 (29.5)	<.001	5.1	5.3	5.6
Direct HDL-cholesterol (mg/dL)	1,740 (29.5)	496 (25.8)	1,344 (31.6)	<.001	42	51	62
Total cholesterol (mg/dL)	2,012 (33.5)	564 (34.8)	1,448 (32.8)	.275	160	187	215
Inflammatory markers (%)							
Serum albumin (g/dL)	2,470 (32.4)	560 (28.4)	1,910 (34.6)	<.001	4.1	4.3	4.5
High-AL ^b (%)	1,874 (26.8)	457 (24.8)	1,417 (27.9)	.030	—	—	—

Note. AL = allostatic load; HDL = high-density lipoprotein.

^ap value reports adjusted Wald test results. ^bAL score computed as sum of all markers, existence of a condition considered as 1 and otherwise as 0. Values above the 75th percentile were defined as high risk for all the biomarkers, with the exception of HDL and serum albumin for which values below the 25th percentile were defined as high risk. High AL was based on those men who had 4 or more biomarkers, which was considered to be high risk.

The distribution of individual biomarkers and AL by family income among NHB and NHW men is presented in Table 2. Men who reported income greater than or equal to \$75,000 were less likely to be above the at-risk cut points for systolic blood pressure, glycohemoglobin, HDL serum albumin, and having four or more high-risk biomarkers compared with men who reported family income less than \$75,000. Men who reported income greater than or equal to \$75,000 were more likely to be above the at-risk cut points for diastolic blood pressure than men who reported income less than \$75,000. No differences were observed between men who reported

income greater than or equal to \$75,000 and those men who reported income less than \$75,000 as it relates to pulse rate, BMI, and total cholesterol.

The association between AL, race, and family income among NHB and NHW men is presented in Table 3. After adjusting for age, education, marital status, insurance status, fair or poor health, and total number of people in the household, NHB men who reported income greater than or equal to \$75,000 (PR = 1.58, 95% CI = [1.28, 1.94]) and NHB men who reported income less than \$75,000 had a higher prevalence of being in the high-AL group (PR = 1.25, 95% CI = [1.09, 1.44]) compared with

Table 3. AL Score^a, Adult White and Black NH Men in National Health and Nutrition Examination Survey 1999–2016 (N = 5,685).

Variables	Model 1		Model 2		Model 3	
	PR	CI-95%	PR	CI-95%	PR	CI-95%
Family income # racial/ethnic (Ref: NH White with income \geq \$75K)						
NH White, <\$75K	1.14	[1.01, 1.30]	1.08	[0.95, 1.23]	1.08	[0.95, 1.22]
NH Black, \geq \$75K	1.56	[1.27, 1.92]	1.58	[1.28, 1.94]	1.58	[1.28, 1.94]
NH Black, <\$75K	1.25	[1.09, 1.44]	1.26	[1.09, 1.45]	1.25	[1.08, 1.45]
Age (years)			1.01	[1.01, 1.01]	1.01	[1.01, 1.02]
Education (Ref: Less than high school)						
High school graduate/GED			1.09	[0.95, 1.25]	1.10	[0.96, 1.27]
Some college and above			0.86	[0.75, 0.97]	0.87	[0.76, 0.99]
Marital status (Ref: Unmarried)						
Married or living with partner			0.97	[0.85, 1.09]	0.97	[0.86, 1.10]
Covered by health insurance						
			1.16	[0.98, 1.37]	1.15	[0.98, 1.37]
Self-rated health (Ref: Excellent/very good/good)						
Fair/poor					1.24	[1.00, 1.53]
Household size	0.99	[0.94, 1.02]	1.04	[0.99, 1.09]	1.04	[1.00, 1.09]

Note. AL = allostatic load; PR = prevalence ratio; CI = confidence interval; GED = General Educational Development; HDL = high-density lipoprotein.

^aFor this analysis, we used the AL score computed as sum of all 8 biomarkers, existence of a condition considered as 1 and otherwise as 0. Values above the 75th percentile were defined as high risk for all the biomarkers, with the exception of HDL and serum albumin for which values below the 25th percentile were defined as high risk. We then created a dummy variable, that is, 1 if AL score \geq 4 or 0 if otherwise, and used that variable for the Poisson model. High AL was based on those men who had 4 or more biomarkers, which was considered to be high risk.

NHW men who reported income greater than or equal to \$75,000.

As a check on the robustness of our findings, we estimated a regression model stratified by income (see Appendix A) and ran regression models using a lower income cut point (\$35,000; see Appendix B). In both sensitivity analyses, we identify similar results to those discussed above.

Discussion

In this article, we sought to determine whether race moderates the association between income and AL among NHB and NHW men in the United States. Using minority stress theory, we expected that higher income would not be associated with lower prevalence of being in the high-AL group for NHB men compared with NHW men. The findings provide strong support for our hypothesis. Relative to White men with family income equal to or greater than \$75,000, NHB men were more likely to be in the high-AL group irrespective of income. Both NHB men with family income below \$75,000 and equal to or greater than \$75,000 were more likely to be in the high-AL group relative to reference group. Relative to NHW men, higher income is not associated with lower risk of being in the high-AL group among NHB men. Our work here indicates that race plays a significant role in the association between income and AL among U.S. men.

Most of the existing research on race, income, and AL to date has been among women. Our findings are consistent with those drawn from women in the United States, which indicate that income does not explain AL variation as well for Black Americans as it does for White Americans (Gaskin et al., 2020; Hickson et al., 2012). Previous research comparing Black women and men suggests that biological dysregulation in the latter group is less tied to socioeconomic status (Hickson et al., 2012). We contribute to the literature by focusing specifically on this association among Black and White men. Our research indicates that, relative to White men, higher income in Black men does not seem to offer protection against being in the high-AL group.

Our main finding aligns with other research on race, socioeconomic status, and health outcomes, particularly that Black men experience diminished or unequal returns of various measures of SES to health (Assari & Bazargan, 2019; Farmer & Ferraro, 2005; Hudson et al., 2012, 2019). In other words, relative to White men, Black men gain less health benefit from socioeconomic resources such as income and education. Furthermore, some studies report that higher income may be associated with higher obesity rates among Black men (Bell & Thorpe, 2020). Our findings suggest that stressors associated with upward mobility that contribute to diminished returns may be a factor in the association between income and AL, given that NHB men have a higher likelihood of having high AL regardless of their income.

According to minority stress theory, members of oppressed groups experience additional stressors, relative to dominant group members, which contribute to poor health. These stressors are chronic and are products of the racial hierarchy. Specifically, structural racism, interpersonal discrimination, and psychosocial factors have been implicated as contributing factors to Black/White men's health disparities. Structural racism refers to the totality of racialized social relations and practices, such as redlining, which have led to deeply entrenched racial residential segregation throughout the United States, as key factors in the creation and reification of racial inequality (Bailey et al., 2017; Bonilla-Silva, 1997; Gee & Ford, 2011). For example, Thorpe and colleagues (2015) reported that social and environmental conditions play a key role in Black–White differences in chronic conditions among men who lived in urban, low-income integrated census tracts in the Exploring Health Disparities in Integrated Communities–Southwest Baltimore (EHDIC-SWB) Study, highlighting the role that residential segregation may play in Black's men health. Similarly, Brown et al. (2015) identified that neighborhood conditions explain much of the Black/White differences in functional limitations among men. Black adults with higher incomes are more likely to live in poorer, more racially segregated neighborhoods than high-income White adults (Darden et al., 2018; Reardon et al., 2015). It is likely that the mechanisms through which racial segregation negatively impacts health among low-income Black adults also affects high-income Black adults, resulting in weak associations between income and AL among Black men.

Structural racism contributes to racial disparities in criminal justice contact. Compared with White men, Black men are at a significantly higher risk for arrests, convictions, and incarceration (Hudson et al., 2019; Wildeman & Wang, 2017). Furthermore, serious criminal justice contact is associated with higher reports of everyday discrimination among Black men (Taylor et al., 2016), which may contribute to poor health. Given the negative impact that criminal justice contact has on health, it is an important consideration here (Archibald et al., 2018; Boen, 2020; Wildeman & Wang, 2017). Qualitative research with Black men suggests that they are cognizant of the role that residential racial segregation, unequal educational opportunities, and racial disparities in criminal justice contact play in their life chances (Hudson et al., 2016).

There is significant evidence that interpersonal discrimination contributes to poor health for Black men (Wheaton et al., 2018; Williams et al., 2019) and reports of racial discrimination increase with higher income and education for Black men (Hudson et al., 2012; Jackson & Williams, 2006). Black middle-class men tend to find themselves in White spaces more often and deal with stressors related to identity management, hypervisibility,

and differential treatment (Hudson et al., 2020), which may contribute to the lower health returns they receive for their higher socioeconomic status (i.e., diminished returns) relative to their White counterparts. Indeed, some scholars conclude that working in predominantly White workspaces may explain why high-SES Black men report more interpersonal discrimination (Assari & Moghani Lankarani, 2018).

This study has significant strengths. First, we use a large, nationally representative sample of Black and White men to build on the income–allostatic link research. Second, we use a higher income cut point (\$75,000) than most previous studies, which allows us to examine whether the association between income and allostatic works differently at higher income brackets. Third, the inclusion of biomarkers was crucial for calculating our measure of biological dysregulation.

Future Research

Our study is not without limitations, which include the use of cross-sectional data. Hence, we are not able to examine causal effects. Future researchers should examine whether changes or stability in income over time affect AL among men in different racialized groups in the United States. It is possible that upward or downward income trajectories may result in different patterns from the ones we identified here. Key historical events, such as the Great Recession, might play an important role, so future researchers might explore how they impact the association explored here because Black and White men were differentially impacted by them (Addo & Darity, 2021; Thomas et al., 2018).

The associations observed here are also not generalizable to Black women or other racial groups. Some research indicates that Black women in higher socioeconomic status groups have similar experiences navigating predominantly White spaces and are subjected to various types of discrimination (Sacks, 2019). There may be similarities in the experiences of higher income Black men and Black women and further research is needed in this area.

One area of focus for future researchers may include differences in exposure to racial violence between Black and White men and what contribution that makes to the diminished health returns among higher income Black men. There are significant racial disparities in hostile police encounters and police killings (Edwards et al., 2019; Geller et al., 2014). Even when they are not victims in police violence, their mental health may be affected by police killings of Black Americans in their respective states (Bor et al., 2018). It is essential to examine how this may contribute to poor health among higher income Black men as increased socioeconomic resources do not necessarily protect against these types of encounters. Our

focus here was on the specific income–AL association and the moderating role that race plays. Future researchers should explore socioeconomic measures that include income, education, occupation, and particularly wealth (Marsh et al., 2007). Given that wealth has been identified as the greatest area of inequality between Black and White Americans (Hamilton et al., 2015), it is important to understand how wealth may be associated with AL among U.S. men.

Implications

The findings of this study suggest that, relative to White men, higher income among Black men is not associated with lower risk of poor health. Public health interventions that focus only on income may not significantly reduce health disparities between Black and White men in the United States. Structural and interpersonal racism, and psychosocial factors, need to be addressed if our goal is to address racial disparities among Black and White men.

Conclusion

In this article, we explored whether race moderated the association between income and AL among Black and White men. We found that, compared with White men with reported family income equal or greater than \$75,000, Black men had higher risk of high AL regardless of their income. This finding is consistent with research on the diminished health returns of higher socioeconomic

status for Black Americans relative to White Americans. The findings here underscore the need to address racism and racism-related stressors at the public policy level. Interventions aimed at reducing health disparities should not focus only on income among Black American men. This study has implications for social workers and clinicians who need to be cognizant of the unique stressors and relative poor health of Black American men.

Future research would benefit from a longitudinal focus on dynamic changes or stability in income and how those are related to biological dysregulation in Black and White men. Those interested in this area would also benefit from exploring a multidimensional socioeconomic status definition, including income, wealth, and other relevant indicators. Future research should investigate what are the exact mechanisms that contribute to this moderation. Although there is significant evidence that racialized minority stress contributes to these findings, we were not able to test this directly. More studies are needed to untangle the complex ways that psychosocial factors may contribute to Black men's health, including how stressors get under the skin to dysregulate systems over time.

Considering the growing number of researchers investigating diminished health returns among Black men, a focus on higher income Black men would provide a greater understanding of the unique stressors and experiences that might lead to lower health returns and inspire interventions targeted toward reducing racial health disparities among U.S. men.

Appendix A. AL Score^a, Adult White and Black NH Men in National Health and Nutrition Examination Survey 1999–2016, Stratified Model by Income.

Variables	Family income ≥ \$75K		Family income < \$75K	
	PR	CI-95	PR	CI-95
Racial/ethnicity (Ref: NH White)				
NH Black	1.57	[1.27, 1.95]	1.16	[1.03, 1.31]
Age (years)	1.02	[1.01, 1.03]	1.01	[1.01, 1.02]
Education (Ref: Less than high school)				
High school graduate/GED	1.11	[0.76, 1.61]	1.06	[0.91, 1.23]
Some college and above	0.67	[0.47, 0.95]	0.94	[0.82, 1.08]
Marital status (Ref: Unmarried)				
Married or living with partner	0.97	[0.73, 1.27]	0.97	[0.85, 1.12]
Covered by health insurance	1.53	[0.91, 2.57]	1.13	[0.95, 1.36]
Self-rated health (Ref: Excellent/very good/good)				
Fair/poor	1.48	[0.93, 2.35]	1.21	[0.95, 1.54]
Household size	1.06	[0.99, 1.14]	1.03	[0.98, 1.10]

Note. AL = allostatic load; NH = non-Hispanic; PR = prevalence ratio; CI = confidence interval; GED = General Educational Development; HDL = high-density lipoprotein.

^aFor this analysis, we used the AL score computed as sum of all eight biomarkers, existence of a condition considered as 1 and otherwise as 0. Values above the 75th percentile were defined as high risk for all the biomarkers, with the exception of HDL and serum albumin, for which values below the 25th percentile were defined as high risk. We then created a dummy variable, that is, 1 if AL score ≥ 4 or 0 if otherwise, and used that variable for the Poisson model. High AL was based on those men who had 4 or more biomarkers, which was considered to be high risk.

Appendix B. Poisson Estimates for AL Score^a, Adult White and Black NH Men in National Health and Nutrition Examination Survey 1999–2016.

Variables	Model 1		Model 2		Model 3	
	PR	CI-95	PR	CI-95	PR	CI-95
Family income (Ref: WNH ≥\$35K)						
NH White, <\$35K	1.26	[1.13, 1.41]	1.22	[1.09, 1.36]	1.121	[1.08, 1.35]
NH Black, ≥\$35K	1.36	[1.19, 1.56]	1.42	[1.24, 1.62]	1.42	[1.24, 1.62]
NH Black, <\$35K	1.17	[1.03, 1.33]	1.22	[1.07, 1.40]	1.21	[1.06, 1.39]
Age (years)			1.01	[1.01, 1.02]	1.01	[1.01, 1.02]
Education (Ref: Less than high school)						
High school graduate/GED			1.12	[0.96, 1.27]	1.12	[0.98, 1.30]
Some college and above			0.89	[0.78, 1.01]	0.90	[0.79, 1.02]
Marital status (Ref: Unmarried)						
Married or living with partner			0.99	[0.87, 1.12]	0.99	[0.87, 1.12]
Covered by health insurance			1.18	[1.00, 1.40]	1.18	[1.00, 1.48]
Self-rated health (Ref: Excellent/very good/good)						
Fair/poor					1.19	[0.96, 1.48]
Household size	0.99	[0.95, 1.03]	1.05	[1.00, 1.09]	1.05	[1.00, 1.09]

Note. AL = allostatic load; NH = non-Hispanic; WNH = White non-Hispanic; PR = prevalence ratio; CI = confidence interval; GED = General Educational Development; HDL = high-density lipoprotein.

^aFor this analysis we used the AL score computed as sum of all 8 biomarkers, existence of a condition considered as 1 and otherwise 0. Values above the 75th percentile were defined as high risk for all the biomarkers, with the exception of HDL and serum albumin, for which values below the 25th percentile was defined as high risk. We then created a dummy variable, that is, 1 if AL score ≥ 4 or 0 if otherwise, and used that variable for the Poisson model.

Interaction test: 0.0005.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by the National Institute on Aging (K02AG059140—R.J.T.) and the National Institute on Minority Health and Health Disparities (U54MD000214—R.J.T.). The funders had no role in study design, data collection, analysis, decision to publish, or manuscript preparation.

ORCID iDs

Carlos D. Tavares  <https://orcid.org/0000-0001-9888-645X>

Hossein Zare  <https://orcid.org/0000-0002-5832-0854>

Darrell Hudson  <https://orcid.org/0000-0003-1385-5887>

Roland J. Thorpe Jr.  <https://orcid.org/0000-0002-4448-4997>

References

- Addo, F. R., & Darity, W. A., Jr. (2021). Disparate recoveries: Wealth, race, and the working class after the Great Recession. *The ANNALS of the American Academy of Political and Social Science*, 695(1), 173–192.
- Archibald, P. C., Parker, L., & Thorpe, R. (2018). Criminal justice contact, stressors, and obesity-related health problems among Black adults in the USA. *Journal of Racial and Ethnic Health Disparities*, 5(2), 387–397. <https://doi.org/10.1007/s40615-017-0382-9>
- Assari, S., & Bazargan, M. (2019). Minorities' diminished returns of educational attainment on hospitalization risk: National Health Interview Survey (NHIS). *Hospital Practices and Research*, 4(3), 86–91. <https://doi.org/10.15171/hpr.2019.17>
- Assari, S., & Moghani Lankarani, M. (2018). Workplace racial composition explains high perceived discrimination of high socioeconomic status African American men. *Brain Sciences*, 8(8), Article 139. <https://doi.org/10.3390/brain-sci8080139>
- Bailey, Z. D., Krieger, N., Agénor, M., Graves, J., Linos, N., & Bassett, M. T. (2017). Structural racism and health inequities in the USA: Evidence and interventions. *Lancet*, 389(10077), 1453–1463. [https://doi.org/10.1016/s0140-6736\(17\)30569-x](https://doi.org/10.1016/s0140-6736(17)30569-x)
- Beckie, T. M. (2012). A systematic review of allostatic load, health, and health disparities. *Biological Research for Nursing*, 14(4), 311–346. <https://doi.org/10.1177/1099800412455688>
- Bell, C. N., & Thorpe, R. J. Jr. (2019). Income and marital status interact on obesity among Black and White men. *American Journal of Men's Health*, 13(1), 1557988319829952. <https://doi.org/10.1177/1557988319829952>
- Bell, C. N., & Thorpe, R. J. Jr. (2020). Association between income and obesity in Black men: The role of work-life

- interference. *Ethnicity and Disease*, 30(4), 629–636. <https://doi.org/10.18865/ed.30.4.629>
- Bell, C. N., Thorpe, R. J. Jr., Bowie, J. V., & LaVeist, T. A. (2018). Race disparities in cardiovascular disease risk factors within socioeconomic status strata. *Annals of Epidemiology*, 28(3), 147–152. <https://doi.org/10.1016/j.annepidem.2017.12.007>
- Boen, C. E. (2020). Criminal justice contacts and psychophysiological functioning in early adulthood: Health inequality in the carceral state. *Journal of Health and Social Behavior*, 61(3), 290–306. <https://doi.org/10.1177/0022146520936208>
- Bonilla-Silva, E. (1997). Rethinking racism: Toward a structural interpretation. *American Sociological Review*, 62(3), 465–480. <https://doi.org/10.2307/2657316>
- Bor, J., Venkataramani, A. S., Williams, D. R., & Tsai, A. C. (2018). Police killings and their spillover effects on the mental health of Black Americans: A population-based, quasi-experimental study. *The Lancet*, 392(10144), 302–310. [https://doi.org/10.1016/S0140-6736\(18\)31130-9](https://doi.org/10.1016/S0140-6736(18)31130-9)
- Borrell, L. N., Rodríguez-Álvarez, E., & Dallo, F. J. (2020). Racial/ethnic inequities in the associations of allostatic load with all-cause and cardiovascular-specific mortality risk in U.S. adults. *PLOS ONE*, 15(2), Article e0228336. <https://doi.org/10.1371/journal.pone.0228336>
- Brown, T. H., Hargrove, T. W., & Griffith, D. M. (2015). Racial/ethnic disparities in men's health: Examining psychosocial mechanisms. *Family & Community Health*, 38(4), 307–318. <https://dx.doi.org/10.1097%2FFCH.0000000000000080>
- Chen, T. C., Parker, J. D., Clark, J., Shin, H. C., Rammon, J. R., & Burt, V. L. (2018). National Health and Nutrition Examination Survey: Estimation procedures, 2011–2014. *Vital Health Statistics*, 2(177), 1–26.
- Chyu, L., & Upchurch, D. M. (2011). Racial and ethnic patterns of allostatic load among adult women in the United States: Findings from the National Health and Nutrition Examination Survey 1999–2004. *Journal of Women's Health*, 20(4), 575–583. <https://doi.org/10.1089/jwh.2010.2170>
- Cobb, R. J., Thorpe, R. J. Jr. & Norris, K. C. (2019). Everyday discrimination and kidney function among older adults: Evidence from the health and retirement study. *The Journals of Gerontology: Series A*, 75(3), 517–521. <https://doi.org/10.1093/gerona/glz294>
- Colen, C. G., Ramey, D. M., Cooksey, E. C., & Williams, D. R. (2018). Racial disparities in health among nonpoor African Americans and Hispanics: The role of acute and chronic discrimination. *Social Science & Medicine*, 199, 167–180. <https://doi.org/10.1016/j.socscimed.2017.04.051>
- Darden, J., Malega, R., & Stallings, R. (2018). Social and economic consequences of Black residential segregation by neighbourhood socioeconomic characteristics: The case of Metropolitan Detroit. *Urban Studies*, 56(1), 115–130. <https://doi.org/10.1177/0042098018779493>
- Edwards, F., Lee, H., & Esposito, M. (2019). Risk of being killed by police use of force in the United States by age, race–ethnicity, and sex. *Proceedings of the National Academy of Sciences*, 116(34), 16793–16798. <https://doi.org/10.1073/pnas.1821204116>
- Fakhouri, T. H. I., Martin, C. B., Chen, T. C., Akinbami, L. J., Ogden, C. L., Paulose-Ram, R., Riddles, M. K., Van de Kerckhove, W., Roth, S. B., Clark, J., Mohadjer, L. K., & Fay, R. E. (2020). An investigation of nonresponse bias and survey location variability in the 2017–2018 National Health and Nutrition Examination Survey. *Vital Health Statistics*, 2(185), 1–36.
- Farmer, M. M., & Ferraro, K. F. (2005). Are racial disparities in health conditional on socioeconomic status? *Social Science & Medicine*, 60(1), 191–204. <https://doi.org/10.1016/j.socscimed.2004.04.026>
- Gaskin, D. J., Thorpe, R. J. Jr. McGinty, E. E., Bower, K., Rohde, C., Young, J. H., LaVeist, T. A., & Dubay, L. (2014). Disparities in diabetes: The nexus of race, poverty, and place. *American Journal of Public Health*, 104(11), 2147–2155. <https://doi.org/10.2105/AJPH.2013.301420>
- Gaskin, D. J., Zare, H., Jackson, J. W., Ibe, C., & Slocum, J. (2020). Decomposing race and ethnic differences in CVD risk factors for mid-life women. *Journal of Racial and Ethnic Health Disparities*, 8, 174–185. <https://doi.org/10.1007/s40615-020-00769-9>
- Gee, G. C., & Ford, C. L. (2011). Structural racism and health inequities: Old issues, new directions. *Du Bois Review: Social Science Research on Race*, 8(1), 115–132. <https://doi.org/10.1017/s1742058x11000130>
- Geller, A., Fagan, J., Tyler, T., & Link, B. G. (2014). Aggressive policing and the mental health of young urban men. *American Journal of Public Health*, 104(12), 2321–2327. <https://doi.org/10.2105/AJPH.2014.302046>
- Geronimus, A. T., Hicken, M., Keene, D., & Bound, J. (2006). “Weathering” and age patterns of allostatic load scores among Blacks and Whites in the United States. *American Journal of Public Health*, 96(5), 826–833. <https://doi.org/10.2105/AJPH.2004.060749>
- Gilbert, K. L., Ray, R., Siddiqi, A., Shetty, S., Baker, E. A., Elder, K., & Griffith, D. M. (2016). Visible and invisible trends in Black men's health: Pitfalls and promises for addressing racial, ethnic, and gender inequities in health. *Annual Review of Public Health*, 37(1), 295–311. <https://doi.org/10.1146/annurev-publhealth-032315-021556>
- Griffith, D. M., Bruce, M. A., & Thorpe, R. J. Jr. (2019). *Men's health equity: A handbook*. Routledge.
- Griffith, D. M., Gilbert, K. L., Bruce, M. A., & Thorpe, R. J. Jr. (2016). Masculinity in men's health: Barrier or portal to healthcare? In J. J. Heidelbaugh (Ed.), *Men's health in primary care* (pp. 19–31). Springer. https://doi.org/10.1007/978-3-319-26091-4_2
- Hamilton, D., Darity, W., Jr. Price, A. E., Sridharan, V., & Tippett, R. (2015). Umbrellas don't make it rain: Why studying and working hard isn't enough for Black Americans. *The New School*, 779, 780–781.
- Hickson, D. A., Diez Roux, A. V., Gebreab, S. Y., Wyatt, S. B., Dubbert, P. M., Sarpong, D. F., Sims, M., & Taylor, H. A. (2012). Social patterning of cumulative biological risk by education and income among African Americans.

- American Journal of Public Health*, 102(7), 1362–1369. <https://doi.org/10.2105/ajph.2011.300444>
- Hobfoll, S. E. (1989). Conservation of resources: A new attempt at conceptualizing stress. *American Psychologist*, 44(3), 513–524. <https://doi.org/10.1037/0003-066X.44.3.513>
- Hobfoll, S. E. (2011). Conservation of resources theory: Its implication for stress, health, and resilience. In S. Folkman (Ed.), *The Oxford handbook of stress, health, and coping* (pp. 127–147). Oxford University Press.
- Hudson, D. L., Banks, A., Holland, D., & Sewell, W. (2019). Understanding health inequalities experienced by Black men: Fundamental links between racism, socioeconomic position, and social mobility. In D. M. Griffith, M. A. Bruce, & R. J. Thorpe (Eds.), *Men's health equity*. Routledge.
- Hudson, D. L., Eaton, J., Lewis, P., Grant, P., Sewell, W., & Gilbert, K. (2016). “Racism?!? . . . Just look at our neighborhoods”: Views on racial discrimination and coping among African American men in Saint Louis *The Journal of Men's Studies*, 24(2), 130–150. <https://doi.org/10.1177/1060826516641103>
- Hudson, D. L., Bullard, K. M., Neighbors, H. W., Geronimus, A. T., Yang, J., & Jackson, J. S. (2012). Are benefits conferred with greater socioeconomic position undermined by racial discrimination among African American men? *Journal of Men's Health*, 9(2), 127–136. <https://doi.org/10.1016/j.jomh.2012.03.006>
- Hudson, D. L., Sacks, T. A., Sewell, W., Holland, D., & Gordon, J. (2020). Surviving the White space: Perspectives on how middle-class Black men navigate cultural racism. *Ethnic and Racial Studies*, 44(14), 2513–2531. <https://doi.org/10.1080/01419870.2020.1834594>
- Jackson, P. B., & Williams, D. R. (2006). The intersection of race, gender, and SES: Health paradoxes. In A. J. Schulz, & L. Mullings (Eds.), *Gender, race, class, & health: Intersectional approaches* (pp. 131–162). Jossey-Bass/Wiley.
- Lewis, T. T., Cogburn, C. D., & Williams, D. R. (2015). Self-reported experiences of discrimination and health: Scientific advances, ongoing controversies, and emerging issues. *Annual Review of Clinical Psychology*, 11(1), 407–440. <https://doi.org/10.1146/annurev-clinpsy-032814-112728>
- Marsh, K., Darity, W. A. Jr., Cohen, P. N., Casper, L. M., & Salters, D. (2007). The emerging Black middle class: Single and living alone. *Social Forces*, 86(2), 735–762.
- McNutt, L.-A., Wu, C., Xue, X., & Hafner, J. P. (2003). Estimating the relative risk in cohort studies and clinical trials of common outcomes. *American Journal of Epidemiology*, 157(10), 940–943. <https://doi.org/10.1093/aje/kwg074>
- Merkin, S. S., Karlamangla, A., Roux, A. D., Shrager, S., Watson, K., & Seeman, T. (2020). Race/ethnicity, neighborhood socioeconomic status and cardio-metabolic risk. *SSM Population Health*, 11, Article 100634. <https://doi.org/10.1016/j.ssmph.2020.100634>
- Meyer, I. H. (2003). Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychological Bulletin*, 129(5), 674–697. <https://doi.org/10.1037/0033-2909.129.5.674>
- National Health and Nutrition Examination Survey. (2018, December 14). *Analytic guidelines, 2011-2014 and 2015-2016*. <https://wwwn.cdc.gov/nchs/data/nhanes/analyticguidelines/11-16-analytic-guidelines.pdf>
- Nuru-Jeter, A. M., Michaels, E. K., Thomas, M. D., Reeves, A. N. Jr., R. J. T., & LaVeist, T. A. (2018). Relative roles of race versus socioeconomic position in studies of health inequalities: A matter of interpretation. *Annual Review of Public Health*, 39(1), 169–188. <https://doi.org/10.1146/annurev-publhealth-040617-014230>
- Phelan, J. C., Link, B. G., & Tehranifar, P. (2010). Social conditions as fundamental causes of health inequalities: Theory, evidence, and policy implications. *Journal of Health and Social Behavior*, 51(1_suppl), S28–S40.
- Rainisch, B. K. W., & Upchurch, D. M. (2013). Socio-demographic correlates of allostatic load among a national sample of adolescents: Findings from the National Health and Nutrition Examination Survey, 1999–2008. *Journal of Adolescent Health*, 53(4), 506–511.
- Rearson, S. F., Fox, L., & Townsend, J. (2015). Neighborhood income composition by household race and income, 1990–2009. *The ANNALS of the American Academy of Political and Social Science*, 660(1), 78–97. <https://doi.org/10.1177/0002716215576104>
- Rieker, P. P., & Bird, C. E. (2005). Rethinking gender differences in health: Why we need to integrate social and biological perspectives. *The Journals of Gerontology Series B*, 60(Special_Issue_2), S40–S47.
- Sacks, T. K. (2019). *Invisible visits: Black middle-class women in the American healthcare system*. Oxford University Press. <https://find.library.duke.edu/catalog/DUKE008750184>
- Seeman, T. E., Singer, B. H., Rowe, J. W., Horwitz, R. I., & McEwen, B. S. (1997). Price of adaptation—allostatic load and its health consequences: MacArthur studies of successful aging. *Archives of Internal Medicine*, 157(19), 2259–2268.
- Sternthal, M. J., Slopen, N., & Williams, D. R. (2011). Racial disparities in health: How much does stress really matter? *Du Bois Review: Social Science Research on Race*, 8(1), 95–113. <https://doi.org/10.1017/S1742058X11000087>
- Taylor, R. J., Miller, R., Mouzon, D., Keith, V. M., & Chatters, L. M. (2016). Everyday Discrimination among African American men: The impact of criminal justice contact. *Race and Justice*, 8(2), 154–177. <https://doi.org/10.1177/2153368716661849>
- Thomas, M. E., Moye, R., Henderson, L., & Horton, H. D. (2018). Separate and unequal: The impact of socioeconomic status, segregation, and the great recession on racial disparities in housing values. *Sociology of Race and Ethnicity*, 4(2), 229–244.
- Thorpe, R. J. Jr., Bell, C. N., Kennedy-Hendricks, A., Harvey, J., Smolen, J. R., Bowie, J. V., & LaVeist, T. A. (2015). Disentangling race and social context in understanding disparities in chronic conditions among men. *Journal of Urban Health*, 92(1), 83–92. <https://doi.org/10.1007/s11524-014-9900-9>
- Thorpe, R. J. Jr., Cobb, R., King, K., Bruce, M. A., Archibald, P., Jones, H. P., Norris, K. C., Whitfield, K. E., & Hudson,

- D. (2020). The association between depressive symptoms and accumulation of stress among Black men in the health and retirement study. *Innovation in Aging*, 4(5), Article igaa047. <https://doi.org/10.1093/geroni/igaa047>
- Thorpe, R. J. Jr., Parker, L. J., Cobb, R. J., Dillard, F., & Bowie, J. (2017). Association between discrimination and obesity in African-American men. *Biodemography and Social Biology*, 63(3), 253–261.
- Turner, R. J., Brown, T. N., & Hale, W. B. (2017). Race, socioeconomic position, and physical health: A descriptive analysis. *Journal of Health and Social Behavior*, 58(1), 23–36.
- Upchurch, D. M., Stein, J., Greendale, G. A., Chyu, L., Tseng, C. H., Huang, M. H., Lewis, T. T., Kravitz, H. M., & Seeman, T. (2015). A longitudinal investigation of race, socioeconomic status, and psychosocial mediators of allostatic load in midlife women: Findings from the study of women's health across the nation. *Psychosomatic Medicine*, 77(4), 402–412. <https://doi.org/10.1097/psy.000000000000175>
- Wheaton, F. V., Thomas, C. S., Roman, C., & Abdou, C. M. (2018). Discrimination and depressive symptoms among African American men across the adult lifecourse. *Journals of Gerontology B*, 73(2), 208–218. <https://doi.org/10.1093/geronb/gbx077>
- Wildeman, C., & Wang, E. A. (2017). Mass incarceration, public health, and widening inequality in the USA. *The Lancet*, 389(10077), 1464–1474. [https://doi.org/10.1016/S0140-6736\(17\)30259-3](https://doi.org/10.1016/S0140-6736(17)30259-3)
- Williams, D. R., Lawrence, J. A., & Davis, B. A. (2019). Racism and health: Evidence and needed research. *Annual Review of Public Health*, 40(1), 105–125. <https://doi.org/10.1146/annurev-publhealth-040218-043750>
- Wilson-Frederick, S. M., Thorpe, R. J. Jr., Bell, C. N., Bleich, S. N., Ford, J. G., & LaVeist, T. A. (2014). Examination of race disparities in physical inactivity among adults of similar social context. *Ethnicity and Disease*, 24(3), 363–369. <https://pubmed.ncbi.nlm.nih.gov/25065080>, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4351739/>
- Zou, G. (2004). A modified Poisson regression approach to prospective studies with binary data. *American Journal of Epidemiology*, 159(7), 702–706. <https://doi.org/10.1093/aje/kwh090>