



# Draft Genome Sequences of Five *Carnobacterium* sp. Strains Isolated from Freshwater Ponds in Belgium

Konrad J. Korzeniowski,<sup>a</sup> Shinjini Mukherjee,<sup>a</sup> Caroline Souffreau<sup>a</sup>

<sup>a</sup>Laboratory of Aquatic Ecology, Evolution and Conservation, KU Leuven, Leuven, Belgium

**ABSTRACT** Strains belonging to the genus *Carnobacterium* are Gram-positive bacteria that are widely distributed in the environment. Here, we report the draft genome sequences of five *Carnobacterium* strains isolated from freshwater ponds located in Flanders, Belgium, and sequenced on an Illumina HiSeq 4000 platform.

Bacteria from the genus *Carnobacterium* have previously been isolated from various food sources (1–3), such as meat (4) and milk (5), and grow in a pH range of 5.5 to 9.1 (1, 3, 6). Fewer reports exist of isolates of this genus from aquatic environments (1, 6–8). To date, 11 species have been reported to belong to the genus (3, 4, 9–13). Here, we present draft genome sequences of five *Carnobacterium* sp. strains isolated from five different freshwater ponds located in Belgium within the scope of experimental work. These ponds were small (<1 ha), shallow (<3 m), and man-made. Table 1 includes the geographic coordinates of the ponds and some water quality variables.

Water (50 ml) from each pond was filtered using a 0.22- $\mu$ m filter and cryopreserved in glycerol at  $-80^{\circ}\text{C}$  before the isolates were cultured on King agar B medium at  $24^{\circ}\text{C}$ . For DNA isolation, five isolates were cultivated at  $24^{\circ}\text{C}$  on King agar B medium for 48 h. Bacterial DNA was isolated using a Qiagen DNeasy UltraClean microbial DNA extraction kit (Qiagen). Initial identification was done by sequencing the 16S rRNA gene using primers 27F and 1492R (14).

The DNA concentration of the samples was measured using a Quant-iT PicoGreen kit (Thermo Fisher Scientific); DNA integrity and purity were determined by electrophoresis (1% agarose gel with Midori Green [Nippon Genetics Europe]) and spectrophotometry ( $A_{260}/A_{280}$ ; Infinite M Nano+ [Tecan]). Library preparation and sequencing were performed at BGI (Shenzhen, China). DNA was fragmented using a Covaris ultrasonicator. Fragmented DNA was combined with end repair mix (Agilent, USA) and purified with an AxyPrep Mag PCR cleanup kit (Thermo Fisher Scientific). Adapters (adapter 1, GATCGGAAGAGCACACGTCTGAACTCCAGTCAC; adapter 2, AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTGA) were added, and adapter-ligated DNA was purified with an AxyPrep Mag PCR cleanup kit. Adapter-ligated fragments (insert size, 300 bp) were enriched by PCR amplification ( $95^{\circ}\text{C}$  for 2 min; 6 cycles at  $98^{\circ}\text{C}$  for 15 s,  $62^{\circ}\text{C}$  for 30 s, and  $72^{\circ}\text{C}$  for 30s;  $72^{\circ}\text{C}$  for 5 min), and the PCR products were purified with an AxyPrep Mag PCR cleanup kit. Sequencing libraries were quantified using an Agilent Technologies 2100 bioanalyzer and an ABI StepOnePlus real-time PCR system. The resulting libraries (<800 bp) were sequenced on an Illumina HiSeq 4000 platform (150-bp paired ends) at BGI.

Trimmomatic v0.38 (15) was used for quality trimming of raw sequences with a Phred quality score cutoff of 33 and a minimum read length of 40. Unicycler v0.4.8 (16) software was used for *de novo* assembly of the genomes. Functional annotation and genome quality assessment were performed using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP). Genome coverage was assessed with BMAP v38.72

**Citation** Korzeniowski KJ, Mukherjee S, Souffreau C. 2020. Draft genome sequences of five *Carnobacterium* sp. strains isolated from freshwater ponds in Belgium. *Microbiol Resour Announc* 9:e00955-19. <https://doi.org/10.1128/MRA.00955-19>.

**Editor** J. Cameron Thrash, University of Southern California

**Copyright** © 2020 Korzeniowski et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Konrad J. Korzeniowski, [konrad.korzeniowski333@gmail.com](mailto:konrad.korzeniowski333@gmail.com).

**Received** 9 September 2019

**Accepted** 26 May 2020

**Published** 18 June 2020

**TABLE 1** Genome sequence statistics summary, pond water quality characteristics, and pond geographic coordinates for the five isolated *Carnobacterium* strains

Variable	Data for <i>Carnobacterium</i> sp. 1290 strain:				
	PL12RED10	PL26RED25	PL17RED31	PL24RED07	PL17GRE32
Genome characteristics					
Total length (bp)	1,976,998	1,974,582	1,957,158	1,975,794	1,996,575
GC content (%)	39.34	39.46	39.48	39.46	39.42
No. of CDS <sup>a</sup>	1,812	1,741	1,800	1,790	1,832
No. of tRNAs	51	51	51	51	52
No. of rRNAs	4	3	3	3	3
No. of reads	6,095,916	6,238,674	5,210,544	5,894,490	6,301,572
Final coverage (%)	457.353	468.178	396.568	444.213	471.107
SRA accession no. for raw reads	<a href="#">SRX7894432</a>	<a href="#">SRX7894431</a>	<a href="#">SRX7894430</a>	<a href="#">SRX7894429</a>	<a href="#">SRX7894428</a>
No. of scaffolds					
Scaffold $N_{50}$ (bp)	151,666	74,489	168,268	74,489	119,417
GenBank accession no. for assembled genomes	<a href="#">WVES00000000.1</a>	<a href="#">WVEP00000000.1</a>	<a href="#">WVEQ00000000.1</a>	<a href="#">WVER00000000.1</a>	<a href="#">WVEO00000000.1</a>
Pond water quality data					
pH	10.14	7.16	6.93	8.91	7.39
Oxygen (mg/liter)	12.495	3.905	2.755	8.94	1.00
Chlorophyll-a ( $\mu\text{g/liter}$ )	72.87	59.52	97.92	219.04	41.90
Pond geographic coordinates					
	51°7'1.7796"N, 4°28'30.4392"E	51°10'18.39"N, 4°24'53.9496"E	50°59'27.6684"N, 3°59'50.0604"E	50°56'5.784"N, 4°51'30.8304"E	51°1'2.7984"N, 3°56'36.3012"E

<sup>a</sup> CDS, coding sequences.

(<https://sourceforge.net/projects/bbmap/>). The sequence statistics of the genomes are summarized in Table 1.

Pairwise comparison analysis using JSpeciesWS (17) (<http://jspecies.ribohost.com/jspeciesws/>) showed that all investigated strains belong to the same species (average nucleotide identity, >95%) and have highest similarity (>97.39%) to *Carnobacterium* sp. 1290. Based on ResFinder (<https://cge.cbs.dtu.dk/services/ResFinder/>) (18), no resistance genes were found. No intact phages were found using PHASTER (19) (<http://phaster.ca/>). Default parameters were used for all software.

**Data availability.** This genome sequencing project has been deposited in the NCBI Sequence Read Archive (SRA) under BioProject number [PRJNA551600](#) and BioSample numbers [SAMN12161540](#) to [SAMN12161544](#). The accession numbers for the individual strains are given in Table 1.

## ACKNOWLEDGMENTS

This study was funded by the KU Leuven Research Fund (project C/16/17/002) and the National Fund for Scientific Research–Flanders (FWO) project G061916N. K.J.K. received a Ph.D. fellowship (FWO grant number 1501320N).

## REFERENCES

- Leisner JJ, Laursen BG, Prévost H, Drider D, Dalgaard P. 2007. *Carnobacterium*: positive and negative effects in the environment and in foods. *FEMS Microbiol Rev* 31:592–613. <https://doi.org/10.1111/j.1574-6976.2007.00080.x>.
- Leroi F, Matamoros S, Pilet MF, Gigout F, Pre H. 2009. Selection and evaluation of seafood-borne psychrotrophic lactic acid bacteria as inhibitors of pathogenic and spoilage bacteria. *Food Microbiol* 26: 638–644. <https://doi.org/10.1016/j.fm.2009.04.011>.
- Yost CK, Guan TY, Holley RA, Peirson M. 2002. *Carnobacterium viridans* sp. nov., an alkaliphilic, facultative anaerobe isolated from refrigerated, vacuum-packed bologna sausage. *Int J Syst Evol Microbiol* 52: 1881–1885. <https://doi.org/10.1099/00207713-52-5-1881>.
- Collins MD, Farrow JAE, Phillips BA, Ferusu S, Jones D. 1987. Classification of *Lactobacillus divergens*, *Lactobacillus piscicola*, and some catalase-negative, asporogenous, rod-shaped bacteria from poultry in a new genus, *Carnobacterium*. *Int J Syst Evol Microbiol* 37:310–316. <https://doi.org/10.1099/00207713-37-4-310>.
- Mora D, Scarpellini M, Franzetti L, Colombo S, Galli A. 2003. Reclassification of *Lactobacillus maltaromicus* (Miller et al. 1974) DSM 20342T and DSM 20344 and *Carnobacterium piscicola* (Collins et al. 1987) DSM 20730T and DSM 20722 as *Carnobacterium maltaromaticum* comb. nov. *Int J Syst Evol Microbiol* 53:675–678. <https://doi.org/10.1099/ijs.0.02405-0>.
- Nicholson WL, Krivushin K, Gilichinsky D, Schuerger AC. 2013. Growth of *Carnobacterium* spp. from permafrost under low pressure, temperature, and anoxic atmosphere has implications for Earth microbes on Mars. *Proc Natl Acad Sci U S A* 110:666–671. <https://doi.org/10.1073/pnas.1209793110>.
- Yanagida F, Chen Y-S, Yasaki M. 2007. Isolation and characterization of lactic acid bacteria from lakes. *J Basic Microbiol* 47:184–190. <https://doi.org/10.1002/jobm.200610237>.
- Voget S, Klippel B, Daniel R, Antranikian G. 2011. Complete genome sequence of *Carnobacterium* sp. 17-4. *J Bacteriol* 193:3403–3404. <https://doi.org/10.1128/JB.05113-11>.

9. Joborn A, Dorsch M, Olsson JC, Westerdahl A, Kjelleberg S. 1999. *Carnobacterium inhibens* sp. nov., isolated from the intestine of Atlantic salmon (*Salmo salar*). *Int J Syst Bacteriol* 49:1891–1898. <https://doi.org/10.1099/00207713-49-4-1891>.
10. Pikuta EV, Marsic D, Bej A, Tang J, Krader P, Hoover RB. 2005. *Carnobacterium pleistocenium* sp. nov., a novel psychrotolerant, facultative anaerobe isolated from permafrost of the Fox Tunnel in Alaska. *Int J Syst Evol Microbiol* 55:473–478. <https://doi.org/10.1099/ijs.0.63384-0>.
11. Franzmann PD, Höpfl P, Weiss N, Tindall BJ. 1991. Psychrotrophic, lactic acid-producing bacteria from anoxic waters in Ace Lake, Antarctica; *Carnobacterium funditum* sp. nov. and *Carnobacterium alterfunditum* sp. nov. *Arch Microbiol* 156:255–262. <https://doi.org/10.1007/BF00262994>.
12. Kim M-S, Roh SW, Nam Y-D, Yoon J-H, Bae J-W. 2009. *Carnobacterium jeotgali* sp. nov., isolated from a Korean traditional fermented food. *Int J Syst Evol Microbiol* 59:3168–3171. <https://doi.org/10.1099/ijs.0.010116-0>.
13. Snauwaert I, Hoste B, De Bruyne K, Peeters K, De Vuyst L, Willems A, Vandamme P. 2013. *Carnobacterium iners* sp. nov., a psychrophilic, lactic acid-producing bacterium from the littoral zone of an Antarctic pond. *Int J Syst Evol Microbiol* 63:1370–1375. <https://doi.org/10.1099/ijs.0.042861-0>.
14. Deng W, Wanapat M, Ma S, Chen J, Xi D, He T, Yang Z, Mao H. 2007. Phylogenetic analysis of 16S rDNA sequences manifest rumen bacterial diversity in gayals (*Bos frontalis*) fed fresh bamboo leaves and twigs (*Sinarumdinaria*). *Asian Australas J Anim Sci* 20:1057–1066. <https://doi.org/10.5713/ajas.2007.1057>.
15. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
16. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
17. Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
18. Zankari E. 2014. Comparison of the Web tools ARG-ANNOT and ResFinder for detection of resistance genes in bacteria. *Antimicrob Agents Chemother* 58:4986. <https://doi.org/10.1128/AAC.02620-14>.
19. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.