

High proportion of Guillain-Barré syndrome associated with chikungunya in Northeast Brazil

Aline de Moura Brasil Matos, MD, Fernanda Martins Maia Carvalho, MD, PhD, Danielle Lima Malta, PhD, Cleonísio Leite Rodrigues, MD, PhD, Alvina Clara Félix, MSc, Claudio Sergio Pannuti, PhD, Amanda Dias da Rocha Lima, MSc, Danilo Lucas Alves Espósito, PhD, Leonilda Maria Barbosa dos Santos, PhD, Felipe von Glehn, MD, PhD, FAAN, Jeová Keny Baima Colares, MD, PhD, Benedito Antônio Lopes da Fonseca, MD, PhD, Augusto César Penalva de Oliveira, MD, PhD, and Camila Malta Romano, PhD

Correspondence
Dr. Romano
cmromano@usp.br

Neurol Neuroimmunol Neuroinflamm 2020;7:e833. doi:10.1212/NXI.0000000000000833

From 2013 to 2015, sanitary authorities reported an increased incidence of Guillain-Barré syndrome (GBS) associated with Zika virus (ZIKV) in French Polynesia, Caribbean, and Brazil.^{1–3} After the end of ZIKV epidemics, GBS cases were still above the usual limits in countries where the arrival of chikungunya virus (CHIKV) was also a concern.³

Here, we report the findings from Hospital Geral de Fortaleza (HGF), a neuroinvasive arboviral disease vigilance center in Ceará State, Northeast Brazil.

Methods

We performed a prospective observational study that enrolled patients aged 15 year or older with the diagnosis of GBS⁴ (Brighton criteria I or II). All consecutive patients fulfilling the inclusion criteria from May 2015 to December 2017 were invited to participate.

Patients were evaluated for demographics, clinics (at admission and 6 months later), serum, and CSF complementary tests, and EMG. Owing to the local epidemics, besides investigating for classic GBS triggers, virologic tests for dengue virus, ZIKV, and CHIKV (i.e., ELISA IgM, IgG, and specific real-time PCR) were performed by a researcher blinded to clinical results. Neurofilament light chain⁵ (NfL) was measured in CSF and related to death, need for mechanical ventilation (MV), and incomplete recovery. CSF from 10 healthy subjects served as the control for NfL.

For comparison, official reports of arboviral systemic infections and total GBS cases/year from 2013 to 2017 were requested to the local state government according the Law #12.527 November 18, 2011.

Continuous data were summarized as median and interquartile range (IQR). Categorical data were presented as counts and percentages. Kendall τ was used for correlations and Mann-Whitney test for comparisons. p values <0.05 were deemed statistically significant. Data were analyzed using SPSS version 25.0. Graphs were constructed using Sigma Plot version 11.0. The study was approved by the HGF ethics committee (CAAE: 56572316.9.0000.5040) and the USP ethics committee (CAAE: 00274418.7.0000.0068) and conducted according to appropriate Brazilian regulations. All subjects provided written informed consent.


From the Universidade de São Paulo (A.d.M.B.M., A.C.F., C.S.P., C.M.R.), Instituto de Medicina Tropical, Faculdade de Medicina; Hospital Geral de Fortaleza (F.M.M.C., C.L.R.), Serviço de Neurologia, CE; Universidade de Fortaleza (F.M.M.C., D.L.M., J.K.B.C.), Programa de Pós-Graduação em Ciências Médicas, CE; Universidade de Campinas (A.D.d.R.L., L.M.B.d.S., F.V.G.), Instituto de Biologia, SP; Universidade de São Paulo (D.L.A.E., B.A.L.d.F.), Faculdade de Medicina de Ribeirão Preto; Instituto de Infectologia Emílio Ribas (A.C.P.d.O.), São Paulo; and Laboratório de Virologia (LIMHC 52) (C.M.R.), Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, Brazil.

Go to Neurology.org/NN for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by PROFAP-HCLIM.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Figure 1 Notified cases of dengue, Zika, and chikungunya in Ceará state and GBS cases in the study site



CHIKV = Chikungunya virus; DENV = Dengue virus; GBS = Guillain-Barré syndrome; ZIKV = Zika virus.

Results

A total of 42 patients with GBS suspicion were admitted. Eight patients were excluded (figure e-1, links.lww.com/NXI/A278). From the 34 remaining, GBS trigger was attributed to CHIKV for 9 patients.⁶ For those, median age was 47 (IQR 31–55), 56% were men and most without comorbidities (56%). The mean number of GBS cases in the previous 2 years of the study was 52 cases/year; from 2015 to 2017, it was 88 cases/year. Most CHIKV-GBS cases concentrated in the CHIKV epidemic peak (figure 1).

At admission, patients were confined to bed/wheelchair (56%), walking with support (33%), or unable to run (11%). During in-hospital stay, 22% required MV and no patient died. Median days in hospital were 16 (IQR 12–25.5). Recovery after 6 months was complete for 33%, whereas 67% remained with minor signs or symptoms (table e-1, links.lww.com/NXI/A278).

Major laboratory results can be found in supplementary material (links.lww.com/NXI/A278). EMG was primary demyelinating (75%) or primary axonal (25%). Although NfL presented higher titles than control (figure e-2), there was no correlation with death, MV or, recovery.

Discussion

We found that 26% of the cases enrolled were associated with CHIKV as an infectious trigger. The increase was coincident with the first local epidemics of CHIKV and followed a ZIKV epidemics. The association might be the result of a molecular mimicry autoimmune mechanism because CHIKV E1 glycoprotein shares homology with contactin-2,⁷ a protein present in the juxtaparanode.⁸

In one of the larger GBS cohort available,⁹ although no laboratory tests are mentioned, there are no reports of rash/arthritis as prodromal symptoms. Despite that, reports of GBS-ZIKV associations are well known,^{1,3} unlike GBS-CHIKV which are rarely reported. Regarding clinical outcomes, differences are also apparent from our GBS-CHIKV to this same cohort. For recovery, all of our patients achieved Huges score of “0” or “1” in 6 months (vs 61%). No patient with GBS-CHIKV died (vs 7%). As for MV, we have similar numbers (22% vs 19%).

A limitation of our study is the small sample size. This was unavoidable because of the rare nature of GBS disease and the seasonality of arbovirus infections. In addition, we were not able to access antiganglioside tests. However, we performed a clinical follow-up of at least 6 months and adopted strict criteria for GBS, allowing proper exclusion of diagnostic mimics.

Our findings suggest CHIKV as an important trigger for GBS during epidemics, overcoming classic triggers as *Campylobacter jejuni*, Epstein-Barr virus, and influenza virus. Good outcomes were a commonplace in our study; however, sanitary authorities of areas affected by CHIKV should be aware of a possible increase in GBS incidence and as a consequence an increased necessity for intensive care unit beds and rehabilitation treatments.

Acknowledgment

The authors would like to thank Professor Steven S. Witkin from Weill Cornell Medicine for constructive comments and kindness in editing the manuscript, Vânia Maria Alves de Araujo from the Hospital Geral de Fortaleza for the laboratory assistance, and Rosa Maria Nascimento Marcusso from Instituto de Infectologia Emilio Ribas for the organizational support and revision of statistical analysis.

Study funding

CNPq/Brazil under the grant number 2016/407429 and Fundação de Amparo à Pesquisa do Estado de São Paulo-FAPESP/Brazil under the grants numbers 2014/26431-0 and 2019/03859-9.

Disclosure

A.d.M.B. Matos reports no disclosures. F.M. Maia Carvalho received research financial support from Conselho Nacional de Desenvolvimento Científico e Tecnológico CNPq/Brazil under the grant number 2016/407429. D.L. Malta, C.L. Rodrigues, A.C. Félix, C.S. Pannuti, A.d.R. Lima, and D.L.A. Espósito report no disclosures. L.M.B. dos Santos received research financial support from Fundação de Amparo à Pesquisa FAPESP/Brazil under the grant number 2014/26431-0. F. von Glehn, J.K.B. Colares, B.A.L. da Fonseca, and A.C.P. de Oliveira report no disclosures. C.M. Romano received research financial support from Programa de fomento as atividades de lideranças científicas dos LIMs do Hospital das Clínicas (PROFAP-LIM/HCFMUSP) #10/2020. Go to Neurology.org/NN for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* January 21, 2020. Accepted in final form May 26, 2020.

Appendix Authors

Name	Location	Contribution
Aline de Moura Brasil Matos, MD	Universidade de São Paulo	Study concept and design, data acquisition, analysis and interpretation of the data, and critical revision of the manuscript from important intellectual content.
Fernanda Martins Maia Carvalho, MD, PhD	Hospital Geral de Fortaleza and Universidade de Fortaleza	Study concept and design, data acquisition, analysis and interpretation of the data, and critical revision of the manuscript from important intellectual content.

Appendix (continued)

Name	Location	Contribution
Danielle Lima Malta, PhD	Universidade de Fortaleza	Study concept and design, data acquisition, and critical revision of the manuscript from important intellectual content.
Cleoniso Leite Rodrigues, MD, PhD	Hospital Geral de Fortaleza	Study concept and design, data acquisition, and critical revision of the manuscript from important intellectual content.
Alvina Clara Félix, MSc	Universidade de São Paulo	Data acquisition and critical revision of the manuscript from important intellectual content.
Claudio Sergio Pannuti, PhD	Universidade de São Paulo	Critical revision of the manuscript from important intellectual content.
Amanda Dias da Rocha Lima, MSc	Universidade de Campinas	Data acquisition and critical revision of the manuscript from important intellectual content.
Danilo Lucas Alves Espósito, PhD	Universidade de São Paulo, Ribeirão Preto	Data acquisition and critical revision of the manuscript from important intellectual content.
Leonilda Maria Barbosa dos Santos, PhD	Universidade de Campinas	Data acquisition and critical revision of the manuscript from important intellectual content.
Felipe von Glehn, MD, PhD, FAAN	Universidade de Campinas	Data acquisition and critical revision of the manuscript from important intellectual content.
Jeová Keny Baima Colares, MD, PhD	Universidade de Fortaleza	Critical revision of the manuscript from important intellectual content.
Benedito Antônio Lopes da Fonseca, MD, PhD	Universidade de São Paulo, Ribeirão Preto	Data acquisition and critical revision of the manuscript from important intellectual content.
Augusto César Penalva de Oliveira, MD, PhD	Instituto de Infectologia Dr. Emilio Ribas, São Paulo	Study concept and design, analysis and interpretation of the data, and critical revision of the manuscript from important intellectual content.
Camila Malta Romano, PhD	Universidade de São Paulo and Hospital das Clínicas, São Paulo	Study concept and design, data acquisition, analysis and interpretation of the data, and critical revision of the manuscript from important intellectual content.

References

1. Cao-Lormeau VM, Blake A, Mons S, et al. Guillain-Barre syndrome outbreak associated with Zika virus infection in French polynesia: a case-control study. *Lancet* 2016; 387:1531–1539.
2. Arias A, Torres-Tobar L, Hernández G, et al. Guillain-Barré syndrome in patients with a recent history of Zika in Cucuta, Colombia: a descriptive case series of 19 patients from December 2015 to March 2016. *J Crit Care* 2017;37:19–23.
3. Mehta R, Soares CN, Medialdea-Carrera R, et al. The spectrum of neurological disease associated with Zika and chikungunya viruses in adults in Rio de Janeiro, Brazil: a case series. *PLoS Negl Trop Dis* 2018;12:e0006212.
4. Fokke C, van den Berg B, Drenthen J, Walgaard C, van Doorn PA, Jacobs BC. Diagnosis of Guillain-Barre syndrome and validation of Brighton criteria. *Brain* 2014; 137:33–43.
5. Rajabally YA, Uncini A. Outcome and its predictors in Guillain-Barre syndrome. *J Neurol Neurosurg Psychiatry* 2012;83:711–718.
6. Center for Disease Control and Prevention. Arboviral diseases, neuroinvasive and non-neuroinvasive 2015 case definition. In: *NNDSS*, ed. Available at: www.cdc.gov/nndss/conditions/arboviral-diseases-neuroinvasive-and-non-neuroinvasive/case-definition/2015/2015. Accessed January 2020.
7. Reddy V, Desai A, Krishna SS, Vasanthapuram R. Molecular mimicry between chikungunya virus and host components: a possible mechanism for the arthritic manifestations. *PLoS Negl Trop Dis* 2017;11:e0005238.
8. Masuda T. Contactin-2/TAG-1, active on the front line for three decades. *Cell Adhes Migration* 2017;11:524–531.
9. Doets AY, Verboon C, van den Berg B, et al. Regional variation of Guillain-Barre syndrome. *Brain* 2018;141:2866–2877.