

Female genital mutilation — a blind spot in Dutch general practice? A case–control study

Ramin Kawous^{1,2*}, Nigar Kerimova³, Maria ETC van den Muijsenbergh^{2,3}

¹Department of Public Health, Erasmus University Medical Centre, Rotterdam, The Netherlands; ²Pharos, Dutch Centre of Expertise on Health Disparities, Utrecht, The Netherlands; ³Department of Primary and Community Care, Radboud University Medical Centre, Nijmegen, The Netherlands

Abstract

Background: Women with female genital mutilation or cutting (FGM/C) often suffer from physical and psychosexual problems related to FGM/C. As gatekeepers to the medical system, GPs are often the first to be consulted about these problems. It is as yet unknown if, and to what extent, Dutch GPs identify women with FGM/C or related health problems.

Aim: To investigate how often Dutch GPs register FGM/C and related health problems.

Design & setting: A case–control study of anonymised patient records was performed in the Netherlands.

Method: Medical records were checked for information on country of origin. Records of women, aged ≥15 years, from countries where FGM/C is practised were compared with those of a case-control.

Results: Although many migrants were registered with the participating GPs, information on country of origin was seldom recorded. Only 68 out of 16 700 patients were identified as women from countries where FGM/C is practised; 12 out of these 68 records contained information about the FGM/C status, but none on the type of FGM/C. There were no significant differences in health problems related to FGM/C between patients with FGM/C and the controls.

Conclusion: FGM/C may be a blind spot for GPs and registration of information on migration background could be improved. A larger sample in a future study is needed to confirm this finding. Given the growing global migration, awareness and knowledge on FGM/C, and other migration-related health issues should be part of GP training.

*For correspondence: r.
kawous@outlook.com

Competing interest: The authors declare that no competing interests exist.

Received: 16 May 2020

Accepted: 21 May 2020

Published: 02 December 2020

©This article is Open Access: CC BY license (<https://creativecommons.org/licenses/by/4.0/>)

Author Keywords: female genital mutilation/cutting, general practice, migrant women, primary health care, culturally competent care

Copyright © 2020, The Authors;
DOI:10.3399/
bjgpopen20X101105

How this fits in

The majority of previous research has focused on FGM/C in maternity settings. There is a need for more research on FGM/C in primary care such as general practice.¹ This is the first study to investigate the provision of health care to women with FGM/C in Dutch general practice.

Introduction

FGM/C consists of ‘all procedures that involve partial or total removal of the external female genitalia or other injury to the female genital organs for non-medical reasons’.² The World Health Organization (WHO) has classified four main types of FGM/C: type I, or clitoridectomy, refers to excision of the prepuce, with or without excision of part or all of the clitoris; type II, or excision, which consists of the clitoris with partial or total removal of the labia minora; type III, the most severe form of FGM/C,

which is known as infibulation, refers to excision of part or all of the external genitalia and stitching or narrowing of the vaginal opening. Type IV is unclassified. It includes all other harmful procedures to the female genitalia for non-medical purposes; for example, pricking, piercing, incising, scraping, and cauterising the genital area.³

FGM/C is predominantly practised in countries in Africa, the Middle East, and Asia. Globally, it is estimated that more than 200 million girls and women have undergone FGM/C,⁴ and that an estimated 68 million girls are at risk of being cut between 2015 and 2030.⁵ As a result of increasing migration of women from countries where FGM/C is concentrated, FGM/C has become common in high-resource countries such as the Netherlands. It is estimated that approximately 41 000 girls and women with FGM/C are living in the Netherlands. About 37% of these girls and women are estimated to have been infibulated.⁶

FGM/C is recognised as a human rights violation² and many countries have increasingly undertaken law reform to prohibit FGM/C.⁷ In the Netherlands, performing any form of FGM/C is forbidden and punishable by law, with an incarceration term of up to 12 years. The law also prohibits any Dutch nationals or permanent Dutch residents from performing FGM/C abroad.^{8–10} However, there is no indication of FGM/C being performed in the Netherlands and no suspected FGM/C case has, as yet, been convicted by a court.¹¹

The Dutch action plan against the practice of FGM/C was established in 2006. The rationale behind the Dutch approach against FGM/C is to prevent FGM/C by accomplishing a behavioural change towards the practice. The prevention policies in the Netherlands consist of, among others, awareness raising on FGM/C among communities concerned and the empowerment of women, as well as training of health professionals and healthcare workers to support women affected by FGM/C, and to identify and prevent girls from being cut.¹² Healthcare providers, social care professionals, and teachers who are concerned that a girl may be at risk of FGM/C have legal obligation to report this.

Recently, the Dutch Society of Obstetrics and Gynaecology (NVOG) has developed guidelines on the management of FGM/C.¹³ Healthcare providers in primary and secondary health care are also able to record FGM/C and they have been urged to practise good record-keeping. Unfortunately, to date, no reliable data are available in the Netherlands regarding FGM/C in primary and secondary care.

FGM/C has been associated with adverse short- and long-term health consequences, including, severe pain, excessive bleeding, urinary tract infections, bacterial vaginosis, painful sexual intercourse, and adverse perinatal outcomes,^{14–16} as well as negative effects on mental health.^{17–19} The migration from countries where FGM/C is practised to high-resource countries will further increase.^{20,21} Consequently, in the near future, healthcare providers are expected to provide care for more women with FGM/C. There is increasing evidence that women with FGM/C receive lower quality of care, owing to *'lack of knowledge, cultural sensitivities associated with the subject leading to silence, stigma and inaction'*.¹ Since most of these studies have been conducted in obstetric settings,^{22–25} there is a need for more research on FGM/C in primary care, such as in general practice.

In the Netherlands, the GP is often the first to encounter health problems of women with FGM/C. GPs function as a 'gatekeeper' to the healthcare system in the Netherlands and all residents are obliged by their (mandatory) health insurance to be registered with a general practice. However, it is as yet unknown if, and to what extent, GPs identify women with FGM/C, nor what care they provide to them. Insight into these aspects are vital in order to evaluate whether any inequities exist in primary health care for women with FGM/C. The aim of this study was to investigate how often in Dutch general practice FGM/C-related health problems were recorded in women from countries where FGM/C is practised, compared with women with other migration background of the same age.

Method

Study design

A case–control study of anonymised patient records of women with a migration background, including women from countries where FGM/C is practised, was carried out between November and December 2017.

Recruitment and study population

The study was executed in five general practices in the Netherlands, known to care for a large migrant community and trained to care for women with FGM/C. In the Netherlands, it is becoming more difficult to encourage GPs to participate in surveys, owing to their demanding work schedules and increasing frequency of being approached for surveys.²⁶ Therefore, the practices were recruited through the researcher's informal network. A purposive sampling method was used, aiming for practices with a high number of patients from very high FGM/C prevalence countries, striving for variation among GPs in terms of sex, size, and geographical location of the general practice.

Patient records of girls and women, aged ≥ 15 years, from very high FGM/C prevalence countries, including Somalia, Egypt, Eritrea, Sierra Leone, and Sudan, were included. The Netherlands has a large number of migrants from these countries in which more than 80% of girls and women of reproductive age have undergone FGM/C. Since country of origin is not routinely registered in general practice, all female patients with a foreign surname were first included. Identifying migrants by means of their surname has proved to be a reliable, second-best method.²⁷ Next, information on country of origin was searched through the first 10 contact records, including the first visit records. Then, notes were searched to retrieve information on country of origin, using the keywords 'origin', 'country of birth' or 'country of origin', 'Somalia', 'Eritrea', 'Egypt', 'Sudan', or 'Sierra Leone'. The data of these women were compared with those of a case-control: female migrant patients in the same age group from countries where FGM/C is not practised. Every next migrant woman from a non-FMG/C practising country was selected from the same age-group on the list after each woman from a country where FGM/C is practised was selected.

It has been well documented that migrants, in general, have at some point differing morbidity patterns and receive lower quality of care than native populations.²⁸ Hence, within the scope of this study, it was found to be more appropriate to compare the data within migrant subgroups, so data of women from countries where FGM/C is practised were compared with female migrant patients from countries where FGM/C is not practised.

Data collection

The researcher (NK), a senior medical student, was allowed to access the complete patient records in each participating general practice, after signing the required confidentiality agreement, in line with national legislation on data protection. NK selected the records based on patients' surnames and then anonymised the data by converting each patient's personal details into an alphanumeric code. Records contained information on every contact with the GP (in person or by telephone), medication prescribed by the GP, and all letters to the GP from other (health) professionals and services.

It was checked whether country of origin, FGM/C status of the patient, type of FGM/C, or health problems often associated with FGM/C were recorded in medical files. The files were searched for health problems, including recurring urinary tract infections (defined as two or more cystitis in 6 months), dysmenorrhoea, sub- or in-fertility, complications during childbirth, painful sexual intercourse, and fear of sexual intercourse. In addition, it was checked whether the FGM/C status of the patient was recorded according to the International Classification of Primary Care (ICPC). The ICPC code-X82

Table 1 Number of women in the FGM/C and control groups in five general practices in the Netherlands

Control group (n = 61)		FGM/C group (n = 68)	
	n		n
Country of origin			
Afghanistan	2	Egypt	8
Armenia	3	Eritrea	7
Azerbaijan	1	Sierra Leone	4
China	3	Somalia	40
Morocco	39	Sudan	9
Pakistan	1		
Syria	1		
Taiwan	2		
Turkey	7		
Vietnam	2		
Mean age, years	39.82		36.13
SD	13.52		14.76
Range	18–74		15–74

FGM/C = female genital mutilation or cutting; SD = standard deviation

refers to injury to the female genital organs. Finally, women with unknown origin were excluded and the data were anonymised prior to analysis.

Data analysis

The data were analysed using SPSS statistical software (version 22). The χ^2 test was used to compare the data from different groups and to determine whether the differences were statistically significant.

Results

A total of 16 700 patient records from five general practices were included in the analysis. Country of origin of patients from FGM/C-prevalent countries was mentioned in 68 cases. As shown in **Table 1**, these women were from (in descending order) Somalia (58.8%), Sudan (13.2%), Egypt (11.8%), Eritrea (10.3%), and Sierra Leone (5.9%), with an average age of 36.18 years (standard deviation [SD] = 14.76). The majority of women in the control group were from Morocco (63.9%), with an average age of 39.82 years (SD = 13.52).

The FGM/C status of patients was recorded in 12 out of the 68 cases where country of origin was documented. Of these patients, 11 had undergone some form of FGM/C. In none of these cases was type of FGM/C classified and recorded. Moreover, the FGM/C status of one patient was recorded using the ICPC code-X82. Other cases were randomly recorded or found in the correspondence.

There were no significant differences in recorded health problems related to FGM/C such as urinary tract infections, dysmenorrhoea, sub- or in-fertility, complications during childbirth, painful sexual intercourse, or fear of sexual intercourse between patients with FGM/C ($n = 12$), patients whose FGM/C status was not recorded ($n = 56$) and patients in the control-group ($n = 61$).

Discussion

Summary

In this first study on FGM/C in five general practices in The Netherlands known to care for a large migrant community, patient demographics such as country of origin were seldom documented in the medical charts. Only 68 out of 16 700 patients were identified as women from very high FGM/C prevalence countries; 12 out of these 68 medical files contained information about the FGM/C status of the patient, and in none of these cases was type of FGM/C classified.

Although the participating practices were known to care for a large migrant community and trained to care for women with FGM/C, it is unclear why information on country of origin was seldom recorded and only 12 records contained information about FGM/C. It is believed that it is highly unlikely that only 12 out of 68 women from very high FGM/C prevalence countries had actually undergone FGM/C. FGM/C may be a blind spot in general practice in the Netherlands. Future research, preferably qualitative, is needed with larger samples to confirm this, and to explore the factors that encourage and hinder GPs in recording and discussing FGM/C with women who have undergone the procedure.

Strengths and limitations

The relatively small numbers of general practices included in the study is acknowledged, which may explain the non-significant difference in recorded health problems related to FGM/C between patients in the FGM/C group and patients in the control group. While healthcare providers are able to systematically record FGM/C, to date, no reliable data are available regarding FGM/C in primary and secondary care. Therefore, the authors believe that it may not be necessary to include more practices before concluding that there is an urgent need for improvement of recognition and recording of FGM in general practice; improvement could be achieved by discussing FGM/C with patients from FGM/C-prevalent countries, and by correctly classifying and recording FGM/C in the medical charts.

Comparison with existing literature

There is limited evidence on FGM/C in primary care. Most studies have been conducted in secondary care and have focused on FGM/C in maternity settings.¹ Nonetheless, the present findings would be consistent with results from available studies among other healthcare providers caring for women with FGM/C in secondary care.²⁹ Several reasons may explain the few recorded cases of FGM/C in

general practice. In general, it is known that most women rarely proactively discuss FGM/C with their healthcare provider. At the same time, healthcare providers refrain from asking about it.^{1,17,30-33} The possible explanation could be owing to cultural taboos around FGM/C in women's own community or fear of being judged, or being fearful as FGM/C is forbidden in their new country of residence.³⁴

In a comprehensive systematic review, Evans *et al*¹ have explored factors influencing the provision of health care related to FGM/C from the perspective of health providers. Feeling of embarrassment, shock, disgust, and horror; uncertainty about how to frame the questions, or '*anxiety about being perceived as culturally insensitive*', language barriers, and a lack of knowledge have been described as barriers to engagement with women with FGM/C.^{1,22-25,30,32,33,35-43}

Because of the sensitive nature of FGM/C, talking about the subject can make health providers feel uncomfortable. Some healthcare providers reported they avoided discussing the topic with their patient, as they did not want to offend, stigmatise, or jeopardise their relationships, assuming that women would initiate a conversation about FGM/C if there was a problem.¹ As a result, FGM/C will probably never be discussed at all, hence not recorded. As mentioned, previous studies have reported expression of strong emotions by healthcare providers, including shock, disgust, and horror. Cutting of women was perceived as an alien and negative practice, with healthcare providers describing and perceiving these women as not 'normal', but mutilated. Also, lack of knowledge has been listed repeatedly as a barrier to identify and manage women with FGM/C.^{1,22-24} For instance, Chalmers *et al*²² reported dissatisfaction of women with care and a perceived lack of knowledge and ability by healthcare providers to care for women during pregnancy and childbirth. Relph *et al*²³ assessed the knowledge, attitude, and training on FGM/C among medical and midwifery professionals and found that although the majority of these professionals were aware of the practice, their ability to identify FGM/C and its associated health complications remains unsatisfactory. Similarly, Zaidi *et al*²⁴ observed deficiencies in knowledge and adherence to guidelines on the management of health complications of FGM/C among healthcare providers. Zenner *et al*²⁵ reported on the quality of obstetric and midwifery care, and noted major deficits in identification, management, and safeguarding.

Implications for research and practice

The findings, along with those of previous reports, support the need for integration of FGM/C and culturally sensitive care in training programmes for healthcare providers. Currently, despite the growing number of books on migration health in primary care,⁴⁴⁻⁴⁷ the subject of FGM/C is not included in the curriculum of most medical, midwifery, and public health training programmes.^{48,49} In addition, specific guidelines on the management of health complications related to FGM/C are not well known among healthcare providers.⁴⁸

As noted by Abdulcadir *et al*,²⁹ recognition of FGM/C is the first step in formulating a diagnosis, before providing information and treating a patient with FGM/C. However, FGM/C may be a blind spot in general practice in the Netherlands. Future research, preferably qualitative, is needed with larger samples to confirm this and to explore the factors that encourage and hinder GPs in recording and discussing FGM/C with women who have undergone it.

Further, registration of information on migration background could be improved and, as migration is on the rise and the population attending general practices gets more and more diverse, GPs should not only be aware and knowledgeable on FGM/C, but also on other migration-related health issues.

Funding

The authors received no specific funding for this work.

Ethical approval

Agreement of all GPs was received after information was given about the study to have insight in the patient records. The information collected from patient records was anonymised and stored safely. Agreement of all the patients was considered implicit due to the fact that getting explicit agreement was very difficult, if not impossible. According to the Dutch law and regulations, this project was exempted from the need for approval through the Ethical committee of the Radboudumc, Nijmegen, because this study involved anonymised patient records.

Provenance

Freely submitted; externally peer reviewed.

Acknowledgements

The authors are grateful to all general practitioners who agreed to participate in this study.

References

1. Evans C, Tweheyo R, McGarry J, et al. Crossing cultural divides: a qualitative systematic review of factors influencing the provision of healthcare related to female genital mutilation from the perspective of health professionals. *PLoS One* 2019; **14**(3): e0211829. DOI: <https://doi.org/10.1371/journal.pone.0211829>
2. World Health Organization. *Eliminating female genital mutilation: an interagency statement* — OHCHR, UNAIDS, UNDP, UNECA, UNESCO, UNFPA, UNHCR, UNICEF, UNIFEM, WHO; http://whqlibdoc.who.int/publications/2008/9789241596442_eng.pdf?ua=1 (accessed 11 Aug 2020).
3. World Health Organization. *Female genital mutilation: an overview*. 1998; https://apps.who.int/iris/bitstream/handle/10665/42042/9241561912_eng.pdf?sequence=1&isAllowed=y (accessed 11 Aug 2020).
4. UNICEF. *Female genital mutilation/cutting: a global concern*; https://data.unicef.org/wp-content/uploads/2016/04/FGMC-2016-brochure_250.pdf (accessed 11 Aug 2020).
5. United Nations Population Fund. *Bending the curve: FGM trends we aim to change*. 2018; https://www.unfpa.org/sites/default/files/resource-pdf/18-053_FGM-Infographic-2018-02-05-1804.pdf (accessed 11 Aug 2020).
6. Kawous R, van den Muijsenbergh METC, Geraci D, et al. The prevalence and risk of female genital mutilation/cutting among migrant women and girls in the Netherlands: an extrapolation method. *PLoS One* 2020; **15**(4): e0230919–16. DOI: <https://doi.org/10.1371/journal.pone.0230919>
7. Williams-Breault BD. Eradicating female genital mutilation/cutting: human rights-based approaches of legislation, education, and community empowerment. *Health Hum Rights* 2018; **20**(2): 223–233.
8. Bijlsma-Schlösser J, Saïd SH, Mohamed MH, et al [Combating female genital mutilation: A policy recommendation] *Bestrijding vrouwelijke genitale verminking — Beleidsadvies (in Dutch)*. Zoetermeer: Raad voor de Volksgezondheid en Zorg; 2005.
9. Ministry of Health Welfare and Sport. [Public health policy, aiming to prevent female genital mutilation] *Preventiebeleid voor de volksgezondheid*. Den Haag (in Dutch). 2005; <https://zoek.officielebekendmakingen.nl/kst-22894-66.pdf> (accessed 11 Aug 2020).
10. Pijpers F, Exterkate M, Jager M. [Female Genital Mutilation. Risk and Safeguarding. Guidance. Youth Health Care (JGZ)] *Standpunt Preventie van Vrouwelijke Genitale Verminking (VGV) door de Jeugdgezondheidszorg (in Dutch)*. Bilthoven: RIVM; 2010.
11. Johnsdotter S., Mestre i Mestre RM *Female genital mutilation in Europe: an analysis of court cases*. Brussels: European Commission; 2015.
12. Pharos. The Dutch chain approach; <https://www.pharos.nl/english/female-genital-mutilation/the-dutch-chain-approach/> (accessed 25 Nov 2020).
13. Nederlandse Vereniging voor Obstetrie & Gynaecologie (NVOG). [Guideline on the management of health complications from female genital mutilation] *Leidraad Medische zorg voor vrouwen en meisjes met vrouwelijke genitale verminking (VGV) (in Dutch)*. Utrecht: Nederlandse Vereniging voor Obstetrie & Gynaecologie (NVOG); 2019.
14. Berg RC, Underland V, Odgaard-Jensen J, et al. Effects of female genital cutting on physical health outcomes: a systematic review and meta-analysis. *BMJ Open* 2014; **4**(11): e006316. DOI: <https://doi.org/10.1136/bmjopen-2014-006316>
15. World Health Organization. *WHO guidelines on the management of health complications from female genital mutilation*. Geneva: WHO; 2016.
16. World Health Organization. Female genital mutilation: key facts. 2019; <https://www.who.int/news-room/fact-sheets/detail/female-genital-mutilation> (accessed 12 Aug 2020).
17. Vloeberghs E, van der Kwaak A, Knipscheer J, van den Muijsenbergh METC. Coping and chronic psychosocial consequences of female genital mutilation in the Netherlands. *Ethn Health* 2012; **17**(6): 677–695. DOI: <https://doi.org/10.1080/13557858.2013.771148>
18. Elnashar A, Abdelhady R. The impact of female genital cutting on health of newly married women. *Int J Gynaecol Obstet* 2007; **97**(3): 238–244. DOI: <https://doi.org/10.1016/j.ijgo.2007.03.008>
19. Applebaum J, Cohen H, Matar M, et al. Symptoms of posttraumatic stress disorder after ritual female genital surgery among Bedouin in Israel: myth or reality? *Prim Care Companion J Clin Psychiatry* 2008; **10**(6): 453–456. DOI: <https://doi.org/10.4088/PCC.v10n0605>
20. Bommers M, Sievers W, Fassmann H. *Migration from the Middle East and North Africa to Europe: Past Developments, Current Status and Future Potentials*. Amsterdam: Amsterdam University Press; 2014.
21. UNICEF. *Female genital mutilation/cutting: what might the future hold?* 2014; https://www.unicef.org/media/files/FGM-C_Report_7_15_Final_LR.pdf (accessed 12 Aug 2020).
22. Chalmers B, Hashi KO, Kowser Omer H. 432 Somali women's birth experiences in Canada after earlier female genital mutilation. *Birth* 2000; **27**(4): 227–234. DOI: <https://doi.org/10.1046/j.1523-536x.2000.00227.x>
23. Relph S, Inamdar R, Singh H, Yoong W. Female genital mutilation/cutting: knowledge, attitude and training of health professionals in inner City London. *Eur J Obstet Gynecol Reprod Biol* 2013; **168**(2): 195–198. DOI: <https://doi.org/10.1016/j.ejogrb.2013.01.004>

24. Zaidi N, Khalil A, Roberts C, Browne M. Knowledge of female genital mutilation among healthcare professionals. *J Obstet Gynaecol* 2007; **27**(2): 161–164. DOI: <https://doi.org/10.1080/01443610601124257>
25. Zenner N, Liao L-M, Richens Y, Creighton SM. Quality of obstetric and midwifery care for pregnant women who have undergone female genital mutilation. *J Obstet Gynaecol* 2013; **33**(5): 459–462. DOI: <https://doi.org/10.3109/01443615.2013.767785>
26. Huibers M, van der Windt D, Boeke J. [General practitioners' participation in scientific research] *De deelname van huisartsen aan wetenschappelijk onderzoek* (in Dutch). *Huisarts Wet* 2002; **45**(9): 454–458.
27. Razum O, Zeeb H, Akgün S. How useful is a name-based algorithm in health research among Turkish migrants in Germany? *Trop Med Int Health* 2001; **6**(8): 654–661. DOI: <https://doi.org/10.1046/j.1365-3156.2001.00760.x>
28. Derose KP, Escarce JJ, Lurie N. Immigrants and health care: sources of vulnerability. *Health Aff* 2007; **26**(5): 1258–1268. DOI: <https://doi.org/10.1377/hlthaff.26.5.1258>
29. Abdulcadir J, Dugerdil A, Boulvain M, et al. Missed opportunities for diagnosis of female genital mutilation. *Int J Gynaecol Obstet* 2014; **125**(3): 256–260. DOI: <https://doi.org/10.1016/j.ijgo.2013.11.016>
30. Vangen S, Johansen REB, Sundby J, et al. Qualitative study of perinatal care experiences among Somali women and local health care professionals in Norway. *Eur J Obstet Gynecol Reprod Biol* 2004; **112**(1): 29–35. DOI: [https://doi.org/10.1016/S0301-2115\(03\)00313-0](https://doi.org/10.1016/S0301-2115(03)00313-0)
31. Byrskog U, Olsson P, Essén B, Allvin M-K. Being a bridge: Swedish antenatal care midwives' encounters with Somali-born women and questions of violence; a qualitative study. *BMC Pregnancy Childbirth* 2015; **15**(1): 1–11. DOI: <https://doi.org/10.1186/s12884-015-0429-z>
32. Thierfelder C, Tanner M, Bodiang CMK. Female genital mutilation in the context of migration: experience of African women with the Swiss health care system. *Eur J Public Health* 2005; **15**(1): 86–90. DOI: <https://doi.org/10.1093/eurpub/cki120>
33. Lazar JN, Johnson-Agbakwu CE, Davis OI, Shipp MP-L. Providers' perceptions of challenges in obstetrical care for Somali women. 2013; <http://downloads.hindawi.com/journals/ogi/2013/149640.pdf> (accessed 12 Aug 2020).
34. Exterkate M. *Female genital mutilation in the Netherlands: prevalence, incidence and determinants*. Utrecht: Pharos; 2013.
35. Ogunsiji O. Australian midwives' perspectives on managing obstetric care of women living with female genital circumcision/mutilation. *Health Care Women Int* 2016; **37**(10): 1156–1169. DOI: <https://doi.org/10.1080/07399332.2016.1215462>
36. Ogunsiji O. Female genital mutilation (FGM): Australian midwives' knowledge and attitudes. *Health Care Women Int* 2015; **36**(11): 1179–1193. DOI: <https://doi.org/10.1080/07399332.2014.992521>
37. Dawson AJ, Turkmani S, Varol N, et al. Midwives' experiences of caring for women with female genital mutilation: insights and ways forward for practice in Australia. *Women Birth* 2015; **28**(3): 207–214. DOI: <https://doi.org/10.1016/j.wombi.2015.01.007>
38. Johansen REB. Care for infibulated women giving birth in Norway: an anthropological analysis of health workers' management of a medically and culturally unfamiliar issue. *Med Anthropol Q* 2006; **20**(4): 516–544. DOI: <https://doi.org/10.1525/maq.2006.20.4.516>
39. Johansen REB. Virility, pleasure and female genital mutilation/cutting. A qualitative study of perceptions and experiences of Medicalized defibulation among Somali and Sudanese migrants in Norway. *Reprod Health* 2017; **14**(1): 25. DOI: <https://doi.org/10.1186/s12978-017-0287-4>
40. Widmark C, Levál A, Tishelman C, Ahlberg BM. Obstetric care at the intersection of science and culture: Swedish doctors' perspectives on obstetric care of women who have undergone female genital cutting. *J Obstet Gynaecol* 2010; **30**(6): 553–558. DOI: <https://doi.org/10.3109/01443615.2010.484110>
41. Widmark C, Tishelman C, Ahlberg BM. A study of Swedish midwives' encounters with infibulated African women in Sweden. *Midwifery* 2002; **18**(2): 113–125. DOI: <https://doi.org/10.1054/midw.2002.0307>
42. Burchill J, Pevalin DJ. Demonstrating cultural competence within health-visiting practice: working with refugee and asylum-seeking families. *Divers Equal Health Care* 2014; **11**(2): 151–159. DOI: <https://doi.org/10.21767/2049-5471.100010>
43. Taskforce on the Health Aspects of Violence Against Women and Children. Responding to violence against women and children — the role of the NHS. 2010; https://www.health.org.uk/sites/default/files/RespondingtoViolenceAgainstWomenAndChildrenTheRoleofTheNHS_guide.pdf (accessed 12 Aug 2020).
44. van den Muijsenbergh METC. Migrant mental health care. In: Dawrick C, ed. *Global Primary Mental Health Care: Practical Guidance for Family Doctors*. 1st edn. London: Routledge; 2019. 194.
45. Kumar BN, Diaz E. *Migrant Health: A Primary Care Perspective*. 1st edn. Oxford: CRC Press Taylor & Francis Group; 2019. 306.
46. Wenzel T, Droždek B. *An Uncertain Safety: Integrative Health Care for the 21st Century Refugees*. New York, NY: Springer; 2019.
47. Oosterberg E, van den Muijsenbergh METC [Care for low literate, migrants and social vulnerable groups in General Practice] *Zorg voor laaggeletterden, migranten en sociaal kwetsbaren in de huisartsenpraktijk* (in Dutch). Utrecht: NHG & Pharos; 2016.
48. Leye E, Ysebaert I, Deblonde J, et al. Female genital mutilation: knowledge, attitudes and practices of Flemish gynaecologists. *Eur J Contracept Reprod Health Care* 2008; **13**(2): 182–190. DOI: <https://doi.org/10.1080/13625180701780957>
49. Jäger F, Schulze S, Hohlfeld P. Female genital mutilation in Switzerland: a survey among gynaecologists. *Swiss Med Wkly* 2002; **132**(19-20): 259–264. DOI: <https://doi.org/2002/19/smw-09912>