

# Analysis of isokinetic muscle strength for sports physiotherapy research in Korean ssireum athletes

Ji-WOONG NOH, PT, MS<sup>1)a</sup>, BYOUNG-SUN PARK, PT, MS<sup>1)a</sup>, MEE-YOUNG KIM, PT, PhD<sup>1)</sup>, LIM-KYU LEE, PT, MS<sup>1, 2)</sup>, SEUNG-MIN YANG, PT, MS<sup>1)</sup>, WON-DEOK LEE, PT, MS<sup>1)</sup>, YONG-SUB SHIN, PT, MS<sup>1)</sup>, JU-HYUN KIM, PT, PhD<sup>3)</sup>, JEONG-UK LEE, PT, PhD<sup>4)</sup>, TAEK-YONG KWAK, PhD<sup>5)</sup>, TAE-HYUN LEE, PhD<sup>6)</sup>, JU-YOUNG KIM, PhD<sup>6)</sup>, JAEHONG PARK, PhD<sup>7)</sup>, JUNGHWAN KIM, PT, PhD<sup>8)\*</sup>

<sup>1)</sup> Laboratory of Health Science and Nanophysiotherapy, Department of Physical Therapy, Graduate School, Yongin University, Republic of Korea

<sup>2)</sup> Commercializations Promotion Agency for R&D Outcomes, Republic of Korea

<sup>3)</sup> Department of Physical Therapy, College of Health Welfare, Wonkwang Health Science University, Republic of Korea

<sup>4)</sup> Department of Physical Therapy, College of Health Science, Honam University, Republic of Korea

<sup>5)</sup> Department of Taekwondo Instructor Education, College of Martial Arts, Yongin University, Republic of Korea

<sup>6)</sup> Department of Combative Martial Arts Training, College of Martial Arts, Yongin University, Republic of Korea

<sup>7)</sup> Department of Social Welfare, College of Public Health & Welfare, Yongin University, Republic of Korea

<sup>8)</sup> Departments of Physical Therapy, College of Public Health & Welfare, Yongin University: Yongin 449-714, Republic of Korea

**Abstract.** [Purpose] The purpose of the present study was to elucidate the muscle conditions such as the isokinetic muscle of Korean ssireum athletes. [Subjects and Methods] This study enrolled 25 elite ssireum athletes. We measured body composition and peak torque at an angular speed at 60°/s using an isokinetic muscle strength dynamometer. [Results] The lean body mass of the left upper limb was significantly higher than that of the right upper limb. However, the lean body mass of the left lower limb was significantly lower than that of the right lower limb. The peak torque for left elbow flexion was significantly higher than that for right elbow flexion. Conversely, the peak torque for left elbow extension was significantly lower than that for right elbow extension. Furthermore, the peak torque for the left knee was significantly lower than that for the right knee for both flexion and extension. [Conclusion] The data from this study elucidate in part the muscle conditions of Korean ssireum athletes, which can be used to establish a reference for the scientific study of sports physiotherapy.

**Key words:** Isokinetic muscle strength, Ssireum athletes, Sports physiotherapy

(This article was submitted Jun. 16, 2015, and was accepted Jul. 16, 2015)

## INTRODUCTION

Ssireum is a traditional Korean sport in which two athletes hold on to a satba (a cloth-sash tied around the waist and the thigh of the right leg) and use strength and various skills to throw their opponent to the ground<sup>1)</sup>. The weight classifications depend on the level of ssireum such as gyungjang, sojang, chungjang, yongjang, yongsa, yeoksa, and jangsa<sup>1)</sup>. Many previous studies have suggested that different sports

require athletes to have different physical characteristics according to the way in which the sport is played<sup>2-5)</sup>. For example, judo athletes have a robust skeleton (that is, strong bone density) and well-developed muscles to enable them to withstand and transmit the forces applied during the fight<sup>6)</sup>. Likewise, ssireum athletes have special physical characteristics such as higher weight, muscle mass, and body fat in order to enhance their physical performance<sup>1)</sup>. In recent years, an isokinetic muscle test has begun to be used in sports science for the assessment of muscle performance and injury prevention in athletes<sup>7-9)</sup>. Isokinetic muscle testing can also be used to estimate muscle imbalance in athletes. Athletes may develop significant muscle imbalance that is specific to the style of the sport they play. Muscle strength imbalance is assessed by comparing the strength of the right muscle group with that of the left muscle group, as well as by determining the agonistic and antagonistic balance relationship. Muscle

<sup>a</sup>The first two authors (Noh JW and Park BS) contributed equally to this work. \*Corresponding author. Junghwan Kim (E-mail: junghwankim3@yongin.ac.kr)

©2015 The Society of Physical Therapy Science. Published by IPEC Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.

strength imbalance as it relates to injuries in athletes has been studied, and bilateral imbalance of muscle strength has been suggested as a risk factor for injury in many sports<sup>10-13</sup>. Agonistic and antagonistic balance also influences injury in athletes. Many previous studies about injury in athletes have shown that imbalance in the hamstring-to-quadriceps peak torque ratio (H:Q ratio) is correlated with a greater incidence of lower extremity injury<sup>7, 14, 15</sup>. Overall muscle imbalance can be estimated through not only muscle strength but also through muscle architecture. A previous study suggested that bilateral structural imbalance of the muscles is related to lower extremity injuries in professional basketball players<sup>16</sup>. Muscle structure imbalance can be estimated through the use of a body composition analyzer, which can analyze lean body mass (LBM), fat-free mass (FFM), fat mass (FM), body fat (BF), and lean mass of each limb<sup>4</sup>. Although many studies on muscle imbalance in different types of sports activities have been conducted, research on ssireum athletes has rarely been performed. Therefore, our study provides information regarding the muscle conditions of ssireum athletes for the purpose of informing decisions related to training and rehabilitation.

## SUBJECTS AND METHODS

The present study enrolled 25 elite ssireum athletes with no physical or psychological conditions who provided voluntary written informed consent to participate in this study. The seven weight categories were as follows: gyungjang (up to 75 kg), sojang (up to 80 kg), chungjang (up to 85 kg), yongjang (up to 90 kg), yongsa (up to 95 kg), yeoksa (up to 110 kg), and jangsa (up to 150 kg)<sup>1</sup>. The subjects were asked to complete a questionnaire administered by individual in-depth interviews that took about 30 min per person. The protocol for the study was approved by the Committee of Ethics in Research of the University of Yongin, in accordance with the terms of Resolution 5-1-20, December 2006. In the present study, the modified somatotype method for the analysis of body type was used<sup>1, 5, 17-20</sup>. To confirm body composition, a 4 pole-8 pole contacted electronic method was applied. LBM, FFM, FM, BF, and lean mass of both limbs of participants were measured using a precision body composition analyzer (InBody 720; Biospace, Korea). Measurements were taken with the participants wearing only ssireum pants in a standing position with slight abduction of both arms<sup>1</sup>. To measure the isokinetic peak torque of the elbow and knee flexion/extension on both sides, we used an isokinetic muscle strength dynamometer (Isomed 2000; Isomed, Germany). For elbow joint testing, participants were positioned sitting on the dynamometer chair and stabilized using straps across the chest and pelvis. The shoulder joint was positioned in 45° abduction with 30° flexion, and the forearm was supine. The mechanical rotation axis of the dynamometer was aligned with the lateral humeral epicondyle and the established range of motion was 20° to 110°. For examination of the knee joint, the participants were positioned sitting on the dynamometer chair and stabilized by straps across the chest and pelvis and femoral region. The mechanical axis of rotation of the dynamometer was aligned to the femoral lateral epicondyle. Resistance was ap-

plied immediately above the ankle joint, and the established range of motion was 20° to 90° of knee flexion. Tests were performed in the reciprocal concentric mode and composed of three repetitions at 60°/s velocity to measure peak torque. Prior to testing, the subjects performed three sub-maximal contractions for familiarization. The examiner verbally encouraged the subjects during the test to help bring forth maximal efforts. Statistical analyses were conducted using PASW Statistics version 18.0 (SPSS/IBM; Armonk, NY, USA) to calculate averages and standard deviations. The measurements are expressed as mean ± standard error (SE). The significance level was set to  $\alpha=0.05$  when performing a paired t-test.

## RESULTS

Table 1 shows the general characteristics and somatotypes of the Korean ssireum athletes. The LBM of the left upper limb was significantly higher than that of the right upper limb. However, the LBM of the left lower limb was significantly lower than that of the right lower limb (Table 2). The peak torque for left elbow flexion was significantly higher than that for right elbow flexion. Conversely, the peak torque for left elbow extension was significantly lower than that for right elbow extension. Furthermore, the peak torque for the left knee was significantly lower than that for the right knee for both flexion and extension (Table 3). The peak torque/body weight (PT/BW) ratio for both, elbows and knees showed a similar pattern to the respective peak torques (Table 3). However, the H:Q ratio was not significantly different between the knee joints (Table 3).

## DISCUSSION

In the present study, body composition results showed that the lean mass of the upper limbs was higher on the left side than on the right; for the lower limbs, lean mass was higher on the right side. Bilateral muscle peak torque in ssireum athletes also showed an imbalance. Unexpectedly, peak torque and the PT/BW ratio for elbow extension was higher on the left side than on the right. Many previous studies have suggested that muscle strength would be higher on the dominant side than on the non-dominant side in athletes irrespective of whether the upper or lower limbs were evaluated<sup>7, 10-12, 21, 22</sup>. According to previous studies, Brazilian national handball players have significantly stronger throwing-side shoulders than non-dominant-side shoulders, and elite karate athletes have greater elbow strength on the dominant side<sup>21, 22</sup>. However, even though almost all participants in our study were right-handed, body lean mass and peak flexor torque were higher for the left upper limb. This can probably be explained by the nature of ssireum. During a ssireum match, athletes face their right shoulders; they grab the "close waist's" satba with the right hand and the far leg's satba with the left hand. Using this position, the athlete pulls the opponent into his body. This technique of ssireum may result in superior left elbow compared to right elbow flexor strength, possibly because the left hands are farther from what they are reaching for during this pulling motion. The risk of bilateral muscle imbalance is greater

**Table 1.** General characteristics of the Korean ssireum athletes included in the study

Variable	Mean $\pm$ standard error/n (%)
Age (yrs)	21.6 $\pm$ 0.7
Gender	
Male (%) / Female (%)	25 (100.0) / - (-)
Height (cm) / Weight (kg)	178.0 $\pm$ 1.0 / 98.1 $\pm$ 4.1
BMI (kg/m <sup>2</sup> )	30.5 $\pm$ 1.1
Weight division	
Gyungjang	5 (20.0)
Sojang	3 (12.0)
Chungjang	4 (16.0)
Yongjang	3 (12.0)
Yongsa	3 (12.0)
Yeoksa	3 (12.0)
Jangsa	4 (16.0)
Career (yr)	10.4 $\pm$ 0.5
Dominant side	
Upper limb	
Right / Left	24 (96.0) / 1 (4.0)
Lower limb	
Right / Left	24 (96.0) / 1 (4.0)
Body composition	
LBM (kg) / FFM (kg)	40.6 $\pm$ 2.6 / 74.3 $\pm$ 2.2
FM (kg) / BF (%)	21.6 $\pm$ 3.8 / 20.7 $\pm$ 2.3
Somatotype	
Endomorphy	-
Mesomorphy	23 (92.0)
Ectomorphy	2 (8.0)
Central	-
Endo. C	4.7 $\pm$ 0.4
Meso. C	6.4 $\pm$ 0.3
Ecto. C	0.7 $\pm$ 0.1

BMI: body mass index; LBM: lean body mass; FFM: fat-free mass; FM: fat mass; BF: body fat; Endo. C: endomorphic component; Meso. C: mesomorphic component; Ecto. C: ectomorphic component

in the lower limbs than in the upper limbs<sup>7, 10–12</sup>). Bilateral muscle imbalance of the lower limbs was suggested as a risk factor for hamstring and anterior cruciate ligament injuries. A previous study suggested that there is an increased rate of injury in cases of a difference of 15% or more in knee flexor or hip extensor strength in collegiate athletes<sup>14</sup>). In addition, an imbalance in the H:Q ratio was shown to correlate with a higher incidence of lower extremity injury. The typical H:Q ratio of a healthy knee ranges from 50 to 80%; it is commonly accepted that an H:Q ratio of 60% or higher is desirable in rehabilitation<sup>8, 15, 23</sup>). In this study, ssireum athletes experienced no issues with respect to H:Q ratios in their knee joints. However, the bilateral muscle strength of the lower limbs in ssireum athletes was imbalanced; thus, ssireum athletes have a potential risk of knee injuries. According to our previous study on injuries in combat sports, almost 50% of ssireum athletes have experienced a knee

**Table 2.** Differences in the lean mass of the limbs on both sides in the Korean ssireum athletes

Variable	Limb side	
	Right side	Left side
Upper limb (kg)	4.4 $\pm$ 0.2	4.6 $\pm$ 0.2**
Lower limb (kg)	11.0 $\pm$ 0.3	10.9 $\pm$ 0.2**

All data are presented as the mean  $\pm$  standard error. \*\*p < 0.01.

**Table 3.** Difference in the isokinetic muscle strength of the limbs on both sides in the Korean ssireum athletes

Variable		Limb side	
		Right side	Left side
Elbow	PT-Flex (Nm)	68.4 $\pm$ 3.0	73.7 $\pm$ 3.5*
	PT-Ext (Nm)	67.2 $\pm$ 2.8	58.0 $\pm$ 3.0*
	PT/BW-Flex (%)	74.6 $\pm$ 3.6	78.7 $\pm$ 4.2
	PT/BW-Ext (%)	71.8 $\pm$ 3.4	62.3 $\pm$ 3.4*
Knee	PT-Flex (Nm)	139.2 $\pm$ 7.4	126.2 $\pm$ 6.7**
	PT-Ext (Nm)	224.7 $\pm$ 13.7	205.7 $\pm$ 10.5*
	PT/BW-Flex (%)	150.0 $\pm$ 7.4	133.4 $\pm$ 5.7**
	PT/BW-Ext (%)	236.4 $\pm$ 11.8	211.7 $\pm$ 7.4*
H:Q ratio (%)		63.4 $\pm$ 2.1	62.0 $\pm$ 1.9

All data are presented as the mean  $\pm$  standard error. \*p < 0.05, \*\*p < 0.01. PT-Flex: peak torque of flexion; PT-Ext: peak torque of extension; PT/BW-Flex: peak torque/body weight of flexion; PT/BW-Ext: peak torque/body weight of extension; H:Q ratio: hamstring-quadriceps ratio

injury. Although other factors are definitely contributory, this result may be evidence of muscle strength imbalance leading to knee joint injuries in ssireum athletes<sup>24</sup>). Balance training exercises, which may involve the use of resistance or unstable surfaces, can decrease lower limb asymmetry with the aim of reducing the risk of injury<sup>25</sup>). We suggest that ssireum athletes undertake balance training exercise to reduce the risk of injury to their knees. In conclusion, this study revealed a bilateral imbalance in muscle condition in ssireum athletes, suggesting that more studies are needed to assist in proper training, injury prevention, and rehabilitation.

## REFERENCES

- 1) Noh JW, Kim JH, Kim J: Somatotype analysis of Korean wrestling athletes compared with non-athletes for sports health sciences. *Toxicol Environ Health Sci*, 2013a, 5: 163–168. [CrossRef]
- 2) Hattori K, Kondo M, Abe T, et al.: Hierarchical differences in body composition of professional Sumo wrestlers. *Ann Hum Biol*, 1999, 26: 179–184. [Medline] [CrossRef]
- 3) Yang SM, Lee WD, Kim JH, et al.: Differences in body components and electrical characteristics between youth soccer players and non-athletes. *Health*, 2013, 5: 1010–1015. [CrossRef]
- 4) Lee WD, Lee LK, Kim B, et al.: Differences in body components and the significance of rehabilitation for taekwondo athletes compared to nonathletes. *Toxicol Environ Health Sci*, 2012, 4: 203–208. [CrossRef]
- 5) Noh JW, Kim JH, Kim MY, et al.: Somatotype analysis of elite boxing athletes compared with nonathletes for sports physiotherapy. *J Phys Ther Sci*, 2014, 26: 1231–1235. [Medline] [CrossRef]

- 6) Lewandowska J, Buško K, Pastuszek A, et al.: Somatotype variables related to muscle torque and power in judoists. *J Hum Kinet*, 2011, 30: 21–28. [[Medline](#)] [[CrossRef](#)]
- 7) Terreri AS, Ambrósio MA, Pedrinelli A, et al.: Isokinetic assessment of the flexor-extensor balance of the knee in athletes with total rupture of the anterior cruciate ligament. *Rev Hosp Clin Fac Med Sao Paulo*, 1999, 54: 35–38. [[Medline](#)] [[CrossRef](#)]
- 8) Kim JH, Kim IH, Lee JU, et al.: Change of muscular activity and dynamic stability of the knee joint due to excessive and repetitive jumping or cutting by female athletes. *J Phys Ther Sci*, 2012, 24: 715–719. [[CrossRef](#)]
- 9) Kanehisa H, Kondo M, Ikegawa S, et al.: Body composition and isokinetic strength of professional Sumo wrestlers. *Eur J Appl Physiol Occup Physiol*, 1998, 77: 352–359. [[Medline](#)] [[CrossRef](#)]
- 10) Drid P, Ostojic SM, Vujkov S, et al.: Physiological adaptations of a specific muscle-imbalance reduction training programme in the elite female judokas. *Arch Budo*, 2011, 7: 61–64.
- 11) Lockie RG, Schultz AB, Jeffriess MD, et al.: The relationship between bilateral differences of knee flexor and extensor isokinetic strength and multi-directional speed. *Isokinet Exerc Sci*, 2012, 20: 211–219.
- 12) Jones PA, Bampouras TM: A comparison of isokinetic and functional methods of assessing bilateral strength imbalance. *J Strength Cond Res*, 2010, 24: 1553–1558. [[Medline](#)] [[CrossRef](#)]
- 13) Ellenbecker TS, Pluim B, Vivier S, et al.: Common injuries in tennis players: exercises to address muscular imbalances and reduce injury risk. *Strength Condit J*, 2009, 31: 50–58. [[CrossRef](#)]
- 14) Cheung RT, Smith AW, Wong P: H:q ratios and bilateral leg strength in college field and court sports players. *J Hum Kinet*, 2012, 33: 63–71. [[Medline](#)] [[CrossRef](#)]
- 15) Evangelidis PE, Pain MT, Folland J: Angle-specific hamstring-to-quadriceps ratio: a comparison of football players and recreationally active males. *J Sports Sci*, 2015, 33: 309–319. [[Medline](#)] [[CrossRef](#)]
- 16) Mangine GT, Hoffman JR, Gonzalez AM, et al.: Bilateral differences in muscle architecture and increased rate of injury in national basketball association players. *J Athl Train*, 2014, 49: 794–799. [[Medline](#)] [[CrossRef](#)]
- 17) Noh JW, Kim JH, Kim J: Somatotype analysis of elite taekwondo athletes compared to non-athletes for sports health sciences. *Toxicol Environ Health Sci*, 2013b, 5: 189–196. [[CrossRef](#)]
- 18) Noh JW, Kim MY, Lee LK, et al.: Somatotype and body composition analysis of Korean youth soccer players according to playing position for sports physiotherapy research. *J Phys Ther Sci*, 2015a, 27: 1013–1017. [[Medline](#)] [[CrossRef](#)]
- 19) Noh JW, Kim JH, Kim J: Somatotype analysis of freestyle wrestlers compared with nonathletes for health science research. *Toxicol Environ Health Sci*, 2014b, 6: 244–250. [[CrossRef](#)]
- 20) Noh JW, Kim JH, Kim J: Somatotype analysis of elite judo athletes compared with nonathletes for health science research. *Toxicol Environ Health Sci*, 2014c, 6: 99–105. [[CrossRef](#)]
- 21) Andrade MS, Vancini RL, de Lira CA, et al.: Shoulder isokinetic profile of male handball players of the Brazilian National Team. *Braz J Phys Ther*, 2013, 17: 572–578. [[Medline](#)] [[CrossRef](#)]
- 22) Scattoni-Silva R, Lessi GC, Lobato DF, et al.: Acceleration time, peak torque and time to peak torque in elite karate athletes. *Sci Sports*, 2012, 27: e31–e37. [[CrossRef](#)]
- 23) Andrade MS, De Lira CA, Koffes FC, et al.: Isokinetic hamstrings-to-quadriceps peak torque ratio: the influence of sport modality, gender, and angular velocity. *J Sports Sci*, 2012, 30: 547–553. [[Medline](#)] [[CrossRef](#)]
- 24) Noh JW, Park BS, Kim MY, et al.: Analysis of combat sports players' injuries according to playing style for sports physiotherapy research. *J Phys Ther Sci*, 2015, 27: 2425–2430. [[CrossRef](#)]
- 25) Sannicandro I, Cofano G, Rosa RA, et al.: Balance training exercises decrease lower-limb strength asymmetry in young tennis players. *J Sports Sci Med*, 2014, 13: 397–402. [[Medline](#)]