



POSTER PRESENTATION

Open Access

Vitamin D deficiency rickets – tip of the ice burg

Navoda Atapattu^{1*}, Subashini Jayasena¹, Eresha Jasinghe¹, Dayanath Bolonnage², Shamy De Silva³

From 8th APPEs Biennial Scientific Meeting
Darwin, Australia. 29 October – 1 November 2014

Introduction

Vitamin D deficiency rickets is an increasingly recognized condition across the world in both developed and developing countries. There is no vitamin D fortification program in Sri Lanka. The main source of Vitamin D is sun exposure. The burden of vitamin D deficiency rickets was not studied Sri Lanka.

Aims

Aims of this study were to identify the number of children affected with Vitamin D deficiency rickets in a single center, affected age groups and the level of vitamin D at which the nutritional rickets appears

Method

The study was conducted in the largest children's hospital in Sri Lanka. The children with radiological rickets were included in the study. Patients with chronic renal failure and liver failure were excluded. The study was conducted from 2012 June to 2014 June. Data obtained from the department of chemical Pathology Lady Ridgeway Hospital Sri Lanka. Vitamin D level was measured using Chemiluminescence method.

Results

Total number of patients presented with rickets was 42. There were 36 children with vitamin D deficiency rickets (F=21). Median age of presentation was 22 months. 22 patients were below 2 years of age (6%). All patients came with complaints of bowing of legs except a 5 month old baby who presented with hypocalcaemic convulsions. The median vitamin D level was 27.9 nmol/L (range 11.5-38.3). There were 18 (50%) children with vitamin D level less than the sample median of 27.9 nmol/L.

Conclusion

Vitamin D deficiency rickets is seen in Sri Lankan children despite the presence of sunshine throughout the year. 61% of the children were below 2 years. Lack of sun exposure due to socio economic changes happened in the recent past would have contributed to vitamin D deficiency rickets. Further community based larger studies are needed to identify the prevalence of vitamin D deficiency and factors contributing to vitamin D deficiency rickets in Sri Lanka.

Authors' details

¹Lady Ridgeway Hospital, Colombo, Sri Lanka. ²Teaching Hospital Karapitiya, Galle, Sri Lanka. ³Faculty of Medicine, University of Colombo, Colombo, Sri Lanka.

Published: 28 April 2015

doi:10.1186/1687-9856-2015-S1-P64

Cite this article as: Atapattu et al.: Vitamin D deficiency rickets – tip of the ice burg. *International Journal of Pediatric Endocrinology* 2015 2015 (Suppl 1):P64.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Lady Ridgeway Hospital, Colombo, Sri Lanka
Full list of author information is available at the end of the article