



Stimulated brains and meditative minds: A systematic review on combining low intensity transcranial electrical stimulation and meditation in humans



Rebecca Divarco^{a,1,*}, Perianen Ramasawmy^{a,1}, Frank Petzke^b, Andrea Antal^a

^a Department of Neurology, University Medical Center Göttingen, Georg-August University, Robert-Koch-Straße 40, Göttingen 37075, Germany

^b Pain Clinic, Department of Anesthesiology, University Medical Center Göttingen, Georg-August University, Göttingen, Germany

ARTICLE INFO

Keywords:

Transcranial electrical stimulation (tES)
Transcranial direct current stimulation (tDCS)
Meditation
Yoga
Brain stimulation

ABSTRACT

Background: Low intensity transcranial electrical stimulation (tES) and meditation are two promising, yet variable, non-pharmacological interventions. Growing research is investigating combined effects of both techniques on one's cognitive, emotional, and physical health.

Objective: This article reviews the current research that combines tES and meditation interventions in healthy and diseased participants. The review considers the intervention parameters and their effects in a well-organized manner.

Method: A systematic search for clinical and experimental published studies was conducted in the PubMed, Cochrane, and transcranial direct current stimulation (tDCS) databases using common keywords for tES and for meditation techniques well defined by previous studies. Unpublished ongoing studies were identified with the ClinicalTrials.gov and DRKS.de clinical trial websites.

Results: 20 published studies and 13 ongoing studies were included for qualitative analysis. 13 published articles studied patients with chronic pain, psychological disorders, cognitive impairment, and movement disorders. Anodal tDCS was the only tES technique while mindfulness meditation was the most common meditation type. Eight studies had a main group effect, with outcome improvement in the active combined intervention. However, most published studies showed improvements after at least one combined intervention with variable effects.

Conclusion: Pairing anodal tDCS with meditation shows promising improvements of the physical, mental, and emotional aspects of daily life. Further studies are required to confirm the relevance of this combination in the clinic.

Introduction

The prevalence for using non-pharmacological interventions in the treatment of diseases is high. This is strongly related to the potential that these interventions have on treating non-curable diseases and conditions in a less invasive and more suitable way (Herguedas, 2021; Lefaucheur et al., 2017; Terranova et al., 2019). Two such interventions are meditation and non-invasive brain stimulation.

Meditation finds its roots in most spiritual traditions of the world, especially Hinduism and Buddhism (Matko & Sedlmeier, 2019). Meditation is defined as a mind-body practice, which comprises at least one of many intentional awareness activities, such as observation, focus, release, production, imagination, and movement, which are reinforced and unified by meta-awareness in an informal or formal setting (Sparby & Sacchet, 2022). Meditation practice might lead to altered states of consciousness, scaling through different stages of development and finally end goals of the practice such as 'awakening' (Nash et al., 2013;

Reddy & Roy, 2019). These meditation experiences, stages, and states may be explained within spiritual or secular frameworks (Sparby & Sacchet, 2022). A systematic review by Ospina et al. (2007) classifies the different types of meditation practices to five broad categories: 1) mindfulness meditation (Tang et al., 2015), 2) Yoga (Varambally & Gangadhar, 2016), 3) Tai Chi (Wayne & Kaptchuk, 2008), 4) mantra meditation (Lynch et al., 2018) and 5) Qi Gong (Feng et al., 2020). Two further commonly used meditation techniques are loving kindness (Hofmann et al., 2011) and compassion meditation, and non-dual meditation (Dahl et al., 2015; Brandmeyer et al., 2019).

In a secular world, meditation offers multiple possibilities to improve symptoms, reduce pain and develop better disease coping and management strategies ranging from psychiatric disorders (Wielgosz et al., 2019) to chronic pain Hilton et al. (2016); Zeidan and Vago (2016) and even eye diseases, such as glaucoma (Dada et al., 2019). The benefits of meditation have been shown in different age groups of patients, ranging from children and adolescents to older adults (Nanthakumar, 2018;

* Corresponding author.

E-mail address: rebecca.divarco@stud.uni-goettingen.de (R. Divarco).

¹ Shared first authorship

Perry-Parrish et al., 2016; Simkin & Black, 2014). Over the decades, meditation, especially mindfulness meditation (MM), has become a very popular method to reduce stress, develop resilience, and promote health and well-being (Huberty et al., 2019; Lemay et al., 2019; Zollars et al., 2019). Many studies implement a sham meditation group as a control group. The most commonly studied and used sham protocol for mindfulness meditation involves guided breathing exercises without any guidance on how to regulate breath or body awareness to control for relaxation and expectation effects. The body position, duration and eye state kept the same as experimental condition (Zeidan et al., 2010; Ahn et al., 2019; Noone & Hogan, 2018). In addition, sham yoga includes basic muscle stretching exercises (Sabina et al., 2005) while sham Qi Gong involves series of mild body movements without meditation and deep breathing guidance (Moon et al., 2020).

The exact mechanisms underlying the practice of meditation is still elusive, with different meditation types having different psychophysiological effects on the practitioner (Brandmeyer et al., 2019). It has been suggested that contemplative practices might improve well-being by modulating maladaptive self-referential thought patterns (Dahl et al., 2015). The practice of meditation including mindfulness may further lead to emotional regulation by bolstering prefrontal cognitive control enhanced amygdala top-down control (Bandmeyer et al., 2019). With most of the contemplative research focusing on mindfulness meditation, it has been proposed that this meditation type exerts its effects via improved self-regulation through enhanced self-awareness, attentional control and emotional regulation (Tang et al., 2015).

On the other hand, low intensity transcranial electric stimulation (tES) techniques are neurophysiological techniques used in both clinical practice and research for safe, effective, and non-invasive brain stimulation in humans (Nitsche & Paulus, 2011; Paulus, 2011). tES intends to directly alter brain function through passing weak currents (<4 mA) through at least two electrodes placed on the scalp (Antal et al., 2017). Two commonly used tES methods include transcranial direct current stimulation (tDCS) and transcranial alternating current stimulation (tACS), which includes transcranial random noise stimulation (tRNS). tDCS is the most frequently used method, it comprises the application of a weak direct current to the scalp (1–2 mA) delivered via at least two electrodes (Brunoni et al., 2011). This leads to a subthreshold change in the neuronal resting membrane potential by modulating the probability with which a neuron fires an action potential in response to input from another neuron (Antal & Herrmann, 2016; Bindman et al., 1964; Creutzfeldt et al., 1962). tDCS can lead to changes in brain activity at both local and network levels (Opitz et al., 2016; Paulus et al., 2012). Repeated tDCS can lead to long lasting changes in cortical excitability through long-term potentiation-like or depression-like mechanisms, which underlie associated behavioral effects (Stagg et al., 2009, 2018). The net effects of the tDCS depends on multiple factors such as polarity, intensity, and duration of the stimulation, the state of the brain, and what the participant is doing during the session, among others (Fregni et al., 2020). Clinical application of repeated tDCS reveals potential therapeutic effects in epilepsy, chronic pain, unipolar depression and stroke (Fregni et al., 2020; Lefaucheur et al., 2017). Furthermore, healthy elderly people demonstrated improved cognitive performance such as different types of memory and learning following a single or consecutive repeated tDCS sessions (Prehn & Flöel, 2015; Siegert et al., 2021). Meanwhile, tACS, including tRNS, is effective at modulating the amplitude, frequency and phase of brain oscillations which leads to behavioral effects (Antal et al., 2022). Recent tACS research has shown potential therapeutic effects on cognitive outcomes in psychiatric and neurodegenerative conditions. (Al Qasem et al., 2022).

Over the years, there has been a growing interest in the development and application of novel and low-cost non-pharmacological therapies to prevent and treat diseases in the clinic or to improve quality of life in non-patient groups. Both meditation and tES have individually shown their ability to relieve symptoms, predominantly in the short term but also long-term. They are also attractive for the low cost and the

possibility of home-based intervention when proper training is provided (Herguedas, 2021; Terranova et al., 2019). However, there are also limitations for each strategy. For example, it can be difficult to control whether someone is properly meditating and to see its effects without significant training (Badran et al., 2017). Meanwhile, tES can vary drastically depending on the stimulation parameters, such as intensity, duration, montage, and number of sessions performed, which the latter influences the effects after the stimulation (Bikson et al., 2019). For both techniques, the subject's current emotional and physical state can also influence the efficacy of the intervention (Fregni et al., 2020).

It has been recently hypothesized that these techniques may provide greater feasibility and efficacy when combined, either simultaneously or subsequently. Currently, there are no systematic reviews for such a topic available. Therefore, this review aims to systematically organize, analyze, and compare the overall effects of studies that implement combined tES and meditation interventions in both patients and healthy individuals, with consideration of the parameters of each technique and their influence on each other. In order to assess the effectiveness of tDCS protocols in combination with meditation, we consider outcomes in healthy individuals and patients separately in the results sections. We also consider clinical and experimental studies irrespective of indication for tES and meditation. We aim to provide an overview of successes and drawbacks of combining these two non-pharmacological methods, either in a concurrent or consequent manner in research. This review also illustrates the variability in both meditation parameters (type, duration) and tDCS parameters (duration, intensity, stimulation site) and their indications for use. Moreover, we also summarize the ongoing clinical studies in this review to demonstrate the current scientific interest in combining tES and meditation in clinical populations, with respect to disease condition, type of tDCS protocol and meditation type.

Methods

Databases

In September 2022, a literature database search for published studies involving both tES and meditation techniques was conducted in the PubMed, tDCS database, Cochrane Library databases. PubMed was primarily used due to its easy navigation and high number of available studies to collect a large amount published studies (U.S. National Library of Medicine n.d.). Meanwhile, the Cochrane Library and tDCS Database were used to find additional studies that were not included in the initial PubMed searches (John Wiley & Sons, Inc, n.d.; TDCS database, n.d.). On the other hand, the tDCS Database (Grossman et al., 2018) was used to search additional studies that were not present in the previous searches that involved tDCS specifically. In addition to published studies, a search for ongoing clinical studies was conducted in Cochrane, ClinicalTrials.gov (NIH U.S. National Library of Medicine, n.d.), and the Deutsches Register Klinischer Studien (DRKS, n.d.; German Clinical Trials Register) databases. The same keywords were used for all databases.

Keyword search

The database search was first done in PubMed, then Cochrane, tDCS database, ClinicalTrials.gov and DRKS. For all appropriate keyword searches, see [Supplementary Materials Table 1](#). All abstracts were screened for relevance. For the ongoing studies that were listed as completed, but without study data posted, the registration code was searched on Google to determine if the study was published. The following terms were used for the tES intervention: "transcranial electrical stimulation", "transcranial direct current stimulation", "tDCS", "transcranial alternating current stimulation", "tACS", "transcranial random noise stimulation", and "trns". Meanwhile, the following terms were used for meditation techniques: "meditation", "mindfulness", "mindful", "mantra", "yoga", "taichi/tai chi", "qigong/qi gong", "mindfulness based stress reduction", "breathing exercises", "non dual",

Table 1

Study design, population, intervention characteristics, and database(s) of which it was found of the selected published studies. For the participants, the listed items are in the following order: *N* = # total subjects analyzed, # of female/male, age [mean ± SD], condition, *n* = # of subjects in the active and combined group. [Witkiewitz et al. \(2019\)](#), [Brown et al. \(2020\)](#), and [Gibson et al. \(2022\)](#) all used the same participants for different experiments. In addition, [Ahn et al. \(2019\)](#) and [Pollonini et al. \(2020\)](#) also used the same participants with different analyses. The * on “mean ± SD” number represents those studies that did not include an overall mean ± SD for all subjects combined, and instead was calculated by taking the averages of the means and SD across the included groups. The ** represents the total number of subjects in both active and combined groups. The *** represents the subjects that were dropped from the previous analysis. The intervention column uses “Arm” for cross over studies and “Group” for parallel studies. If not at the same time, the timing of the meditation + tES technique is stated in the parentheses following the condition. Those that have **** mean that the intervention order and type was not clearly stated. The following acronyms are used: tDCS = transcranial direct current stimulation, MM = mindfulness meditation, LKM = loving kindness meditation, MBCT = mindfulness based cognitive training, MBRP = mindfulness-based relapse prevention, MBSAT = mindfulness-based substance abuse treatment, MBSR = mindfulness based stress reduction. For sham meditation techniques type, see [Table 3 and 4](#), and for definitions, see Supplementary Materials Table 2.

Study	Study design	Participants	Dropouts	Intervention	Database (s)
Badran et al. (2017)	Randomized, sham-controlled, double-blind, cross over, clinical, pilot	<i>N</i> = 15, 7/8, 28.2 ± 6.2, healthy, <i>n</i> = 15	Not stated	Arm 1: real MM & active tDCS (1 mA); Arm 2: real MM & active tDCS (2 mA); Arm 3: real MM & sham tDCS;	PubMed Cochrane tDCS Database ClinicalTrials.gov
Hunter et al. (2018)	Randomized, double-blind, parallel	<i>N</i> = 29, 11/18, 27.5 ± 5.45*, healthy, <i>n</i> = 16	4	Group 1: real MM & active tDCS; Group 2: sham MM & sham tDCS;	PubMed Cochrane
Ahn et al. (2019)	Randomized, sham-controlled, double-blind, single center, parallel, clinical, pilot	<i>N</i> = 30, 18/12, 59.47 ± 6.91, knee osteoarthritis in older adults, <i>n</i> = 15	8	Group 1: real MM & active home-based tDCS; Group 2: sham MM & sham home-based tDCS;	PubMed Cochrane ClinicalTrials.gov
Hung (2019)	Randomized, sham-controlled, parallel, clinical, pilot	<i>N</i> = 26, 18/8, 47.85 ± 10.51, treatment resistant unipolar depression, <i>n</i> = 8	Not stated	Group 1: mindfulness training**** & active tDCS; Group 2: active tDCS only; Group 3: sham tDCS only;	PubMed Cochrane
Monnart et al. (2019)	Randomized, controlled, parallel, clinical, pilot	<i>N</i> = 31, 20/11, 50.16 ± 6.73*, treatment-resistant depressed patients, <i>n</i> = 15	15	Group 1: real MBCT (2nd) & active tDCS (1st); Group 2: sham MBCT (2nd) & active tDCS (1st);	PubMed Cochrane
Robinson et al. (2019)	Randomized, sham-controlled, double-blind, parallel, pilot	<i>N</i> = 87, 63/24, 20.16 ± 4.34, healthy college students, <i>n</i> = 26**	4	Group 1: real LKM & active tDCS (2 mA, F3); Group 2: real LKM & active tDCS (2 mA, CP6); Group 3: real LKM & sham tDCS (0.1 mA, F3); Group 4: real LKM & sham tDCS (0.1 mA, CP6); Group 5: sham LKM & active tDCS (2 mA, F3); Group 6: sham LKM & active tDCS (2 mA, CP6); Group 7: sham LKM & sham tDCS (0.1 mA, F3); Group 8: sham LKM & sham tDCS (0.1 mA, CP6);	PubMed Cochrane tDCS Database
Witkiewitz et al. (2019)	Randomized, sham-controlled, double-blind, parallel, clinical	<i>N</i> = 84, 34/50, 52.27 ± 13.00, older adults with alcohol use disorder, <i>n</i> = 47	Not stated	Group 1: real MBRP & active tDCS; Group 2: real MBRP & sham tDCS;	PubMed Cochrane
Clarke et al. (2020)	Randomized, sham-controlled, single blind, parallel	<i>N</i> = 97, 75/22, 22.12 ± 5.51*, healthy young adults, <i>n</i> = 25	2	Group 1: real MM & active tDCS; Group 2: real MM & sham tDCS; Group 3: sham MM & active tDCS; Group 4: sham MM & sham tDCS;	PubMed Cochrane
Brown et al. (2020)	Randomized, sham-controlled, double-blind, parallel, clinical, secondary analysis of Witkiewitz et al. (2019)	<i>N</i> = 68, 32/36, 52.16 ± 13.6, older adults with alcohol use disorder, <i>n</i> = 36	16	Group 1: real MBRP & active tDCS; Group 2: real MBRP & sham tDCS;	PubMed Cochrane
Pollonini et al. (2020)	Randomized, sham-controlled, double-blind, clinical, pilot study, secondary analysis of Ahn et al. (2019)	<i>N</i> = 19, 10/9, 60 ± 7.7, knee osteoarthritis in older adults, <i>n</i> = 11	11***	Group 1: real MM & active tDCS; Group 2: sham MM & sham tDCS;	PubMed Cochrane

(continued)

Table 1 (Continued)

Study	Study design	Participants	Dropouts	Intervention	Database (s)
Alizadehgoradel et al. (2021)	Randomized, sham-controlled, single-blind, parallel, clinical	$N = 64, 0/64, 19.46 \pm 1.15$, youths with methamphetamine addictions, $n = 17$	16	Group 1: real MBSAT (2nd) & active tDCS (1st); Group 2: active tDCS only; Group 3: real MBSAT only; Group 4: sham tDCS only;	PubMed Cochrane
Brooks et al. (2021)	Randomized, sham-controlled, double-blind, two site parallel, clinical, pilot study	$N = 26, 12/14, 68.65 \pm 5.45^*$, cognitive complaints and symptoms of depression/anxiety in older adults, $n = 12$	9	Group 1: real MBSR & active tDCS; Group 2: real MBSR & sham tDCS;	PubMed Cochrane ClinicalTrials.gov
Danilewitz et al. (2021)	Randomized, sham-controlled, double-blind, cross over	$N = 22, 14/8, 29.05 \pm 4.56$, healthy, $n = 22$	2	Arm 1: real yoga (2nd) & active tDCS (1st); Arm 2: real yoga (2nd) & sham tDCS (1st);	PubMed Cochrane
Liao et al. (2021)	Randomized, sham-controlled, double-blind, parallel, clinical	$N = 20, 13/7, 72.85 \pm 4.35^*$, older adults with mild cognitive impairments, $n = 10$	1	Group 1: real tai chi & active tDCS; Group 2: real tai chi & sham tDCS;	PubMed Cochrane
Nishida et al. (2021)	Randomized, sham-controlled, double-blind, parallel, clinical	$N = 56, 34/22, 39.80 \pm 10.06$, healthy to subclinical, $n = 27$	2	Group 1: real TW-FM (MM) & active tDCS; Group 2: real TW-FM (MM) & sham tDCS;	PubMed Cochrane
Park et al. (2021)	Randomized, sham-controlled, double-blind, cross over	$N = 5, 2/3, 52 \pm 4$, clinically definite functional movement disorder, $n = 5$	Not stated	Arm 1: real yoga (2nd) & active tDCS (1st); Arm 2: real yoga (2nd) & sham tDCS (1st);	PubMed Cochrane
Pimenta et al. (2021)	Randomized, sham-controlled, double-blind, parallel, clinical, pilot	$N = 24, 24/0, 32.91 \pm 9.96^*$, females with chronic migraine, $n = 12$	7	Group 1: real MM & active tDCS; Group 2: real MM & sham tDCS;	PubMed Cochrane
Gibson et al. (2022)	Randomized, sham-controlled, double-blind, parallel, clinical, secondary analysis of Witkiewitz et al. (2019)	$N = 84, 34/50, 52.27 \pm 13.00$, older adults with alcohol use disorder, $n = 47$	Not stated	Group 1: real MBRP & active tDCS; Group 2: real MBRP & sham tDCS;	PubMed Cochrane ClinicalTrials.gov
Ramasawmy et al. (2022)	Randomized, sham-controlled, placebo-controlled, double-blind, parallel, clinical, pilot	$N = 30, 28/2, 53.60 \pm 9.48$, adults with fibromyalgia syndrome, $n = 10$	6	Group 1: real MM & active tDCS; Group 2: real MM & sham tDCS;	Cochrane DRKS.de
Sefat et al. (2022)	Randomized, sham-controlled, double-blind, cross over	$N = 18, 5/13, 27.04 \pm 4.51$, healthy, $n = 18$	4	Group 3: no intervention; Arm 1: real yoga (1st) & active tDCS (2nd); Arm 2: real yoga (1st) & sham tDCS (2nd);	PubMed Cochrane

“transcendental” and “loving kindness”. These meditation keywords were chosen based on the five broad categories of meditation defined by Ospina et al. (2007), and the various meditation techniques listed by Brandmeyer et al. (2019). Supplementary Materials Table 1 describes the search process and the search results for each appropriate database.

Eligibility criteria

The process of identifying, screening, and assessing for eligibility of studies is summarized in a PRISMA flow diagram (Fig. 1). Studies were included if they met the following criteria: 1) included at least one arm of intervention where there was combination of tES (tDCS, tACS, or tRNS) and one of the following meditation types, such as mindfulness meditation, mantra meditation, yoga, tai chi, qigong, non-dual meditation, transcendental meditation, loving kindness meditation or meditation involving breathing exercises, 2) randomized design, and 3) human participants, part of either healthy or clinical groups.

Studies were excluded if they: 1) contained transcranial magnetic stimulation (TMS), transcutaneous electrical stimulation, cranial electrical stimulation, neuro- or bio-feedback, theory of mind therapy, physical therapy, behavioral therapy, electroacupuncture, and/or motor/guided imagery as an intervention, 2) were designed as a review or meta-

analysis, 3) were not completed or accessible (i.e. abstracts only, poster abstract, no abstract or complete paper available, or not able to be translated) and/or 4) were designed as a case study. The blinding status of the studies was not an exclusion factor.

Analysis

Studies were first given priority of review if they had healthy participants or patients. Then, the studies were selected for analysis if they included an arm of intervention where tES and a form of meditation were applied at the same time or in a sequential manner. Finally, these studies were analyzed and discussed based on the efficacy of the combination of the two techniques.

Results

Study design

Out of the 276 studies searched, 20 studies published between 2017 and 2022 (Fig. 2A) and 13 ongoing studies met all inclusion criteria. All the included studies, whether published or ongoing, included tDCS as the tES technique and no studies were or are being performed with tACS or tRNS (Table 1 & 2). Out of the published articles, there were only 17




Fig. 1. PRISMA flow diagram depicting identification, screening (databases and reports), and inclusion strategies for the selection of the published and ongoing studies. * Includes duplicate records that were returned for multiple keyword searches. ** Includes ongoing and published studies. The PRISMA diagram was modified from Page et al. (2021).

unique studies as Brown et al. (2020) and Gibson et al. (2022) were secondary analyses for Witkiewitz et al. (2019), and Pollonini et al. (2020) was the secondary analysis of the Ahn et al. (2019) study. The published studies included 664 participants, after correcting for secondary analyses publications using the same participants, comprising of 56.93% females and 43.07% males (Table 1). Seven out of the 20 studies were conducted in healthy individuals (Badran et al., 2017; Clarke et al., 2020; Danilewitz et al., 2021; Hunter et al., 2018; Nishida et al., 2021; Robinson et al., 2019; Sefat et al., 2022), while the remaining studies tested the combination of tDCS and meditation in patient groups diagnosed with conditions such as chronic pain (Ahn et al., 2019; Pimenta et al., 2021; Pollonini et al., 2020; Ramasawmy et al., 2022), psychological disorders (Alizadehgoradel et al., 2021; Brooks et al., 2021; Brown et al., 2020; Gibson et al., 2022; Hung, 2019; Monnart et al., 2019; Witkiewitz et al., 2019), mild cognitive impairment (Liao et al., 2021), and movement disorders (Park et al., 2021). All the ongoing studies test the combined intervention in diseased populations (Table 2). For a list of the published and ongoing studies and their respective or anticipated study details (design, population/patient information, & intervention) and source information (title, authors, & year published/expected), see Tables 1 and 2, respectively.

Out of the seven published studies with healthy participants, one study (Clarke et al., 2020) had a double-blinded study design, where both the participants and the experimenters were blinded to the intervention (Table 1). In the patient groups, ten studies were double-blinded, while Alizadehgoradel et al. (2021) was single blinded

and Monnart et al. (2019) and Hung (2019) were not blinded (Table 1, Fig. 2B). In addition, four studies had a cross over design, with all participants receiving both sham and real tDCS protocols (Badran et al., 2017; Danilewitz et al., 2021; Park et al., 2021; Sefat et al., 2022), while the remaining studies followed a parallel design (Table 1).

Meditation parameters

The most common type of meditation used mindfulness techniques (Mindfulness Meditation (MM)), accounting for 15 out of the 20 published studies (75%) and 12 out of the 13 ongoing studies (92%). Mindfulness was used in this review as an umbrella term for any intervention that included some aspect of MM. In this review, MM was used to describe an audio-guided mindfulness meditation while sitting ($n = 6$) or walking ($n = 1$). Treadmill walking focused meditation (TW-FM) is a type of MM practiced during walking. Some meditative interventions included psychoeducation aspects to it and other tasks, such as mindfulness-based stress reduction training (MBSR; $n = 1$), mindfulness-based relapse prevention (MBRP; $n = 3$), mindfulness-based substance abuse treatment (MBSAT; $n = 1$), mindfulness based cognitive therapy (MBCT; $n = 1$). Hung (2019) was the only study that lacked information on the nature of the mindfulness training. In the published studies, the rest of the studies used yoga ($n = 3$), tai chi ($n = 1$), and loving kindness meditation (LKM; $n = 1$) as meditation interventions (Table 3 & 4). As for the ongoing studies, one study is using a type of MM which is called




Fig. 2. Characteristics of the published studies. A) Number of studies published according to year. B) Hierarchical representation the characteristics of the published studies. The order of significance from the inside-out is the condition of the participants, whether it is double-blinded, and the meditation technique. The outer most layer is the study itself. All studies included tDCS as the tES technique. MM = mindfulness meditation, MBCT = Mindfulness-Based Cognitive Therapy, MBRP = Mindfulness-Based Relapse Prevention, MBSAT = Mindfulness-Based Substance Abuse Treatment, MBSR = Mindfulness-Based Stress Reduction, TW-FM = Treadmill Walking with Focused Meditation. The * represents the studies that have a cross over design. The ** represents the studies where tES and meditation technique are not implemented concurrently.

breathing and attention training (BAT; Table 2). The technique, duration, number of sessions and timepoints in respect to tDCS is listed in Tables 3 & 4, where the tables differ according to their participant population information. The detailed description of the meditation technique used in each published study is summarized in Supplementary Materials Table 2.

tDCS parameters

For the selected publications, the stimulation parameters are listed in Table 3 & 4 according to participant population groups and visualized in Fig. 3. The most common duration of stimulation was 20 min (65%, Fig. 3). The only studies that used 30-min stimulations were done at 2 mA (35%, Fig. 3). The montages of electrode placement for the 20 published studies were very different (Table 3 & 4). The most frequent montage was the left (anode, F3) and right (cathode, F4) dorsolateral prefrontal cortex (dlPFC) (25%, Alizadehgoradel et al., 2021; Danilewitz et al., 2021; Hung, 2019; Monnart et al., 2019; Sefat et al., 2022). Generally, most of the studies (95%) stimulated the frontal lobe with the exception being Park et al. (2021), who stimulated at the right temporoparietal junction (tPJ). It should also be noted that Robinson et al. (2019) also stimulated the tPJ along with the dlPFC. In addition, the overall number of sessions varied greatly across the tDCS parameters.

Review of Selected Groups of Studies

tDCS + meditation in healthy people

Impact on psychological and emotional wellbeing

In 2017, Badran et al. were the first to combine tDCS with meditation, donning the term E-meditation with the goal of facilitating meditation practice in naïve meditators to potentially enhance wellbeing. With respect to the meditation ability of the 15 included participants, those receiving 1 mA anodal tDCS applied at the left ventrolateral prefrontal cortex paired with MM showed higher ability to act with awareness as a measure of mindfulness level, compared to sham and frontal tDCS at 2 mA group. They found that E-meditation with tDCS at both 1 mA and 2 mA showed relevant mathematical improvements in positive affect of the participants (Badran et al., 2017). To further test the potential impact of E-meditation on facilitating mindfulness acquisition, Nishida et al. (2021) investigated brain function using electroencephalography (EEG) and anxiety levels following an intervention of 1 mA anodal tDCS applied at the left dlPFC concurrently paired with TW-FM. Participants in the active group reported one week later significantly lower state anxiety and current density of the alpha band at the rostral anterior cingulate cortex (rACC) which was significantly correlated to the anxiety scores than the sham group (sham tDCS + TW-FM). The authors suggest

Table 2

Title, study design, population, intervention characteristics, and database(s) of which it was found of ongoing studies. For the participants, the listed items are in the following order: condition, age target, and expected enrollment (N). The – represents the ongoing studies that do not have a year of expected completion listed. For the meditation techniques specified, the following acronyms are used: MM = mindfulness meditation, MBCT = mindfulness based cognitive training, MBSR = mindfulness based stress reduction, BAT = Breathing and Attention Training. In addition, tDCS = transcranial direct current stimulation, DLPFC = dorsolateral prefrontal cortex, rTMS = repetitive transcranial magnetic stimulation.

Registration Identifier	Title	Study Design	Participants	Intervention	Primary Outcomes	Expected Year of Completion	Database (s)
IRCT20191214045723N1	The effect and comparison of education on mindfulness-based stress reduction and combination method mindfulness-based stress reduction/ MBSR and transcranial direct current stimulation/tDCS in the severity of stress, quality of life and cognitive functions of migraine sufferers	Randomized, controlled, double blind, parallel, Clinical	Young adults and adults (18–60 yrs) with migraines, <i>N</i> = 45	MBSR with active tDCS, MBSR alone, or neither;	Changes in severity of stress, quality of life and cognitive functions	2019	Cochrane
NCT03897699	Mindful breathing and neuromodulation for depression in young people	Randomized, controlled, quadruple blind, parallel, Clinical	Adolescents and young adults (16–24 yrs) with depression, <i>N</i> = 68	Mindful Breathing Training (MBT) with active or sham tDCS;	Change in DLPFC Connectivity	2021	Cochrane Clinical-Trials.gov
NCT04652869	Mindfulness + tDCS to reduce urgency incontinence in women	Randomized, controlled, open label, parallel, clinical	Adult (40+ yrs) women with urge urinary incontinence, <i>N</i> = 60	MM with active tDCS, MM only, or active tDCS only;	Feasibility (percent enrolled completed) Post-study survey (acceptability and compliance of study) Cue reactivity questionnaire Reaction time to Urinary Stroop Task Questionnaire for evaluating female lower urinary tract symptoms (ICIQ-FLUTS) Urge Urinary Incontinence (UUI) Incidence	2022	Cochrane Clinical-Trials.gov
NCT04012853	Remotely supervised tDCS for persistent post-traumatic headache: A pilot study	Randomized, sham-controlled, quadruple blind, parallel, clinical, pilot	Adults (20–60 yrs) with persistent post traumatic headache associated with mild traumatic brain injury (mTBI), <i>N</i> = 24	MM with active or sham tDCS;	Number of moderate to severe headache days per month	2022	ClinicalTrials.gov
NCT05005013	A telehealth tDCS approach to decrease cannabis use: Towards reducing multiple sclerosis disability	Randomized, controlled, double blind, parallel, clinical	Adults (21–65 yrs) with cannabis use disorder, <i>N</i> = 46	MM with active or sham tDCS;	Changes in Kessler Psychological Distress Scale (K10) Positive and Negative Affect Schedule (PANAS-SF) Marijuana Craving Questionnaire (MCQ-17) Cannabis Withdrawal Scale (CWS) scores Number of Daily Sessions and Frequency of Cannabis Use	2024	ClinicalTrials.gov
NCT03708159	Maintenance of response after rTMS for depression using tDCS (START)	Randomized, controlled, triple blind, multi-site, parallel, clinical	Young adults and adults (18–85 yrs) with major depressive disorder with recent acute rTMS treatments, <i>N</i> = 200	MM with active or sham tDCS;	Depressive symptoms or response to rTMS treatments as measured by the Hamilton Rating Scale for Depression	2024	ClinicalTrials.gov
NCT05460676	Reducing tobacco smoking: a transcranial direct current stimulation (TDCS) telehealth study	Randomized, sham-controlled, double blind, parallel, clinical	Adults (21–75 yrs) who smoke cigarettes and are distressed, <i>N</i> = 46	MM with active or sham tDCS;	Global measure of distress from the Kessler Psychological Distress Scale and weekly cigarette use	2024	ClinicalTrials.gov

(continued on next page)

Table 2 (Continued)

Registration Identifier	Title	Study Design	Participants	Intervention	Primary Outcomes	Expected Year of Completion	Database (s)
NCT04375072	Combination therapy of home-based transcranial direct current stimulation and mindfulness-based meditation for self-management of clinical pain and symptoms in older adults with knee osteoarthritis	Randomized, controlled, triple blind, parallel, clinical	Knee osteoarthritis in older adults (50–85 yrs), <i>N</i> = 200	MM with active or sham tDCS;	Change in clinical pain as assessed by the numeric rating scale (NRS) for pain	2025	Cochrane Clinical-Trials.gov
NCT03884374	Pain relief for osteoarthritis through combined treatment (PROACT)	Randomized, sham-controlled, double-blind, cross over, clinical	Knee osteoarthritis in non-Hispanic black and white adults (45–85 yrs), <i>N</i> = 240	BAT mindfulness technique (MM) with active or sham tDCS and sham BAT with active or sham tDCS;	Change in Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)	2027	PubMed Cochrane Clinical-Trials.gov
DRKS00029024	Delving in the behavioral and neural correlates of pain perception using a therapy combining mindfulness meditation and transcranial direct current stimulation for the treatment of fibromyalgia syndrome	Randomized, sham-controlled, double blind, clinical	Fibromyalgia syndrome in adults (30–70 yrs), <i>N</i> = 80	MM with active or sham tDCS;	Change in clinical pain as assessed by the numeric rating scale (NRS) for pain and quality of life with the Fibromyalgia Impact Questionnaire	–	Cochrane DRKS.de
ChiCTR1800015629	Clinical research on influence of taijiquan exercise and transcranial direct current stimulation on cognitive function of patients with mild cognitive impairment	Randomized, sham controlled, no blinding specified, parallel, clinical	Mild cognitive impairment in older adults (50–75 yrs), <i>N</i> = 156	Tai Chi with active or sham tDCS, or walking with active or sham tDCS;	Montreal Cognitive Assessment scale (MoCA)	–	Cochrane
IRCT20090716002195N3	Efficacy of mindfulness-based intervention and transcranial direct current stimulation (tDCS) on cognitive disorders, and emotional problems in patients with stroke: A randomized clinical trial	Randomized, Controlled, double blind, parallel, clinical	Adults (30–70 yrs) with stroke, <i>N</i> = 72	MBSR with active tDCS, MBSR only, active tDCS only, or neither;	Addenbrooke's Cognitive Examinations and Beck's depression and anxiety inventories	–	Cochrane
IRCT20201004048920N1	Effectiveness of mindfulness-based cognitive therapy separately and combined with tDCS on worry, emotion regulation, pain severity and quality of life in fibromyalgia patients	Randomized, controlled, no blinding, parallel, clinical	Adults (30–50 yrs) with Fibromyalgia, <i>N</i> = 48	Mindfulness-Based Cognitive Therapy (MBCT) with active tDCS, MBCT only, active tDCS only, or neither;	Cognitive emotion regulation, Short-form McGill Pain, Short-Form 36 (SF-36, quality of life), and Pennsylvania State Worry Questionnaires	–	Cochrane

that the immediate reduction in rACC alpha activity might modulate the long-term activity of the amygdala, hence reducing anxiety level (Nishida et al., 2021).

Robinson et al. (2019) used a more robust study design with 8 arms to test the effects of concurrent anodal tDCS-LKM intervention on positive affect and emotional intelligence, varying the electrode placement (left dlPFC; right tPJ), current intensity (2 mA-active; 0.1 mA-sham) and meditation type (real LKM, control). Participants who received anodal tDCS at the tPJ with real LKM had a significantly higher positive affect to neutral and positive images after stimulation regardless of current strength than the other electrode position. However, they did not find any main effect of stimulation current strength, electrode placement, or LKM group on emotional intelligence or positive affect (Robinson et al., 2019). More recently, Clarke et al. (2020) investigated the combination of anodal tDCS over the left dlPFC and a guided body scan, a type of MM, on anxious responses to induced worry. They found significantly higher anxiety reactivity to intentional worry in participants receiving active tDCS regardless of active or sham MM condition, which they further argue that the enhanced emotional reactivity is rather due to the interaction between tDCS and networks involved in negative cognitive patterns such as worry (Clarke et al., 2020).

Impact on cognitive function

Hunter et al. (2018) were the first to test the combination of MM and anodal tDCS at 2 mA over the right inferior frontal gyrus on working memory (WM) and attentional resource allocation using EEG. They found a group specific improvement in WM performance for the active group, but not for the sham group (real MM and sham tDCS). The observed larger P3 amplitude and redistribution of frontal and posterior theta power during the N-back WM task is in line with the neural efficiency hypothesis, which proposes a more efficient brain functioning with higher cognitive capacity level (Neubaue & Fink, 2009).

Along the same line of interest, Danilewitz et al. (2021) and Sefat et al. (2022) paired yoga with tDCS targeting WM. Danilewitz et al. (2021) found that one session of yoga following anodal 2 mA (left dlPFC) tDCS failed to show any improvement on WM or mindfulness level compared to a sham session of yoga combined with sham tDCS in a cross-over design. However, a placebo effect was observed with higher WM performances after first session than second one. Sefat et al. (2022) had a similar intervention design to Danilewitz et al. (2021), but rather, participants performed yoga prior to stimulation. Despite finding no significant differences in power spectral density, they found that anodal tDCS over the left dlPFC at 2 mA and yoga had significantly increased functional connectivity in the frontal lobe, as measured using EEG. They argued that this increased connectivity could lead to changes in executive and attentional functions.

tDCS + meditation in diseased patients

Chronic pain

Combining meditation and tDCS opens a novel avenue in chronic pain management through their potential additive effects. Ahn et al. (2019) showed that patients suffering from knee osteoarthritis reported significant improvements of clinical pain and quantitative sensory assessments than the sham group (sham tDCS + sham MM) after 10 sessions of MM concurrently paired with 2 mA anodal tDCS over the primary motor cortex of the hemisphere contralateral to the affected knee. Pollonini et al. (2020) analyzed a participant subset from Ahn et al. (2019) to test the neuronal responses due to heat evoked pain using functional near-infrared spectroscopy. They showed significant reduction in pain intensity only in the active treatment group, but not in the sham group. Following active treatment, there was an increase in oxyhemoglobin

activation in the superior motor and somatosensory cortices next to the anode (Pollonini et al., 2020). Recently, we tested the efficacy of 10 sessions with concurrent MM and 2 mA anodal tDCS over the left primary motor cortex in patients with fibromyalgia syndrome who received a brief 5-day training in MM (Ramasawmy et al., 2022). We found a significant improvement in the quality of life in patients in the active groups compared to the sham group (sham tDCS + real MM) and to a waiting-list control group, without any improvement in pain intensity, pain sensitivity, sleep and psychological symptoms.

Lastly, in a study on migraine headache, Pimenta et al. (2021) found that 4 weeks (3 days per week) of therapy combining MM and 2 mA anodal tDCS for 20 min over the left dlPFC concurrently in patients with chronic migraine improved results of both mindfulness and clinical headache pain assessments. Both the sham group (sham tDCS + real MM) and the active group benefitted from the therapy, but the effect sizes were larger in the active group than sham one.

Psychological disorders

Four published studies investigated the effects of combined intervention on substance abuse disorders. However, a double-blinded study testing the combined effects of MBRP with tDCS in patients with alcohol use disorders led to three separate investigations of the data (Brown et al., 2020; Gibson et al., 2022; Witkiewitz et al., 2019). In this experiment, participants received either 8 sessions of 2 mA anodal tDCS over the right inferior frontal gyrus or sham tDCS concurrently with MBRP training. Witkiewitz et al. (2019) demonstrated a significant decrease in alcohol consumption and craving across all groups post intervention, with effects lasting up to two months. However, there was no specific group effect observed, yet the more MBRP sessions participants took part in, the greater decrease in alcohol consumption occurred (Witkiewitz et al., 2019). Despite the lack of combined effects of tDCS and MBRP, the mindfulness intervention alone led to decreases in craving in response to alcohol cues was also supported by electrophysiological measurements (Brown et al., 2020). An interesting observation by Gibson et al. (2022) was that participants in the sham group with more mindfulness intervention sessions attended reported larger decreases in craving than the active tDCS group. However, no interaction effect for self-reported mindfulness was reported. Contrastingly, in a methamphetamine abuse study, Alizadehgoradel et al. (2021) evaluated the impact of combining anodal tDCS over the left dlPFC with MBSAT on executive function in young adults with methamphetamine addictions. They found that the combination of tDCS and MBSAT significantly improved most of the executive tasks and craving after intervention, while only the executive tasks were still improved one month later (Alizadehgoradel et al., 2021).

On the other hand, three studies were interested in effects of the combination therapy on depressive and anxiety disorders. Hung (2019) tested the therapeutic effects of combining 2 mA anodal tDCS over the left dlPFC and mindfulness training in patients with treatment resistant depression as compared to active tDCS and sham tDCS monotherapies. Hung (2019) found that the first e-mindfulness session after two weeks of tDCS training significantly decreased the depression ratings in all three groups, and the effect persisted over the remaining 6 weekly mindfulness trainings (including 2 e-mindfulness sessions). In a similar study with treatment resistant depression (Monnart et al., 2019), patients also received anodal tDCS over the left dlPFC but prior to undergoing MBCT or Jacobsen relaxation (sham MBCT). Nevertheless, their results show a significant improvement of anxiety, depression and cognitive symptoms post intervention compared to baseline in both groups, regardless of MBCT or relaxation, after 8 consecutive sessions with tDCS. However, those that received MBCT maintained more positive clinical and cognitive symptoms after an additional session 2 weeks after the 8 sessions of treatment (Monnart et al., 2019). Brooks et al. (2021) also examined the

Table 3

Intervention parameters for selected published studies with healthy participants. Meditative characteristics and transcranial direct current stimulation (tDCS) parameters are listed according to each study. The * represents the studies with cross over designs. For the meditation techniques, the following acronyms are used: MM = mindfulness meditation, LKM = loving kindness meditation, TW-FM = treadmill walking with focused meditation. The positions of the electrodes are based on the standard 10–20 EEG system and the other locations are listed as: vIPFC = ventrolateral prefrontal cortex, IFG = inferior frontal gyrus, dlPFC = dorsolateral prefrontal cortex, tPJ = temporoparietal junction. The montage of the tDCS electrode is listed as anode first then cathode location. Concerning the electrode size, wherever the shape of the electrode was stated, it is written in parentheses. In the tDCS intensity column, the current density is inside parentheses. The ** represents the ramp information for the tDCS stimulation was not clearly stated in the study but implied to have occurred due to specification of sham ramping method and therefore, the details of the ramps are elaborated on.

Study	Meditation Characteristics			tDCS Parameters							
	Technique	Duration	No. of sessions	Timepoint	Montage	Size of electrode	tDCS Duration	Intensity	Sham conditions	No. of sessions	Timepoint
Badran et al. (2017)*	MM	20 min	3 (2)	Each condition only once a week separated by a week; Same time as tDCS;	-F8 (right vIPFC); -Left supraorbital;	25 cm2	20 min; No ramp stated;	1 mA or 2 mA	Yes; No intensity stated; No ramp stated;	3 (2)	Each condition only once a week separated by a week; Same time as meditation;
Hunter et al. (2018)	MM	30 min	20 (8)	Once every weekday for four weeks; Same time as tDCS for two of the weekdays; Immediately after baseline EEG;	-F10 (right IFG); -Left upper arm;	11 cm2	30 min; No ramp stated;	2 mA	Yes; 0.1 mA; No ramp stated;	8	Two nonconsecutive days a week for four weeks; Same time as training;
Robinson et al. (2019)	LKM and relaxing podcasts (sham LKM)	30 min	1	Same time as tDCS;	-F3 (left dlPFC) or CP6 (right tPJ); -Contralateral triceps;	25 cm2	30 min No ramp stated	2 mA	Yes; 0.1 mA; No ramp stated;	1	Same time as meditation;
Clarke et al. (2020)	MM and Mind-wandering (sham MM)	14 min	1	Same time as tDCS;	-F3 (left dlPFC); -Left superior trapezius muscle (reference);	30 cm2	20 min; 30 s ramp up at the beginning and 30 s ramp down at the end	2 mA (0.07 mA/cm2)	Yes; 0 and 1.0 mA; 30 s ramp to 1 mA for 1 min at beginning with 30 second ramp down to 0 mA for the remaining 18 min	1	Same time as meditation/mind-wandering;
Danilewitz et al. (2021)*	Traditional Hatha Yoga	40 min	2 (1)	Once a week for two weeks; Immediately after tDCS;	-F3 (left dlPFC); -F4 (right dlPFC);	35 cm2	20 min; 30 s ramp up at the beginning and 30 s ramp down at the end;	2 mA (0.08 mA/cm2)	Yes; 0 and 2 mA; 30 s ramp up to 2 mA and immediate 30 s ramp down to 0 mA for the remaining time;	2 (1)	Each condition only once a week separated by a week; Immediately before Yoga;
Nishida et al. (2021)	MM: TW-FM	20 min	1	Same time as tDCS; Encouraged to engage in walking mindfulness for 1 week;	-F5 (left dlPFC); -Left shoulder;	35 cm2 (circular)	20 min; 30 s ramp up at the beginning and 30 s ramp down at the end;	1 mA	Yes; 0 and 1 mA; Description of sham protocol is not clear;	1	Same time as meditation;
Sefat et al. (2022)*	Hatha yoga	40 min	2 (1)	Once a week for 2 weeks; Immediately before tDCS;	-F3 (dlPFC); -F4 (right dlPFC);	25 cm2	20 min; No ramp stated**;	2 mA (0.08 mA/cm2)	Yes; 0 and 2 mA; 30 s ramp up to 2 mA and then 30 s ramp down to 0 mA for the remaining time;	2 (1)	Each condition only once separated by a week; Immediately after yoga;

Table 4

Intervention parameters for selected published studies with diseased participants. Meditative characteristics and transcranial direct current stimulation (tDCS) parameters are listed according to each study. The * represents the studies with cross over designs. For the meditation techniques, the following acronyms are used: MM = mindfulness meditation, MBCT = mindfulness based cognitive training, MBRP = mindfulness-based relapse prevention, MBSAT = mindfulness-based substance abuse treatment, MBSR = mindfulness based stress reduction. The positions of the electrodes are based on the standard 10–20 EEG system and the other locations are listed as: M1 = primary motor cortex, dlPFC = dorsolateral prefrontal cortex, IFG = inferior frontal gyrus, tPJ = temporoparietal junction. The montage of the tDCS electrode is listed as anode first then cathode location. Concerning the electrode size, wherever the shape of the electrode was stated, it is written in parentheses. In the tDCS intensity column, the current density is inside parentheses. The ** represents the ramp information for the tDCS stimulation was not clearly stated in the study but implied to have occurred due to specification of sham ramping method and therefore, the details of the ramp are elaborated on.

Study	Meditation Characteristics Technique	Duration	No. of sessions	Timepoint	tDCS Parameters						
					Montage	Size of electrode	tDCS Duration	Intensity	Sham conditions	No. of sessions	Timepoint
Ahn et al. (2019)	MM	20 min	10	Once every week-day for two weeks; Same time as tDCS;	-M1 of the hemisphere contralateral to the affected knee; -Supraorbital region ipsilateral to the affected knee;	35 cm ²	20 min; 10 s ramp up at the beginning and 10 s ramp down at the end;	2 mA	Yes; 0, 2, 0 mA; 10 s ramp up to 30 s of 2 mA then 10 s ramp down at beginning only;	10	Once every week-day for two weeks; Same time as meditation;
Hung (2019)	Mindfulness training (lack of information about nature of training)	1 hr	6	One session a week for six weeks; Same time as tDCS (timing not clearly stated);	-F3 (left dlPFC); -F4 (right dlPFC);	Not stated	30 min; No ramp stated**;	2 mA	Yes; 0 and 2 mA; 2 mA current only at beginning and end, duration of current (initial current and ramp not stated);	12 (6)	Ten working days and then one day on the following 4th and 6th week; Same time as mindfulness training (timing not clearly stated);
Monnart et al. (2019)	MBCT and Jacobson Relaxation (sham MBCT)	2 hr (30 min sham)	9	Eight consecutive days with one session two weeks later; Immediately following tDCS;	-F3 (left dlPFC); -F4 (right dlPFC);	35 cm ² (rect-angular)	20 min; 15 s ramp up at the beginning and 15 s ramp down at the end;	2 mA	No sham	9	Eight consecutive days with one session two weeks later; Immediately before MBCT/relaxation;
Witkiewitz et al. (2019)	MBRP	2 hr	8	Once a week for eight weeks; First 30 mins of guided meditation practice at the same time as tDCS and the following 90 min of discussions of mindfulness;	-F10 (right IFG); -Left upper arm;	15 cm ²	30 min; 30 s ramp up at the beginning and 30 s ramp down at the end;	2 mA	Yes; 0 and 2 mA; 30 s ramp up to 2 mA then 30 s ramp down to 0 mA at the beginning and end; 0 mA for 28 min;	8	Once a week for eight weeks; Rolling group sessions; Same time as guided meditation practice;
Brown et al. (2020)	MBRP	2 hr	8	Once a week for eight weeks; First 30 mins of guided meditation practice at the same time as tDCS and the following 90 min of discussions of mindfulness;	-F10 (right IFG); -Left upper arm;	15 cm ²	30 min; 30 s ramp up at the beginning and 30 s ramp down at the end;	2 mA	Yes; 0 and 2 mA; 30 s ramp up to 2 mA then 30 s ramp down to 0 mA at the beginning and end; 0 mA for 28 min;	8	Once a week for eight weeks; Rolling group sessions; Same time as guided meditation practice;
Pollonini et al. (2020)	MM	20 min	10	Once every week-day for two weeks; Same time as tDCS;	-M1 of the hemisphere contralateral to the affected knee; -Supraorbital region contralateral to the affected knee;	35 cm ² (rect-angular)	20 min; 10 s ramp up at the beginning and 10 s ramp down at the end;	2 mA	Yes; 0, 2, 0 mA; 10 s ramp up to 30 s of 2 mA then 10 s ramp down at beginning only;	10	Once every week-day for two weeks; Same time as meditation;
Alizadehgoradel et al. (2021)	MBSAT	50 min	12	Two sessions a week for six weeks;	-F3 (left dlPFC); -F4 (right dlPFC);	Not stated	20 min; No ramp stated**;	1.5 mA	Yes; 0 and 1.5 mA; 30 s ramp up to 1.5 mA for 30 s with a ramp	12	Two sessions a week at least 72-hrs apart for six weeks;

(continued on next page)

Table 4 (Continued)

Study	Meditation Characteristics Technique	tDCS Parameters				No. of sessions	Timepoint
		Duration	Montage	Size of electrode	tDCS Duration		
				Intensity	Sham conditions		Timepoint
Brooks et al. (2021)	MBSR	2.5 hr	-Fz; -Iz;	25 cm ²	30 min; 30 s ramp up at the beginning and 30 s ramp down at the end;	57 (9 including retreat)	Immediately before MBSAT; Once a week for eight weeks; Same time as guided meditation part of MBSR; Asked to complete at home for the remaining 6 days of the week;
Liao et al. (2021)	Yang Style Tai Chi	40 min	-F3 (left dlPFC); -Fp4 (right supraorbital);	35 cm ²	20 min; 10 s ramp up at the beginning; Duration of ramp down not stated;	36	Three times a week for 12 weeks; Same time as Tai Chi for the first 20 min
Park et al. (2021)*	Yoga	1 h	-CP6 (right rPJ); -Left supraorbital;	35 cm ² (rect-angular)	20 min; 8 s ramp up at the beginning and 8 s ramp down at the end;	16 (2)	Two times a week for eight weeks; One hour after tDCS when tDCS was performed;
Pimenta et al. (2021)	MM	At least 20 min**	-F3 (left dlPFC); -Fp2 (right supraorbital);	25 cm ²	20 min; No ramp stated;	28 (12)	Daily for four weeks; Same time as tDCS when tDCS was performed;
Gibson et al. (2022)	MBRP	2 hr	-F10 (right IFG); -Left upper arm;	15 cm ²	30 min; 30 s ramp up at the beginning and 30 s ramp down at the end;	8	Once a week rolling group sessions for eight weeks; First 30 mins of guided meditation practice at the same time as tDCS and the following 90 min of discussions of mindfulness;
Ramasawmy et al. (2022)	MM	20 min (+ 5 min body scan)	-Left primary motor cortex; -Right supraorbital;	16 cm ² (anode; square); 50 cm ² (cathode; Rectangular)	20 min; 15 s ramp up at the beginning and 15 s ramp down at the end;	15 (10)	Once every week day for three weeks, where the first week was for training only and the last 2 weeks at the same time as tDCS;

(2021) and Sefat et al. (2022) had more than one active session. Five showed significant improvements without having any main or group effects (25%: Hung, 2019; Monnart et al., 2019; Pimenta et al., 2021; Robinson et al., 2019; Witkiewitz et al., 2019).

Four other studies did not show a significant effect of intervention but showed numerical improvements in the appropriate measurements or interaction effects (Badran et al., 2017; Brooks et al., 2021; Brown et al., 2020; Park et al., 2021), where two had specific improvements in the active group (active tDCS and meditation technique; Badran et al., 2017; Brown et al., 2020). The other remaining two studies had either improved across all groups with larger effect size in the active group (Brooks et al., 2021), or only showed general therapeutical improvements (Park et al., 2021).

In general, only five studies applied tDCS non-simultaneously with a meditation technique. In Sefat et al. (2022), participants practiced yoga prior to tDCS stimulation, while in the other four studies participants received tDCS stimulation prior to practicing in yoga (Danilewitz et al., 2021; Park et al., 2021), MBCT (Monnart et al., 2019), or MBSAT (Alizadehgoradel et al., 2021). In addition, out of the twenty studies included, only three studies included a sham meditation condition with active tDCS (Clarke et al., 2020; Monnart et al., 2019; Robinson et al., 2019). Meanwhile, three other studies included a sham tDCS with sham MM as the only control group (Ahn et al., 2019; Hunter et al., 2018; Pollonini et al., 2020).

Discussion

Summary of results from selected studies

The above studies show promising results for the combined effects of tDCS and meditation on emotion regulation, psychological wellbeing, working memory, and pain perception. More specifically, all but two studies (Danilewitz et al., 2021: no significant results; Clarke et al., 2020: worsened symptoms) showed general improvements in either mindfulness behavior or primary clinical outcomes after one or more sessions. This aligns with previous studies claiming that tDCS is more effective with more sessions (Stagg et al., 2018). The studies that showed improvements in their outcomes, but did not necessarily have significant effects, typically improved when meditation training and practice was given.

However, the role of combined therapy on the overall improvements is not as clear across the remaining studies. Pimenta et al. (2021) had a larger effect size for the group with active tDCS and MM despite including additional MM training. Robinson et al. (2019) found significant results at a specific location with LKM, regardless of tDCS intensity, which could be explained by the lack of LKM and tDCS sessions. In addition, Gibson et al. (2022), one of the secondary analyses from Witkiewitz et al. (2019), interestingly only showed significant effects in the sham tDCS group and MBSR training group, but also had no interaction effect on mindfulness.

Methodological considerations

In general, studies are inconsistent in terms of participant population, sample size, and study design. For example, most of the studies have small sample sizes for the active condition (mean $n = 19.7$) which diminishes the statistical power. Moreover, the clinical condition of the participants is very inconsistent across all studies, making it difficult to compare findings across the studies. It is known that the effects of meditation depend on the physical and emotional state of the subject especially based on their own personal goals and beliefs (Wolsko et al., 2004). On the other hand, tDCS effects also vary depending on whether the subject has had any caffeine or medicine that day, the mental and physical condition, as well as their age and education (Antal et al., 2008; Terranova et al., 2019). Moreover, if participants are aware of the

conditions they are receiving, it can influence how they feel (placebo effect), and if experimenters are aware of the condition, it could skew their interpretation of the data. Therefore, having at least a double blinded study is crucial, yet 20% of the studies did not have such a design (Alizadehgoradel et al., 2021; Clarke et al., 2020; Hung, 2019; Monnart et al., 2019).

The studies that did not show a significant effect of intervention could be due to having cross over designs (Badran et al., 2017; Park et al., 2021), not having tDCS and meditation technique at the same time (Park et al., 2021), or not having a proper sham mediation technique (Badran et al., 2017; Brooks et al., 2021; Brown et al., 2020; Park et al., 2021). However, more research is necessary to confirm whether the combined techniques is sufficient for a non-invasive treatment. On the other hand, it is possible that Nishida et al. (2021) had such distinct results since participants were encouraged to practice mindful walking at home for 1 week prior to the last measurements and Sefat et al. (2022) had increased functional connectivity but not in power spectral density (EEG) possibly due to having a cross over design and including yoga as its meditation technique, which its effects have not been as thoroughly investigated in combination with tDCS.

Moreover, most of the studies did not control for meditation type, and thus only looked at the impact active tDCS on meditation. In general, meditation control conditions are important to investigate and determine, and thus should also be implemented alone. However, it is also important to have proper tDCS controls as comparing active tDCS combined with real meditation to a condition where sham tDCS is paired with control meditation makes it challenging to determine what is causing the effect. Therefore, monotherapies of both interventions should be included similarly to Hung (2019) and Alizadehgoradel et al. (2021). Lastly, the parameters of the sham tDCS condition was variable and needs to be further investigated so that there is a more universal sham condition in general. These make it challenging to discern whether the effects in the active tDCS and meditation group is due to the combination of effects or just active tDCS altering the brain responses from the meditation.

In addition to the study and participant characteristics, the number of sessions as well as the duration, and parameters of both tES and meditation techniques are essential for interpreting the results. It has been previously shown that repeated tDCS shows better offline effects (Nitsche & Paulus, 2011), which could explain why the studies with one active session does not show very conclusive effects. However, these studies, among some of the others, mostly did not include a proper training for learning meditation. Ramasawmy et al. (2022) was the only study to include MM training prior to the active sessions, while three other studies included training before and during the total experiment (Brooks et al., 2021; Park et al., 2021; Pimenta et al., 2021). Meditation especially can be difficult for those trying to meditate for the first time. On the other hand, the montage, intensity, and duration of tDCS used by these studies could also influence the final effects. In this review, many studies use the dlPFC as the anodal stimulation site, but it has been argued across majority of the studies that tDCS could have effects across the brain in addition to this location and that the site of stimulation is important for online and offline effects (Nitsche & Paulus, 2011). This means that other areas of the brain, when stimulated, may show better results when combined with meditation.

Expanding on the online and offline effects of tDCS, the interaction between the two techniques' role in manipulating the brain could significantly influence the result. It is possible, as mentioned in Gibson et al. (2022), that the active tDCS and meditation group might be not showing an impact in these studies. They argue that mindfulness techniques cause the ongoing neural activity, which tDCS causes changes in, to be dispersed and more dynamic and thus the effects of tDCS are potentially diminished. In addition, the online and offline effects of both techniques

could be cancelling each other out from the brain activity of mindfulness shifting between focused attention and monitoring (Gibson et al., 2022). Overall, it may be beneficial for future studies to implement more modeling and analysis of individual brain currents and circuitry with EEG or fMRI to help discern the precise techniques necessary (Terranova et al., 2019).

Finally, the timing of the tES and the meditation could also influence the impact shown on the measured outcomes. Out of the seven studies with healthy participants, five of them implemented the tDCS and meditation intervention at the exact same time (Badran et al., 2017; Clarke et al., 2020; Hunter et al., 2018; Nishida et al., 2021; Robinson et al., 2019). All but one (Clarke et al., 2020) showed improvements with the combined intervention. Lastly, the two remaining studies with healthy participants used tDCS and yoga one before another and found no combined effect on working memory (Danilewitz et al., 2021) but showed that there were functional connections occurring (Sefat et al., 2022). For the participants with mental or physical conditions, the results were even more variable.

The ongoing studies promote promising studies that investigate more thoroughly the impacts of tDCS and meditation/meditation techniques. Future studies should include larger sample sizes and precision of double blinding and study designs. In addition, investigating the longer lasting effects of the intervention will be very important for understanding the efficacy of these interventions. It is much easier for someone who is undergoing difficulties to do as little as possible to result in long term improvements, especially when it comes to lifestyle practices. Finally, the overall interaction between tDCS and meditation techniques still needs further analysis with proper control groups so that the influence of one on the other could be revealed.

Limitations

This review has a few limitations. Firstly, this review is a traditional, systematic, qualitative clinical review on studies that involve a form of tDCS and meditation techniques. Therefore, there is no quantitative analysis on the included studies regarding their significance and clinical relevance. In addition, the general grouping of meditation was broadly defined and considered studies that included combined therapies of mindfulness and meditation techniques. Finally, some of the studies included were not the complete published papers, but rather manuscripts, letters, or proceedings, making it challenging to accurately display the results.

Conclusion

Here, we investigated and discussed 20 published studies that used a combination of tDCS and a meditation technique with 664 participants comprising of both healthy and diseased participants. Furthermore, the most common combined techniques were anodal tDCS with mindfulness meditation. This study provides the general overview that the combination of tDCS and meditation shows promising yet inconclusive results of improvements on the physical, mental, and normal aspects of daily life. This review suggests that further studies need to improve on the mentioned limitations of study design and experimental stimulation and meditation training limitations in order to produce more reliable results. Combining tES and meditation is important to examine as both are non-invasive techniques that in combination have shown promising, but mostly not significant results of unclear clinical relevance. Their “combined” potential to enhance short- and long-term effects deserves further study.

Acknowledgments

We acknowledge the support by the Open Access Publication Funds of the Göttingen University. In addition, RD & PR was supported by the

MSc/PhD program “Neurosciences” – International Max Planck Research School at the Georg August University Göttingen.

Supplementary materials

Supplementary material associated with this article can be found in the online version at doi:10.1016/j.ijchp.2023.100369.

References

- Ahn, H., Zhong, C., Miao, H., Chaoul, A., Park, L., Yen, I. H., et al. (2019). Efficacy of combining home-based transcranial direct current stimulation with mindfulness-based meditation for pain in older adults with knee osteoarthritis: A randomized controlled pilot study. *Journal of Clinical Neuroscience*, 70, 140–145. doi:10.1016/j.jocn.2019.08.047.
- Alizadehgoradel, J., Imani, S., Nejati, V., Vanderhasselt, M.-A., Molaei, B., Salehinejad, M. A., et al. (2021). Improved executive functions and reduced craving in youths with methamphetamine addiction: Evidence from combined transcranial direct current stimulation with mindfulness treatment. *Clinical Psychopharmacology and Neuroscience*, 19(4), 653–668. doi:10.9758/cpn.2021.19.4.653.
- Al Qasem, W., Abubaker, M., & Kvašňák, E. (2022). Working memory and transcranial-alternating current stimulation—State of the art: Findings, missing, and challenges. *Frontiers in Psychology*, 13. doi:10.3389/fpsyg.2022.822545.
- Antal, A., Boros, K., Poreisz, C., Chaieb, L., Terney, D., & Paulus, W. (2008). Comparatively weak after-effects of transcranial alternating current stimulation (tacs) on cortical excitability in humans. *Brain Stimulation*, 1(2), 97–105. doi:10.1016/j.brs.2007.10.001.
- Antal, A., & Herrmann, C. S. (2016). Transcranial alternating current and random noise stimulation: Possible mechanisms. *Neural Plasticity* 1–12. doi:10.1155/2016/36168072016.
- Antal, A., Alekseichuk, I., Bikson, M., Brockmüller, J., Brunoni, A. R., Chen, R., et al. (2017). Low intensity transcranial electric stimulation: Safety, ethical, legal regulatory and application guidelines. *Clinical Neurophysiology*, 128(9), 1774–1809. doi:10.1016/j.clinph.2017.06.001.
- Antal, A., Luber, B., Brem, A.-K., Bikson, M., Brunoni, A. R., Cohen Kadosh, R., et al. (2022). Non-invasive brain stimulation and neuroenhancement. *Clinical Neurophysiology Practice*, 7, 146–165. doi:10.1016/j.cnp.2022.05.002.
- Badran, B. W., Austelle, C. W., Smith, N. R., Glusman, C. E., Froeliger, B., Garland, E. L., et al. (2017). A double-blind study exploring the use of transcranial direct current stimulation (tDCS) to potentially enhance mindfulness meditation (e-meditation). *Brain Stimulation*, 10(1), 152–154. doi:10.1016/j.brs.2016.09.009.
- Bikson, M., Esmailpour, Z., Adair, D., Kronberg, G., Tyler, W. J., Antal, A., et al. (2019). Transcranial electrical stimulation nomenclature. *Brain Stimulation*, 12(6), 1349–1366. doi:10.1016/j.brs.2019.07.010.
- Bindman, L. J., Lippold, O. C., & Redfearn, J. W. (1964). The action of brief polarizing currents on the cerebral cortex of the rat (1) during current flow and (2) in the production of long-lasting after-effects. *The Journal of Physiology*, 172(3), 369–382. doi:10.1113/jphysiol.1964.sp007425.
- Brandmeyer, T., Delorme, A., & Wahbeh, H. (2019). The neuroscience of meditation: Classification, phenomenology, correlates, and mechanisms. *Progress in Brain Research*, 1–29. doi:10.1016/bs.pbr.2018.10.020.
- Brooks, H., Oughli, H. A., Kamel, L., Subramanian, S., Morgan, G., Blumberger, D. M., et al. (2021). Enhancing cognition in older persons with depression or anxiety with a combination of mindfulness-based stress reduction (MBSR) and transcranial direct current stimulation (tDCS): Results of a pilot randomized clinical trial. *Mindfulness*, 12(12), 3047–3059. doi:10.1007/s12671-021-01764-9.
- Brown, D. R., Jackson, T. C., Claus, E. D., Votaw, V. R., Stein, E. R., Robinson, C. S., et al. (2020). Decreases in the late positive potential to alcohol images among alcohol treatment seekers following mindfulness-based relapse prevention. *Alcohol and Alcoholism*, 55(1), 78–85. doi:10.1093/alcal/agz096.
- Brunoni, A. R., Amadera, J., Berbel, B., Volz, M. S., Rizzerio, B. G., & Fregni, F. (2011). A systematic review on reporting and assessment of adverse effects associated with transcranial direct current stimulation. *International Journal of Neuropsychopharmacology*, 14(8), 1133–1145. doi:10.1017/s1461145710001690.
- Clarke, P. J. F., Sprylian, B. F., Hirsch, C. R., Meeten, F., & Notebaert, L. (2020). TDCS increases anxiety reactivity to intentional worry. *Journal of Psychiatric Research*, 120, 34–39. doi:10.1016/j.jpsychi.2019.10.013.
- Creutzfeldt, O. D., Fromm, G. H., & Kapp, H. (1962). Influence of transcortical D-C currents on cortical neuronal activity. *Experimental Neurology*, 5(6), 436–452. doi:10.1016/0014-4886(62)90056-0.
- Dada, T., Ramesh, P., & Shakrawal, J. (2019). Meditation: A polypill for comprehensive management of glaucoma patients. *Journal of Glaucoma*, 29(2), 133–140. doi:10.1097/ijg.0000000000001406.
- Dahl, C. J., Lutz, A., & Davidson, R. J. (2015). Reconstructing and deconstructing the self: Cognitive mechanisms in meditation practice. *Trends in Cognitive Sciences*, 19(9), 515–523. doi:10.1016/j.tics.2015.07.001.
- Danilewitz, M., Gao, S., Salehinejad, M., Ge, R., Nitsche, M., & Vila-Rodriguez, F. (2021). Effect of combined yoga and transcranial direct current stimulation intervention on working memory and mindfulness. *Journal of Integrative Neuroscience*, 20(2), 367. doi:10.31083/jjin2002036.

- Feng, F., Tuchman, S., Denninger, J. W., Fricchione, G. L., & Yeung, A. (2020). Qigong for the prevention, treatment, and rehabilitation of covid-19 infection in older adults. *The American Journal of Geriatric Psychiatry*, 28(8), 812–819. doi:10.1016/j.jagp.2020.05.012.
- Fregni, F., El-Hagrassy, M. M., Pacheco-Barrios, K., Carvalho, S., Leite, J., Simis, M., et al. (2020). Evidence-based guidelines and secondary meta-analysis for the use of transcranial direct current stimulation in neurological and psychiatric disorders. *International Journal of Neuropsychopharmacology*, 24(4), 256–313. doi:10.1093/ijnp/pyaa051.
- Gibson, B. C., Votaw, V. R., Stein, E. R., Clark, V. P., Claus, E., & Witkiewitz, K. (2022). Transcranial direct current stimulation provides no additional benefit to improvements in self-reported craving following mindfulness-based relapse prevention. *Mindfulness*, 13(1), 92–103. doi:10.1007/s12671-021-01768-5.
- Deutsches Register Klinischer Studien. Federal Institute for Drugs and Medical Devices. <https://drks.de/search/de>.
- Grossman, P., Alekseichuk, I., de Lara, G., Paneri, K., Kunz, P., Turi, Z. et al. (2018). Transcranial direct current stimulation studies open database (tdcs-OD). <https://doi.org/10.1101/369215>
- Herguedas, A. J. U. (2021). Non-pharmacological interventions in preventive, rehabilitative and restorative medicine. *Alternative Medicine - Update*. doi:10.5772/intechopen.94187.
- Hilton, L., Hempel, S., Ewing, B. A., Apaydin, E., Xenakis, L., Newberry, S., et al. (2016). Mindfulness meditation for chronic pain: Systematic review and meta-analysis. *Annals of Behavioral Medicine*, 51(2), 199–213. doi:10.1007/s12160-016-9844-2.
- Hofmann, S. G., Grossman, P., & Hinton, D. E. (2011). Loving-kindness and compassion meditation: Potential for psychological interventions. *Clinical Psychology Review*, 31(7), 1126–1132. doi:10.1016/j.cpr.2011.07.003.
- Huberty, J., Green, J., Glissmann, C., Larkey, L., Puzia, M., & Lee, C. (2019). Efficacy of the mindfulness meditation mobile app “calm” to reduce stress among college students: Randomized controlled trial. *JMIR MHealth and UHealth*, 7(6). doi:10.2196/14273.
- Hung, G. C. L. (2019). Proceedings #3: Effects of combining transcranial direct current stimulation with mindfulness training in patients with treatment-resistant depression: A pilot study. *Brain Stimulation*, 12(2). doi:10.1016/j.brs.2018.12.172.
- Hunter, M.A., Lieberman, G., Coffman, B.A., Trumbo, M.C., Armenta, M.L., Robinson, C.S. H. et al. (2018). Mindfulness-based training with transcranial direct current stimulation modulates neuronal resource allocation in working memory: A randomized pilot study with a nonequivalent control group. *Heliyon*, 4(7). <https://doi.org/10.1016/j.heliyon.2018.e00685>
- John Wiley & Sons, Inc. (n.d.). *About the cochrane library*. Cochrane Library. <https://www.cochranelibrary.com/about/about-cochrane-library>.
- Lefaucheur, J.-P., Antal, A., Ayache, S. S., Benninger, D. H., Brunelin, J., Cogiamanian, F., et al. (2017). Evidence-based guidelines on the therapeutic use of transcranial direct current stimulation (tdcs). *Clinical Neurophysiology*, 128(1), 56–92. doi:10.1016/j.clinph.2016.10.087.
- Lemay, V., Hoolahan, J., & Buchanan, A. (2019). Impact of a yoga and meditation intervention on students' stress and Anxiety Levels. *American Journal of Pharmaceutical Education*, 83(5), 7001. doi:10.5688/ajpe7001.
- Liao, Y. Y., Liu, M. N., Wang, H. C., Walsh, V., & Lau, C. I. (2021). Combining transcranial direct current stimulation with Tai Chi to improve dual-task gait performance in older adults with mild cognitive impairment: A randomized controlled trial. *Frontiers in Aging Neuroscience*, 13. doi:10.3389/fnagi.2021.766649.
- Lynch, J., Prihodova, L., Dunne, P. J., O'Leary, C., Breen, R., Carroll, Á., et al. (2018). Mantra meditation programme for emergency department staff: A qualitative study. *BMJ open*, 8(9). doi:10.1136/bmjopen-2017-020685.
- Matko, K., & Sedlmeier, P. (2019). What is meditation? proposing an empirically derived classification system. *Frontiers in Psychology*, 10. doi:10.3389/fpsyg.2019.02276.
- Moon, S., Sarmiento, C. V. M., Steinbacher, M., Smirnova, I. V., Colgrove, Y., Lai, S.-M., et al. (2020). Can qigong improve non-motor symptoms in people with parkinson's disease - a pilot randomized controlled trial? *Complementary Therapies in Clinical Practice*, 39, 101169. doi:10.1016/j.ctcp.2020.101169.
- Monnart, A., Vanderhasselt, M. A., Schroder, E., Campanella, S., Fontaine, P., & Kornreich, C. (2019). Treatment of resistant depression: A pilot study assessing the efficacy of a tdcs-mindfulness program compared with a tdcs-relaxation program. *Frontiers in Psychiatry*, 10. doi:10.3389/fpsyg.2019.00730.
- Nanthakumar, C. (2018). The benefits of yoga in children. *Journal of Integrative Medicine*, 16(1), 14–19. doi:10.1016/j.joim.2017.12.008.
- Nash, J. D., Newberg, A., & Awasthi, B. (2013). Toward a unifying taxonomy and definition for meditation. *Frontiers in Psychology*, 4. doi:10.3389/fpsyg.2013.00806.
- Neubauer, A. C., & Fink, A. (2009). Intelligence and Neural Efficiency: Measures of brain activation versus measures of functional connectivity in the brain. *Intelligence*, 37(2), 223–229. doi:10.1016/j.intell.2008.10.008.
- NIH U.S. National Library of Medicine. (n.d.). *Find a study*. ClinicalTrials.gov. <https://clinicaltrials.gov/>.
- Nishida, K., Morishima, Y., Pascual-Marqui, R. D., Minami, S., Yamane, T., Michikura, M., et al. (2021). Mindfulness augmentation for anxiety through concurrent use of transcranial direct current stimulation: A randomized double-blind study. *Scientific Reports*, 11(1). doi:10.1038/s41598-021-02177-3.
- Nitsche, M. A., & Paulus, W. (2011). Transcranial direct current stimulation – update 2011. *Restorative Neurology and Neuroscience*, 29(6), 463–492. doi:10.3233/rnm-2011-0618.
- Noone, C., & Hogan, M. J. (2018). A randomised active-controlled trial to examine the effects of an online mindfulness intervention on Executive Control, critical thinking and key thinking dispositions in a university student sample. *BMC Psychology*, 6(1). doi:10.1186/s40359-018-0226-3.
- Opitz, A., Falchier, A., Yan, C. G., Yeagle, E. M., Linn, G. S., Megevang, P., et al. (2016). Spatiotemporal structure of intracranial electric fields induced by transcranial electric stimulation in humans and nonhuman primates. *Scientific Reports*, 6(1), 1–11. doi:10.1038/srep31236.
- Ospina, M. B., Bond, K., Karkhaneh, M., Tjosvold, L., Vandermeer, B., Liang, Y., et al. (2007). Meditation practices for Health: State of the research. *Evidence report/technology assessment*, 155. <https://pubmed.ncbi.nlm.nih.gov/17764203/>.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., et al. (2021). The Prisma 2020 statement: An updated guideline for reporting systematic reviews. *BMJ (Clinical research ed.)*. doi:10.1136/bmj.n71.
- Paulus, W. (2011). Transcranial electrical stimulation (TES – tdcs; trms, tacs) methods. *Neuropsychological Rehabilitation*, 21(5), 602–617. doi:10.1080/09602011.2011.557292.
- Paulus, W., Antal, A., & Nitsche, M. A. (2012). Physiological basis and methodological aspects of transcranial electric stimulation (tDCS, tACS, and tRNS). *A reference book for transcranial brain stimulation* (pp. 93–106). Taylor and Francis Group essay.
- Park, J. E., Hong, J. Y., & Lee, S. Y. (2021). Transcranial direct current stimulation and yoga for functional movement disorders. *The Neurologist*, 26(6), 231–236. doi:10.1097/nrl.0000000000000345.
- Perry-Parrish, C., Copeland-Linder, N., Webb, L., & Sibinga, E.M.S. (2016). Mindfulness-based approaches for children and youth. *Current Problems in Pediatric and Adolescent Health Care*, 46(6), 172–178. <https://doi.org/10.1016/j.cppeds.2015.12.006>
- Pimenta, L. D., de Araújo, E. L., Silva, J. P., Franca, J. J., Brito, P. N., de Holanda, L. J., et al. (2021). Effects of synergism of mindfulness practice associated with transcranial direct-current stimulation in chronic migraine: Pilot, randomized, controlled, double-blind clinical trial. *Frontiers in Human Neuroscience*, 15. doi:10.3389/fnhum.2021.769619.
- Pollonini, L., Montero-Hernandez, S., Park, L., Miao, H., Mathis, K., & Ahn, H. (2020). Functional near-infrared spectroscopy to assess central pain responses in a nonpharmacologic treatment trial of osteoarthritis. *Journal of Neuroimaging*, 30(6), 808–814. doi:10.1111/jon.12782.
- Prehn, K., & Flöel, A. (2015). Potentials and limits to enhance cognitive functions in healthy and pathological aging by tDCS. *Frontiers in Cellular Neuroscience*, 9. doi:10.3389/fncel.2015.00355.
- Ramasawmy, P., Khalid, S., Petzke, F., & Antal, A. (2022). Pain reduction in fibromyalgia syndrome through pairing transcranial direct current stimulation and mindfulness meditation: A randomized, double-blinded, sham-controlled pilot clinical trial. *Frontiers in Medicine*, 9. doi:10.3389/fmed.2022.908133.
- Reddy, J. S., & Roy, S. (2019). Understanding meditation based on the subjective experience and traditional goal: Implications for current meditation research. *Frontiers in Psychology*, 10. doi:10.3389/fpsyg.2019.01827.
- Robinson, C., Armenta, M., Combs, A., Lamphere, M. L., Garza, G. J., Neary, J., et al. (2019). Modulating affective experience and emotional intelligence with loving kindness meditation and transcranial direct current stimulation: A pilot study. *Social Neuroscience*, 14(1), 10–25. doi:10.1080/17470919.2017.1397054.
- Sabina, A. B., Williams, A., Wall, H. K., Bansal, S., Chupp, G., & Katz, D. L. (2005). Yoga intervention for adults with mild-to-moderate asthma: A pilot study. *Annals of Allergy, Asthma & Immunology*, 94(5), 543–548. doi:10.1016/s1081-1206(10)61131-3.
- Siegrist, A., Diedrich, L., & Antal, A. (2021). New methods, Old Brains—A systematic review on the effects of tdcs on the cognition of elderly people. *Frontiers in Human Neuroscience*, 15. doi:10.3389/fnhum.2021.730134.
- Simkin, D. R., & Black, N. B. (2014). Meditation and mindfulness in clinical practice. *Child and Adolescent Psychiatric Clinics of North America*, 23(3), 487–534. doi:10.1016/j.chc.2014.03.002.
- Sefat, O., Salehinejad, M. A., Danilewitz, M., Shalhaf, R., & Vila-Rodriguez, F. (2022). Combined yoga and transcranial direct current stimulation increase functional connectivity and synchronization in the frontal areas. *Brain Topography*, 35(2), 207–218. doi:10.1007/s10548-022-00887-z.
- Sparby, T., & Sacchet, M. D. (2022). Defining meditation: Foundations for an activity-based phenomenological classification system. *Frontiers in Psychology*, 12. doi:10.3389/fpsyg.2021.795077.
- Stagg, C. J., Best, J. G., Stephenson, M. C., O'Shea, J., Wylezinska, M., Kineses, Z. T., et al. (2009). Polarity-sensitive modulation of cortical neurotransmitters by transcranial stimulation. *Journal of Neuroscience*, 29(16), 5202–5206. doi:10.1523/JNEUROSCI.4432-08.2009.
- Stagg, C. J., Antal, A., & Nitsche, M. A. (2018). Physiology of transcranial direct current stimulation. *The Journal of ECT*, 34(3), 144–152. doi:10.1097/ycet.0000000000000510.
- Tang, Y.-Y., Hölzel, B. K., & Posner, M. I. (2015). The neuroscience of mindfulness meditation. *Nature Reviews Neuroscience*, 16(4), 213–225. doi:10.1038/nrn3916.
- Terranova, C., Rizzo, V., Cacciola, A., Chillemi, G., Calamuneri, A., Milardi, D., et al. (2019). Is there a future for non-invasive brain stimulation as a therapeutic tool? *Frontiers in Neurology*, 9. doi:10.3389/fneur.2018.01146.
- TDCS database. tDCS database. (n.d.). Retrieved from <http://tdcsdatabase.com/>.
- U.S. National Library of Medicine. (n.d.). *About - pubmed*. National Center for Biotechnology Information. <https://pubmed.ncbi.nlm.nih.gov/about/>.
- Varambally, S., & Gangadhar, B. N. (2016). Current status of yoga in mental health services. *International Review of Psychiatry*, 28(3), 233–235. doi:10.3109/09540261.2016.1159950.
- Wayne, P. M., & Kaptchuk, T. J. (2008). Challenges inherent to t'ai chi research: Part II—Defining the intervention and Optimal Study Design. *The Journal of Alternative and Complementary Medicine*, 14(2), 191–197. doi:10.1089/acm.2007.7170b.
- Wielgosz, J., Goldberg, S. B., Kral, T. R. A., Dunne, J. D., & Davidson, R. J. (2019). Mindfulness meditation and psychopathology. *Annual Review of Clinical Psychology*, 15(1), 285–316. doi:10.1146/annurev-clinpsy-021815-093423.
- Witkiewitz, K., Stein, E. R., Votaw, V. R., Wilson, A. D., Roos, C. R., Gallegos, S. J., et al. (2019). Mindfulness-based relapse prevention and transcranial

- direct current stimulation to reduce heavy drinking: A double-blind sham-controlled randomized trial. *Alcoholism: Clinical and Experimental Research*. doi:[10.1111/acer.14053](https://doi.org/10.1111/acer.14053).
- Wolsko, P. M., Eisenberg, D. M., Davis, R. B., & Phillips, R. S. (2004). Use of mind-body medical therapies. *Journal of General Internal Medicine*, 19(1), 43–50. doi:[10.1111/j.1525-1497.2004.21019.x](https://doi.org/10.1111/j.1525-1497.2004.21019.x).
- Zeidan, F., Johnson, S. K., Gordon, N. S., & Goolkasian, P. (2010). Effects of brief and sham mindfulness meditation on mood and cardiovascular variables. *The Journal of Alternative and Complementary Medicine*, 16(8), 867–873. doi:[10.1089/acm.2009.0321](https://doi.org/10.1089/acm.2009.0321).
- Zeidan, F., & Vago, D. R. (2016). Mindfulness meditation-based pain relief: A mechanistic account. *Annals of the New York Academy of Sciences*, 1373(1), 114–127. doi:[10.1111/nyas.13153](https://doi.org/10.1111/nyas.13153).
- Zollars, I., Poirier, T. I., & Pailden, J. (2019). Effects of mindfulness meditation on mindfulness, mental well-being, and perceived stress. *Currents in Pharmacy Teaching and Learning*, 11(10), 1022–1028. doi:[10.1016/j.cptl.2019.06.005](https://doi.org/10.1016/j.cptl.2019.06.005).