

Low-level carbon monoxide exposure affects BOLD fMRI response

Caroline Bendell¹, Shakeeb H Moosavi¹  and Mari Herigstad² 

Journal of Cerebral Blood Flow & Metabolism
2020, Vol. 40(11) 2215–2224
© The Author(s) 2019



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/0271678X19887358
journals.sagepub.com/home/jcbfm



Abstract

Blood oxygen level dependent (BOLD) fMRI is a common technique for measuring brain activation that could be affected by low-level carbon monoxide (CO) exposure from, e.g. smoking. This study aimed to probe the vulnerability of BOLD fMRI to CO and determine whether it may constitute a significant neuroimaging confound. Low-level (6 ppm exhaled) CO effects on BOLD response were assessed in 12 healthy never-smokers on two separate experimental days (CO and air control). fMRI tasks were breath-holds (hypercapnia), visual stimulation and fingertapping. BOLD fMRI response was lower during breath holds, visual stimulation and fingertapping in the CO protocol compared to the air control protocol. Behavioural and physiological measures remained unchanged. We conclude that BOLD fMRI might be vulnerable to changes in baseline CO, and suggest exercising caution when imaging populations exposed to elevated CO levels. Further work is required to fully elucidate the impact on CO on fMRI and its underlying mechanisms.

Keywords

Blood oxygen level dependent, carbon monoxide, fMRI, physiological confounds, smoking

Received: 11 March 2019; Revised 5 September 2019; Accepted: 7 October 2019

Introduction

One of the most common methods used to measure brain function in humans is fMRI, of which blood oxygen level dependent (BOLD) fMRI is arguably the most mainstream technique. BOLD fMRI is an indirect measure of brain activation, based on changes in the ratio of oxygenated to deoxygenated blood in the brain, which depends on cerebral metabolic rate (CMRO₂), cerebral blood volume (CBV) and cerebral blood flow (CBF).¹ These factors may be altered as part of the experimental design or as unintended confounds, potentially affecting BOLD response.

Carbon monoxide (CO) is a toxic gas that can act as a cerebral vasodilator.^{2,3} Increases in CBF with elevations in CO have been shown in animal models^{3–7} as well as in humans.^{8,9} Low-level CO exposure is common, through inhalation of cigarette smoke or air pollution. Due to its high affinity for haemoglobin, CO immediately enters and can linger in the bloodstream for several hours after inhalation. Smokers typically have persisting elevated levels of CO bound to haemoglobin in their blood (carboxyhaemoglobin, COHb), which is reflected in higher exhaled levels of CO

(6 ppm in exhaled air or above¹⁰) compared to non-smokers (1–5 ppm). As smoking behaviour is associated with, e.g., socioeconomic status and disease status, elevated COHb may significantly influence neuroimaging results on the group level in certain demographic groups. For example, if low-level CO causes a baseline increase in CBF, this might affect fMRI outcome as the BOLD response for any given task is assessed by comparing task-related signal to baseline signal. An increased baseline could artificially dampen or alter the time course of the observed task-specific BOLD response.^{11–13} To our knowledge, the effect of CO on fMRI signal has not been investigated

¹Biological and Medical Sciences, Oxford Brookes University, Oxford, UK
²Biomolecular Sciences Research Centre, Department of Biosciences and Chemistry, Sheffield Hallam University, Sheffield, UK

Corresponding author:

Mari Herigstad, Department of Biosciences and Chemistry, Sheffield Hallam University, Owen Building 752, City Campus, Sheffield S1 1WB, UK.

Email: m.herigstad@shu.ac.uk

in previous studies, except as part of cigarette smoking, where its impact is confounded by other (vasoactive) tobacco components. The rationale for this study was to test if CO could alter BOLD fMRI response in response to three different fMRI tasks.

One method to test cerebral vascular function is carbon dioxide (CO₂) exposure. Hypercapnia induces a strong CBF increase,¹⁴ and has often been used as a cerebrovascular challenge in fMRI studies^{11,15–17} due to its global and reproducible effect on BOLD response.^{16,18} Raising baseline CBF by CO₂ inhalation can furthermore reduce or cause delays or non-linearity in the vascular responsiveness to subsequent hypercapnia.^{11,12} In this study, we therefore used hypercapnia derived through breath holds as a tool to investigate the effect of CO on BOLD response, as it is a robust, reproducible stimulus, and susceptible to changes in baseline CBF. We hypothesised that low-level CO inhalation would significantly reduce global BOLD response during hypercapnia. To determine whether the effect of CO extended to common fMRI paradigms, we also included a simple visual stimulation and motor task, hypothesising that CO would dampen BOLD response in brain regions associated with these tasks.

Materials and methods

Participants

We recruited 12 (8 F, age 25.3 ± 4.3 years) healthy never-smokers to the study. Never-smokers were chosen to ensure a uniform sample group as smokers typically have varying levels of COHb and may exhibit variation in, e.g., craving. Exclusion criteria were MRI contraindications, smoking history, history of cardiorespiratory or neurological disease, and pregnancy. Female participants were on hormonal contraceptives. All participants gave written, informed consent. The study was approved by Oxford Brookes University Research Ethics Committee (approval number 140840) and carried out in accordance with the Declaration of Helsinki. The sample size was determined by a formal statistical power calculation

(fMRIpower software package, www.fmripower.org¹⁹). At alpha level 0.05, 11 subjects were found to provide at least 80% power to detect an effect. Twelve were recruited to ensure that the study was powered in case of any unforeseen events (e.g. subject dropout).

Protocol

Participants were asked to attend a preliminary laboratory visit. During this visit, medical history and state and trait anxiety inventory (STAI) questionnaires were completed.²⁰ A CO inhalation test was conducted to let the participant familiarise themselves with the breathing system and the CO exposure. Participants were asked to breathe on a custom-made breathing system through a mouthpiece with their nose occluded, and were given time for their breathing to stabilise before commencing the experiment. A full description of the breathing system can be found in the supplement. After stable breathing had been recorded for 5 min, CO was added to the inspired air over 5 min, out of sight of the participant. Following CO administration, five more minutes of stable breathing was recorded. During the experiment, ECG, pulse pressure and saturation were continuously measured. Expired CO was measured before, immediately after, and 10 min after the breathing test (Micro+ Smokelyzer, Intermedical Ltd, Kent). The Smokelyzer kit is suitable for non-invasive, repeated assessments of expired CO in humans, and its output was compared with COHb values (blood samples) prior to the study to ensure correct readings.

MRI scans were conducted on two separate days (Figure 1). Participants were asked to complete the state anxiety part of STAI on arrival and no more than 15 min after the end of the experiment on each day. Whilst in the scanner, participants were asked to undertake the following tasks: breath holds, a visual stimulation task, a motor task and a simple reaction time task. Breath holds were conducted end-expiration, signalled by visual cues and lasting 15 s. Participants were instructed to follow each breath-hold with an expiration (rather than inspiration) to obtain accurate end-tidal PCO₂ values. The visual stimulation was a




Figure 1. Schematic of protocol. fMRI tasks included breath holds (T1), visual stimulation (8 Hz flashing checkerboard, T2) and a (right hand) finger tapping task (T3). Two sets of BOLD scans (each 10 min 6 s) were obtained on each experimental day, separated by a 5 min breathing intervention (air or CO, order randomized and counterbalanced) during which a structural scan was acquired.

flashing checkerboard (8 Hz, lasting 10 s). The motor task was tapping of the right index finger, signalled by visual cues and lasted 15 s. The reaction time task required participants to immediately press a button upon the appearance of a red dot on the screen (24 appearances, random intervals). These tasks were conducted twice, once before the breathing intervention (baseline) and once after (post-intervention). The participants received the gas mixtures in the scanner, and their head was kept in the same position for the intervention and subsequent scan. On one day, the intervention was air, and on the other day, the intervention was CO. Participants were not aware of which intervention would be given on any of the days and the order of the interventions was randomised and balanced. Participants were asked verbally after each protocol if they felt any change in their breathing, and if they could guess which protocol they had undertaken. Training in all fMRI tasks was given by an experimenter prior to the first scan on each day, to ensure that the participant could reliably complete these on their own in the scanner. Expired CO measurements were made before the first scan, immediately after the second scan (~20 min after the breathing intervention) and 10 min after the second scan (~30 min after the breathing intervention). An extended protocol section can be found in the supplement.

MRI data acquisition

Imaging was performed at the University of Oxford Centre for Clinical Magnetic Resonance Research with a Siemens 3 Tesla TIM-Trio scanner, using a 12-channel head coil. Participants were given two fMRI scans (BOLD echo-planar image acquisition, time repetition (TR) = 3000 ms, time echo (TE) = 30 ms, field-of-view = 192 × 192 mm, voxel-size = 3 × 3 × 3 mm, 45 slices) on each day, separated by the intervention period (air or CO). A structural T1-weighted, whole-brain scan (MPRAGE, TR = 2040 ms, TE = 4.7 ms, flip angle = 8°, voxel-size = 1 × 1 × 1 mm) was collected and used for image registration.

Heart rate (HR) and pulse oximetry (SaO₂, multigas monitor, 9500, MR Equipment), ECG, respiration (respiratory bellows around the chest) and end-tidal partial pressures of oxygen (P_{ET}O₂) and CO₂ (P_{ET}CO₂; Datex, Normocap) were continuously measured throughout the scans. ECG data were observed throughout. All physiological data were sampled at 50 Hz and recorded along with scan volume triggers via PowerLab 16/35 using LabChart (ADInstruments).

Data analysis

fMRI data processing was carried out within FSL (Oxford Centre for Functional Magnetic Resonance Imaging of the Brain (FMRIB) Software Library), using FEAT (FMRI Expert Analysis Tool) Version 6.0. The cluster Z threshold was set to 3.1 and a corrected cluster significance threshold to $p = 0.05$.

Prestatistic processing of the data included MCFLIRT motion correction,²¹ spatial smoothing with a full-width-half-maximum Gaussian kernel of 5 mm and high-pass temporal filtering (Gaussian-weighted least-squares straight line fitting, high-pass filter cut-off of 60 s). FSL motion outliers (<https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSLMotionOutliers>) were used to detect and regress out large motion artifacts. Data were modelled using FMRIB's Improved Linear Model (FILM) with local autocorrelation correction.²² Images were registered to the MNI152 standard space using an affine registration between the EPI and T1-weighted scan and a nonlinear registration between the T1-weighted scan and the MNI standard brain.

General linear models (GLMs) with multiple explanatory variables (EVs) incorporating timing values for the different events were designed to describe the data. The haemodynamic response function (HRF) was modelled using a standard gamma waveform. A physiological noise modelling tool was used to regress out the effects of physiological noise.²³ A 6-s haemodynamic delay was assumed and contrast images were used for higher level analyses as appropriate. An end-tidal CO₂ regressor was used to analyse the BOLD response change associated with the breath-hold challenge. This was done by extracting the breath by breath P_{ET}CO₂ data and convolving this with an HRF (e.g. see Bright and Murphy²⁴). This approach models the breath hold challenge response with the recorded P_{ET}CO₂ values and thus makes no assumption about breath hold length. This analysis fits the signal to the P_{ET}CO₂ data and returns statistical maps of significant changes in BOLD response (thresholded zstats). These statistical maps were also converted to %BOLD/mmHg CO₂ in a second, separate analysis of cerebrovascular reactivity (CVR).

A fixed-effects model was used to generate contrast of parameter estimate (COPE) images of the mean signal for all scans as well as the difference between the baseline and post-intervention scans for each participant on each experimental day. The baseline versus post-intervention difference COPE images were calculated to compensate for any variation in baseline between days and account for potential test-retest variability. This was done by forcing random effects variance to zero in FLAME (FMRIB's Local Analysis of Mixed Effects).^{25,26}

Voxelwise statistical analysis was extended to a group level, in a mixed-effects analysis using FLAME²⁶ with automatic outlier de-weighting, and Z statistic images were thresholded using clusters determined by $Z > 3.1$ and a $p < 0.05$ (corrected) cluster significance threshold. Means of COPE images were calculated for all conditions. Group analyses compared COPE images between protocols for each task using a whole-brain approach, and with the following contrasts of interest: $P_{ET}CO_2$ values with breath holds, presentation of flashing checkerboards (visual task) and finger-tapping. An analysis using the older standard of a 2.3 cluster-forming threshold was also conducted, and is included in the Supplement.

STAI questionnaires were scored according to their respective manuals and compared using paired non-parametric t-tests (Mann–Whitney U test). Reaction times were averaged for each participant and compared using Student's t-test (paired). Physiological data were analysed using custom-written MATLAB scripts and

compared using Student's t-tests (paired). Data obtained during the motor task were used for the comparison of end-tidal gases between protocols.

Results

Psychological and physiological data

There were no significant differences between protocols in anxiety scores ($t(11)=0.61$, $p=0.55$) or reaction times ($t(23)=1.1$, $p=0.29$). None of the participants was able to discern between CO and air inhalations, nor did they report any change in breathing. $P_{ET}O_2$ was reduced between baseline and post-intervention scans in both protocols, but no significant difference was found between protocols ($t(11)=-0.58$, $p=0.57$). There was no change in $P_{ET}CO_2$ or HR between scans or protocols. CO values increased significantly in the CO protocol ($p < 0.0001$, Figure 2), but not air ($p=0.10$). COHb values (estimated from exhaled CO) also showed a significant increase in the CO protocol ($p < 0.0001$, Table 2), but not air. The modest rise in COHb (from $1.1\% \pm 0.1$ to $1.5\% \pm 0.2$) highlights the low level of CO used in the study. Participant details are shown in Table 1, and physiological data in Table 2. Extended participant demographics and physiological data, including $P_{ET}CO_2$ and $P_{ET}O_2$ averages and CO_2 traces for all tasks, can be found in the supplement.

fMRI data

For fMRI results, significance denotes thresholded, cluster corrected, signal (cluster-forming threshold of 3.1, $p < 0.05$).²⁷

Breath hold task (Figure 3). The rise in CO_2 with breath holds caused BOLD response change (increase) throughout the grey matter during all scans. Figure 3 shows pre-intervention and post-intervention BOLD response for both air and CO protocols, and group (protocol) contrasts between the pre- versus post-intervention difference maps. These Z score maps




Figure 2. Exhaled CO (ppm). Baseline, post-scan (~20 min after end-inhalation) and 10 min post-scan (~30 min after end-intervention). Individual values plus average and standard deviation (bold line).

Table 1. Participant details and behavioural data.

	Preliminary visit	MRI visit (CO)	MRI visit (air)
Sex (F/M)	8/4	8/4	8/4
Age (years)	25.3 (4.3)	25.3 (4.3)	25.3 (4.3)
BMI (kg/m^2)	23.6 (3.0)	23.6 (3.0)	23.6 (3.0)
Trait anxiety score	35.4 (7.2) [23–47]	N/A	N/A
State anxiety score	31.1 (8.6) [21–55]	27.0 (4.3) [21–35]	28.2 (4.4) [23–37]
RT change (post > pre)	N/A	15.2 (24.3)	10.0 (21.8)

Note: Mean(SD). Range included for STAI scores.
BMI: body mass index; RT: reaction time.

Table 2. Physiological data.

	MRI visit (CO)		MRI visit (Air)	
	Baseline	Post-intervention	Baseline	Post-intervention
P _{ET} CO ₂ (%)	5.5 (0.7)	5.5 (0.7)	5.3 (0.7)	5.4 (0.7)
P _{ET} O ₂ (%)	15.5 (1.2)	15.0 (0.9)*	15.5 (0.7)	14.9 (0.7)**
HR (bpm)	69.3 (12.5)	66.2 (8.3)	71.3 (15.0)	63.5 (8.2)
CO (ppm)	2.9 (1.0)	5.7 (0.7)***	3.0 (0.7)	2.7 (0.8)
COHb (%)	1.1 (0.1)	1.5 (0.2)***	1.1 (0.1)	1.1 (0.1)

Note: Mean(SD). P_{ET}CO₂, P_{ET}O₂ and HR obtained during visual stimulation task. CO obtained pre-MRI and post-MRI (20–25 min after CO delivery). Estimated COHb included. Paired t-tests within visit * $p < .05$, ** $p < .01$, *** $p < .0001$.




Figure 3. BOLD fMRI response associated with breath-by-breath end-tidal CO₂ during the breath hold task. Whole-brain analysis. Images are colour-rendered statistical maps (Z scores) superimposed on a standard (MNI) brain. Significant regions are displayed with a threshold of $Z > 3.1$ with a cluster probability threshold of $p < 0.05$ (corrected for multiple comparisons). Maps are BOLD response associated with air and CO inhalation (pre- and post-intervention), pre versus post-intervention difference maps for each protocol (Δ), and contrasts between protocols (contrast between the pre- vs. post-intervention difference maps). For contrasts, blue-lightblue indicates where BOLD response following CO (i.e. CO(post>pre)) was lower than BOLD response following air – i.e. on the day the participants inhaled CO, the BOLD response was reduced in the post-inhalation scan, but this did not occur on the day the participants inhaled air. This difference between protocols was significant. In no area was BOLD response following CO increased compared to BOLD response following Air. **A** shows areas where change in BOLD fMRI response for the CO protocol correlates with individual rise in CO level. Red-yellow indicates a positive correlation.

indicate a significant linear regression of the end-tidal CO₂ and BOLD response, as expected with rises in P_{ET}CO₂. This was seen for the control protocol and the baseline scan (pre) for the CO protocol, but not after CO inhalation. Following CO inhalation, activation was significantly reduced in the left insula, premotor cortex, left secondary somatosensory cortex and in the brain stem (see Supplement Figure S6). Group comparisons showed lower significant BOLD response change in the CO protocol compared to air in the left operculum and insula.

CVR maps for the breath hold task can be found in the supplement (Figure S5). Statistical comparisons showed no difference in mean %BOLD/mmHg between baseline (pre-intervention) scans, but a statistically significant difference in post-intervention scans (CVR lower in the CO protocol, $p = 0.048$).

Visual task (Figure 4). The flashing checkerboard generated significant BOLD activation in the visual cortex for all scans. Group comparisons showed lower activation in response to the task in the CO protocol compared to air. This was observed in the visual cortex.

Motor task (Figure 5). The finger-tapping task generated significant task-related BOLD activation in the left primary and secondary somatosensory cortices, the left premotor and primary motor cortices, the left thalamus and the visual cortex for all scans. Group comparisons showed lower activation in response to the task in the CO protocol compared to air in the visual cortex. The group analysis at a lower cluster-forming threshold also showed higher activation in the CO protocol compared to air in the premotor cortex (Figure S10).

Discussion

Key findings

In this study, we show that a small amount of inhaled CO, raising expired levels from ~3ppm to ~6 ppm, significantly alters BOLD response in never-smokers.




Figure 4. BOLD fMRI response during visual stimulus. Whole-brain analysis. Images are colour-rendered statistical maps superimposed on a standard (MNI) brain. Significant regions are displayed with a threshold of $Z > 3.1$ with a cluster probability threshold of $p < 0.05$ (corrected for multiple comparisons). Maps are BOLD response associated with air and CO inhalation (pre- and post-intervention), pre versus post-intervention difference maps for each protocol (Δ), and contrasts between protocols (contrast between the pre- vs. post-intervention difference maps). For contrasts, blue-lightblue indicates where BOLD response following CO (i.e. CO(post>pre)) was lower than BOLD response following air – i.e. on the day the participants inhaled CO, the BOLD response was reduced in the post-inhalation scan, but this did not occur on the day the participants inhaled Air. This difference between protocols was significant. In no area was (CO(post>pre)) greater than (Air(post>pre)).

This suggests that CO, even in low doses, might be a confound in BOLD fMRI. Systematic differences in COHb between, e.g., a patient group consisting of a greater proportion of smokers and a control group of predominantly non-smokers could generate group differences that are CO-related rather than associated with the specific research outcome. This could affect the results of clinical trials and patient-oriented neuroscience research. Given that absolute measures of CBF were not obtained in this study, the mechanism underlying the observed impact remains unknown (although potential mechanisms are discussed below). Future studies should incorporate flow measurements using techniques such as arterial spin labelling to further elucidate the effect of CO on BOLD fMRI.




Figure 5. BOLD fMRI response during motor task. Whole-brain analysis. Images are colour-rendered statistical maps superimposed on a standard (MNI) brain. Significant regions are displayed with a threshold of $Z > 3.1$ with a cluster probability threshold of $p < 0.05$ (corrected for multiple comparisons). Maps are BOLD response associated with air and CO inhalation (pre- and post-intervention), and contrasts between protocols (contrast between the pre- vs. post-intervention difference maps). For contrasts, blue-lightblue indicates where the BOLD response following CO (i.e. CO(post>pre)) was lower than the BOLD response following air.

Discussion of findings

Studies have shown that global baseline increases in CBF can reduce or alter task-related BOLD response. For example, Cohen et al.²⁸ used experimentally induced hypercapnia to reduce visual activation, and Brown et al.²⁹ showed that the cerebral vasodilator acetazolamide can dampen motor activation. Similarly, Halani et al.¹¹ showed that BOLD response could be modulated by changes in baseline CBF (induced by hypercapnia) as this altered the time course of cerebrovascular responses. Yet this effect has not yet been linked to CO exposure.

To probe the vulnerability of the BOLD response to COHb elevation, we employed a low-level increase in inhaled CO, raising exhaled levels to the lowest associated with tobacco smoking. Using this minimal level, we observed significant effects on BOLD response

during a hypercapnic challenge and during commonly used visual and motor tasks. The large impact of low-level CO exposure on common fMRI paradigms such as a simple flashing checkerboard and finger-tapping tasks highlights the relevance of the present findings.

The effect of CO on BOLD response was not uniform. BOLD response changes associated with visual and motor tasks were impacted by CO exposure. Compared to air, the BOLD response changes were lower for the visual task and lower in the visual cortex during the motor task, but analysis using a lower cluster-forming threshold of 2.3 also showed an area of activation in the motor cortex that was higher compared to air during the motor task (see Figure S10). The reduction in visual cortex signal mirrors that for the visual task, and may be associated with the visual instructions on screen throughout tapping intervals. The agreement in CO-impact on BOLD response in the visual cortex between these two separate and different tasks is encouraging. However, the higher BOLD response in the motor cortex after CO exposure compared to after air, despite only surviving at a lower cluster-forming threshold (Figure S10, S11), could indicate that the impact of CO on global fMRI signal might be complex. Studies have shown that hypercapnia may affect BOLD response differently depending on the type of task and activated brain regions. For example, Kastrup et al.³⁰ reported that BOLD response changes with hypercapnia were greater in the visual cortex than in the sensorimotor cortex, possibly due to the location of large veins and/or neural activity associated with respiratory stimuli.³¹ Bright et al.¹² have shown that there are regional differences in optimal haemodynamic delay under hypercapnic conditions, with the visual cortex trending towards lower optimal delay than, e.g., the parietal lobe. It is possible that the regional variations observed in our study is, in part, due to variations in optimal haemodynamic delay although further studies are required to fully elucidate the underlying mechanism.

Potential mechanisms

At present, we cannot be certain of the mechanism(s) underlying CO mediation of BOLD response. Acute CO exposure can cause cerebral vasodilation both directly³² and indirectly through nitric oxide,³³ and changes in CBF can impact BOLD response in a variety of ways.³⁴ Increases in baseline CBF can create ceiling effects, thus reducing task-related signal.³⁴ For example, hypercapnia has been shown to increase baseline CBF, reduce BOLD activation^{28,34} and alter the time course of the BOLD response^{11,12} in a potentially region-specific manner.¹² While it is difficult to draw direct comparisons between hypercapnia and CO

exposure, particularly during breath holds as CO and CO₂ may interact,³⁵ it is possible that similar mechanisms underlie our findings. Hypercapnia has also been shown to reduce CVR¹³ (but see also¹²). Indeed, we observed that CVR following CO inhalation was significantly reduced compared to CVR following air inhalation, suggesting that CO may affect BOLD response at least partly through changes in CVR. In summary, both alteration of the BOLD response time course and changes in CVR may explain the impact of CO on BOLD response in our study.

Other mechanisms that may contribute to the observed BOLD response change include the formation of COHb at the expense of oxyhaemoglobin. This may cause increased CBF through the development of hypoxia,³⁶ and may be augmented by the presence of hypercapnia.³⁵ While we observed reduced P_{ET}O₂ during the second scan on each experimental day, this was similar for both protocols, and may thus rather be due to altered breathing patterns during the experimental protocol despite pre-scan acclimatization to the breathing system. Furthermore, P_{ET}O₂ remained within normal range throughout the experiment. It is therefore unlikely that hypoxia is the cause of the observed group differences. Hypoxia may, however, contribute to BOLD response changes at higher doses of CO. Another way in which CO could reduce BOLD response is by shifting the oxygen dissociation curve to the left, reducing oxygen availability. While this is unlikely to be the mechanism in the present study, given the low levels of COHb observed, the impact of CO on physiology is complex and should not be ignored.

CO may also slightly inhibit cell respiration even under normoxic conditions³⁷ and it remains unknown whether the observed effect on BOLD response is linked in part to metabolic modulation. Similarly, we cannot rule out the possibility that CO altered BOLD response through its role as an endogenous neurotransmitter.³⁸

Participants showed no change in reaction times with CO compared to air, no difference in anxiety scores, and were not able to tell which protocol they were undertaking when prompted. It is thus unlikely that the effect on BOLD response observed in our study is driven by behavioural factors.

While further work is required to elucidate the precise mechanism underlying our findings, it is clear that CO can alter BOLD response, and should be considered a non-negligible neuroimaging confound. Further work should include formal comparisons between smokers and non-smokers to determine the impact of CO on BOLD response in a wider population beyond our tightly-controlled sample.

Brain regions

As the purpose of the study was to assess whether there is an overall, global effect of CO on BOLD response rather than interrogating specific neural responses, tasks were not linked to behavioural measures. Consequently, the following interpretation of BOLD response change patterns is speculative in nature as it relies upon reverse inference, and will be kept short. We observed BOLD response reductions after CO for the visual task in the visual cortex (as expected for this task), although this was not found in the contrast between protocols. The BOLD response change outside of the visual cortex was not significant in either mean analysis, suggesting that protocol differences were driven by small variations in signal between protocols (see e.g. Herigstad et al.³⁹). Similarly, group analysis showed reduced BOLD response change in response to the breath hold task in the left insula, which is associated with breathing challenges and anticipation of the same.^{39–42} Mean contrasts also highlighted differences in the premotor cortex, left secondary somatosensory cortex, left supramarginal gyrus^{39,40} and in the brain stem.^{40,43} The reason for the lateralisation remains unknown, but may be due to left-lateralisation associated with reading.⁴⁴ It is thus possible that there are effects of CO on BOLD response that are specific to respiratory processing centres, and that tasks probing such regions could be particularly susceptible to CO effects, although further studies incorporating appropriate behavioural measures are required to determine if this is the case.

Implications for neuroimaging and clinical trials

In this study, we show that low-level CO exposure may significantly alter BOLD response. Due to its affinity for haemoglobin, CO is not readily removed and therefore its effects on signal could persist for some time following inhalation. Here, CO assessments made following the scan (approximately 20 and 30 min after the intervention) show steady, elevated levels of exhaled CO (Figure 2). This level of CO exhalation is at the lower end of that associated with smokers, with mean exhaled values being more than 20 ppm in outpatient groups.¹⁰ It remains unknown if higher levels of CO exposure will have a greater effect (i.e. a dose-dependent effect similar to that observed in rat aortas⁴⁵). Furthermore, the findings observed in this paper suggest that the effect may be region- and/or task-dependent, which could complicate any potential adjustments for COHb during analysis.

Smoking is associated with a range of diseases, including cardiorespiratory diseases, cancers, dementia and cognitive decline⁴⁶ and several mental disorders,⁴⁷

as well as demographic factors such as socioeconomic status, education and income level.⁴⁸ CO exposure through cigarette smoking could therefore constitute a significant confound in neuroimaging research. Differences in COHb may occur both longitudinally (e.g. if smoking participants or patients are encouraged to stop smoking) and whenever participants or patients are compared with controls that are not precisely matched for smoking behaviour. Furthermore, the possibility for dose-dependent effects means that it may not be sufficient to match simply for 'smoker' and 'non-smoker', but rather the amount of COHb present in the blood stream. Given that only a small increase in COHb might affect BOLD response, this confound should be monitored carefully, particularly in clinical trials.

Conclusions

We conclude that even small amounts of inhaled CO might significantly alter BOLD response during simple tasks such as breath hold, visual stimulation and finger-tapping. Further research is required to assess the precise underlying mechanism of this effect as well as generalisability to a wider population including smokers. We suggest that care should be taken to include CO as a potential confound in neuroimaging research when appropriate, for example in studies on clinical populations with greater/lower prevalence of smokers.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Acknowledgements

We would like to thank Steve Knight for his generous assistance with data collection, and Dr Olivia Faull, Dr Anja Hayen and Dr Kyle Pattinson for their invaluable feedback on the analysis and manuscript. This study was funded by the Oxford Brookes University Central Research Fund.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Authors' contributions

CB contributed to the data acquisition, manuscript draft and approval. SM contributed to the concept and design, manuscript revision and approval. MH contributed to the concept and design, data acquisition, analysis, interpretation of data, manuscript draft and approval.

Supplemental material

Supplemental material for this article is available online.

ORCID iDs

Shakeeb H Moosavi  <https://orcid.org/0000-0002-9709-7675>

Mari Herigstad  <https://orcid.org/0000-0003-1053-2038>

References

- Kim SG and Ogawa S. Biophysical and physiological origins of blood oxygenation level-dependent fMRI signals. *J Cereb Blood Flow Metab* 2012; 32: 1188–1206.
- Hanafy KA, Oh J and Otterbein LE. Carbon monoxide and the brain: time to rethink the dogma. *Curr Pharm Des* 2013; 19: 2771–2775.
- Komuro T, Borsody MK, Ono S, et al. The vasorelaxation of cerebral arteries by carbon monoxide. *Exp Biol Med* 2001; 226: 860–865.
- Kanu A, Whitfield J and Leffler CW. Carbon monoxide contributes to hypotension-induced cerebrovascular vasodilation in piglets. *Am J Physiol Heart Circ Physiol* 2006; 291: H2409–H2414.
- Doblar DD, Santiago TV and Edelman NH. Correlation between ventilatory and cerebrovascular responses to inhalation of CO. *J Appl Physiol Respir Environ Exerc Physiol* 1977; 43: 455–462.
- Meilin S, Rogatsky GG, Thom SR, et al. Effects of carbon monoxide on the brain may be mediated by nitric oxide. *J Appl Physiol* 1996; 81: 1078–1083.
- Mendelman A, Zarchin N, Meilin S, et al. Blood flow and ionic responses in the awake brain due to carbon monoxide. *Neurol Res* 2002; 24: 765–772.
- Benignus VA, Petrovick MK, Newlin-Clapp L, et al. Carboxyhemoglobin and brain blood flow in humans. *Neurotoxicol Teratol* 1992; 14: 285–290.
- Paulson OB, Parving HH, Olesen J, et al. Influence of carbon monoxide and of hemodilution on cerebral blood flow and blood gases in man. *J Appl Physiol* 1973; 35: 111–116.
- Middleton ET and Morice AH. Breath carbon monoxide as an indication of smoking habit. *Chest* 2000; 117: 758–763.
- Halani S, Kwinta JB, Golestani AM, et al. Comparing cerebrovascular reactivity measured using BOLD and cerebral blood flow MRI: the effect of basal vascular tension on vasodilatory and vasoconstrictive reactivity. *Neuroimage* 2015; 110: 110–123.
- Bright MG, Donahue MJ, Duyn JH, et al. The effect of basal vasodilation on hypercapnic and hypocapnic reactivity measured using magnetic resonance imaging. *J Cereb Blood Flow Metab* 2011; 31: 426–438.
- Zhao P, Alsop DC, Abduljalil A, et al. Vasoreactivity and peri-infarct hyperintensities in stroke. *Neurology* 2009; 72: 643–649.
- Kim SG and Ugurbil K. Comparison of blood oxygenation and cerebral blood flow effects in fMRI: estimation of relative oxygen consumption change. *Magn Reson Med* 1997; 38: 59–65.
- Rostrup E, Law I, Blinkenberg M, et al. Regional differences in the CBF and BOLD responses to hypercapnia: a combined PET and fMRI study. *Neuroimage* 2000; 11: 87–97.
- Pattinson KT, Governo RJ, MacIntosh BJ, et al. Opioids depress cortical centers responsible for the volitional control of respiration. *J Neurosci* 2009; 29: 8177–8186.
- Pattinson KT, Rogers R, Mayhew SD, et al. Pharmacological fMRI: measuring opioid effects on the BOLD response to hypercapnia. *J Cereb Blood Flow Metab* 2007; 27: 414–423.
- Tancredi FB and Hoge RD. Comparison of cerebral vascular reactivity measures obtained using breath-holding and CO₂ inhalation. *J Cereb Blood Flow Metab* 2013; 33: 1066–1074.
- Mumford JA and Nichols TE. Power calculation for group fMRI studies accounting for arbitrary design and temporal autocorrelation. *Neuroimage* 2008; 39: 261–268.
- Spielberger CD, Gorsuch RL, Lushene R, et al. *Manual for the state-trait anxiety inventory*. Palo Alto, USA: Consulting Psychologists Press, 1989.
- Jenkinson M, Bannister P, Brady M, et al. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* 2002; 17: 825–841.
- Woolrich MW, Ripley BD, Brady M, et al. Temporal autocorrelation in univariate linear modeling of fMRI data. *Neuroimage* 2001; 14: 1370–1386.
- Brooks JC, Beckmann CF, Miller KL, et al. Physiological noise modelling for spinal functional magnetic resonance imaging studies. *Neuroimage* 2008; 39: 680–692.
- Bright MG and Murphy K. Reliable quantification of BOLD fMRI cerebrovascular reactivity despite poor breath-hold performance. *Neuroimage* 2013; 83: 559–568.
- Beckmann CF, Jenkinson M and Smith SM. General multilevel linear modeling for group analysis in fMRI. *Neuroimage* 2003; 20: 1052–1063.
- Woolrich MW, Behrens TE, Beckmann CF, et al. Multilevel linear modelling for fMRI group analysis using Bayesian inference. *Neuroimage* 2004; 21: 1732–1747.
- Eklund A, Nichols TE and Knutsson H. Cluster failure: why fMRI inferences for spatial extent have inflated false-positive rates. *Proc Natl Acad Sci U S A* 2016; 113: 7900–7905.
- Cohen ER, Ugurbil K and Kim SG. Effect of basal conditions on the magnitude and dynamics of the blood oxygenation level-dependent fMRI response. *J Cereb Blood Flow Metab* 2002; 22: 1042–1053.
- Brown GG, Eyler Zorrilla LT, Georgy B, et al. BOLD and perfusion response to finger-thumb apposition after acetazolamide administration: differential relationship to global perfusion. *J Cereb Blood Flow Metab* 2003; 23: 829–837.
- Kastrup A, Kruger G, Glover GH, et al. Regional variability of cerebral blood oxygenation response to hypercapnia. *Neuroimage* 1999; 10: 675–681.
- Corfield DR and McKay LC. Regional cerebrovascular responses to hypercapnia and hypoxia. *Adv Exp Med Biol* 2016; 903: 157–167.

32. Peers C, Boyle JP, Scragg JL, et al. Diverse mechanisms underlying the regulation of ion channels by carbon monoxide. *Br J Pharmacol* 2015; 172: 1546–1556.
33. Thorup C, Jones CL, Gross SS, et al. Carbon monoxide induces vasodilation and nitric oxide release but suppresses endothelial NOS. *Am J Physiol* 1999; 277: F882–F889.
34. Mark CI, Mazerolle EL and Chen JJ. Metabolic and vascular origins of the BOLD effect: implications for imaging pathology and resting-state brain function. *J Magn Reson Imaging* 2015; 42: 231–246.
35. Levin BC, Paabo M, Gurman JL, et al. Toxicological interactions between carbon monoxide and carbon dioxide. *Toxicology* 1987; 47: 135–164.
36. Harris AD, Murphy K, Diaz CM, et al. Cerebral blood flow response to acute hypoxic hypoxia. *NMR Biomed* 2013; 26: 1844–1852.
37. D'Amico G, Lam F, Hagen T, et al. Inhibition of cellular respiration by endogenously produced carbon monoxide. *J Cell Sci* 2006; 119: 2291–2298.
38. Verma A, Hirsch DJ, Glatt CE, et al. Carbon monoxide: a putative neural messenger. *Science* 1993; 259: 381–384.
39. Herigstad M, Hayen A, Evans E, et al. Dyspnea-related cues engage the prefrontal cortex: evidence from functional brain imaging in COPD. *Chest* 2015; 148: 953–961.
40. Hayen A, Wanigasekera V, Faull OK, et al. Opioid suppression of conditioned anticipatory brain responses to breathlessness. *Neuroimage* 2017; 150: 383–394.
41. Herigstad M, Faull OK, Hayen A, et al. Treating breathlessness via the brain: changes in brain activity over a course of pulmonary rehabilitation. *Eur Respir J* 2017; 50: 1701029.
42. Faull OK and Pattinson KT. The cortical connectivity of the periaqueductal gray and the conditioned response to the threat of breathlessness. *Elife* 2017; 6: e21749.
43. Pattinson KT, Mitsis GD, Harvey AK, et al. Determination of the human brainstem respiratory control network and its cortical connections in vivo using functional and structural imaging. *Neuroimage* 2009; 44: 295–305.
44. Strother L, Zhou Z, Coros AK, et al. An fMRI study of visual hemifield integration and cerebral lateralization. *Neuropsychologia* 2017; 100: 35–43.
45. Decaluwe K, Pauwels B, Verpoest S, et al. Divergent mechanisms involved in CO and CORM-2 induced vasorelaxation. *Eur J Pharmacol* 2012; 674: 370–377.
46. Ostergaard L, Engedal TS, Moreton F, et al. Cerebral small vessel disease: capillary pathways to stroke and cognitive decline. *J Cereb Blood Flow Metab* 2016; 36: 302–325.
47. Tidey JW and Miller ME. Smoking cessation and reduction in people with chronic mental illness. *BMJ* 2015; 351: h4065.
48. Casetta B, Videla AJ, Bardach A, et al. Association between cigarette smoking prevalence and income level: a systematic review and meta-analysis. *Nicotine Tob Res* 2016; 19: 1401–1407.