



Complete Genome Sequence of *Lactococcus piscium* CNCM I-4031, a Bioprotective Strain for Seafood Products

Laurent Marché,^a Taous Saraoui,^{a,b*} Benoit Remenant,^{a*} Monique Zagorec,^a Hervé Prévost,^a Christine Delbarre-Ladrat,^b Françoise Leroi,^b Marie France Pilet^a

UMR1014 SECALIM, INRA, Oniris, Nantes, France^a; Ifremer, Laboratoire Ecosystèmes Microbiens et Molécules Marines pour les Biotechnologies (EM³B), Nantes, France^b

ABSTRACT *Lactococcus piscium* CNCM I-4031 is a psychrotrophic foodborne lactic acid bacterium showing potential interest for the biopreservation of seafood products due to its inhibition properties toward pathogenic and spoilage bacteria. The analysis of its genome will provide a better understanding of the mechanisms of interaction between these bacteria.

Lactococcus piscium is a psychrotrophic species first isolated in 1990 from rainbow trout (1). Studies have reported the presence of this species in a large variety of food products, including meat, seafood, and vegetables (2, 3). *L. piscium* CNCM I-4031 (also named EU2241), isolated from raw salmon packs (4), might present interests for seafood biopreservation: it improves the sensory quality of seafood and limits the growth of spoilage (*Brochothrix thermosphacta* and *Serratia proteamaculans*) or pathogenic (*Listeria monocytogenes*) bacteria (5–7). Although the mechanisms of growth inhibition have not yet been elucidated, a previous study suggested the requirement for a contact between *L. piscium* CNCM I-4031 and the target-inhibited strain as well as putative involvement of quorum sensing (8).

The whole-genome sequencing of *L. piscium* CNCM I-4031 was carried out by Eurofins MWG Operon laboratories (Ebersberg, Germany) using GS FLX/FLX+ technologies with shotgun and 8-kbp-long paired-end libraries. The genome sequencing coverage is 63×. The assembly realized with Newbler 2.6 (9) showed an N_{50} of 144,750 bp for 31 contigs. The final assembly gave two scaffolds with seven contigs and a genome size of 2.26 Mbp, in accordance with results obtained by pulsed-field gel electrophoresis (PFGE) (data not shown). The first scaffold (2.06 Mb), composed of six contigs, is the complete sequence of the chromosome, with a G+C content of 38.74%. The second scaffold (one contig) formed a 20-kb plasmid with a G+C content of 34.36%. Automatic annotation and manual curation were performed on the whole genome thanks to the MicroScope platform (10, 11) (Genoscope, Évry, France). The results showed that *L. piscium* CNCM I-4031 chromosome contains 2,205 predicted coding DNA sequences (CDSs), as observed in *Lactococcus lactis* genomes (around 2,500).

Thirty-one percent of the predicted CDSs were assigned to proteins of unknown functions, 40% were predicted as enzymes, 6% as transcriptional factors, and 4% as proteins of cell structure. The remaining proteins are involved in diverse cell processes (including carriers, lipoproteins, receptors, and other membrane proteins) or from an extrachromosomal origin. Seventy-five genes (3.4%) were manually curated as carbohydrate-active enzymes (CAZymes) that degrade or modify carbohydrates or create glycosidic bonds.

Received 14 November 2016 **Accepted** 22 November 2016 **Published** 26 January 2017

Citation Marché L, Saraoui T, Remenant B, Zagorec M, Prévost H, Delbarre-Ladrat C, Leroi F, Pilet MF. 2017. Complete genome sequence of *Lactococcus piscium* CNCM I-4031, a bioprotective strain for seafood products. Genome Announc 5:e01510-16. <https://doi.org/10.1128/genomeA.01510-16>.

Copyright © 2017 Marché et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marie France Pilet, marie-france.pilet@oniris-nantes.fr.

* Present address: Taous Saraoui, Laboratoire de Biologie, Clermont Université, Université d'Auvergne, IUT Aurillac, Clermont-Ferrand, France; Benoit Remenant, Laboratoire de la Santé des Végétaux (LSV), Anses, Angers, France.

The plasmid encompasses 26 CDSs, including a complete ribose gene cluster, *rbsRKDACB*. The *rbs* genes are involved in ribose utilization but have also been reported to be responsible for internalization of autoinducer II (AI-2) involved in quorum sensing (12, 13).

The comparison of *L. piscium* CNCM I-4031 genome with that of *L. piscium* MKFS47, a spoiler strain of meat products (14), revealed 384 *L. piscium* CNCM I-4031-specific genes.

Accession number(s). This whole-genome shotgun project has been deposited in ENA under the accession no. [FLZT01000001](https://ena.ebi.ac.uk/ena/record/FLZT01000001) to [FLZT01000007](https://ena.ebi.ac.uk/ena/record/FLZT01000007). The versions described in this paper are the first versions.

ACKNOWLEDGMENTS

The LABGeM (CEA/IG/Genoscope & CNRS UMR 8030) and the France Génomique National infrastructure (funded as part of Investissement d'Avenir Program managed by Agence Nationale pour la Recherche, contract ANR-10-INBS-09) are acknowledged for support within the MicroScope annotation platform. Taous Saraoui was the recipient of a Ph.D. fellowship program from the French Ministry of Higher Education and Research. This study was financially supported by the Région des Pays de la Loire, France within the context of the COM-BACT research program (contract 2014 07957).

REFERENCES

- Williams AM, Fryer JL, Collins MD. 1990. *Lactococcus piscium* sp. nov. a new *Lactococcus* species from salmonid fish. FEMS Microbiol Lett 68: 109–113. <https://doi.org/10.1111/j.1574-6968.1990.tb04132.x>.
- Sakala RM, Hayashidani H, Kato Y, Kaneuchi C, Ogawa M. 2002. Isolation and characterization of *Lactococcus piscium* strains from vacuum-packaged refrigerated beef. J Appl Microbiol 92:173–179. <https://doi.org/10.1046/j.1365-2672.2002.01513.x>.
- Vihavainen E, Lundström HS, Susiluoto T, Koort J, Paulin L, Auvinen P, Björkroth KJ. 2007. Role of broiler carcasses and processing plant air in contamination of modified-atmosphere-packaged broiler products with psychrotrophic lactic acid bacteria. Appl Environ Microbiol 73: 1136–1145. <https://doi.org/10.1128/AEM.01644-06>.
- Matamoros S, Pilet MF, Gigout F, Prévost H, Leroi F. 2009. Selection and evaluation of seafood-borne psychrotrophic lactic acid bacteria as inhibitors of pathogenic and spoilage bacteria. Food Microbiol 26: 638–644. <https://doi.org/10.1016/j.fm.2009.04.011>.
- Fall PA, Pilet MF, Leduc F, Cardinal M, Duflos G, Guérin C, Joffraud JJ, Leroi F. 2012. Sensory and physicochemical evolution of tropical cooked peeled shrimp inoculated by *Brochothrix thermosphacta* and *Lactococcus piscium* CNCM I-4031 during storage at 8°C. Int J Food Microbiol 152: 82–90. <https://doi.org/10.1016/j.ijfoodmicro.2011.07.015>.
- Leroi F, Cornet J, Chevalier F, Cardinal M, Coeuret G, Chaillou S, Joffraud JJ. 2015. Selection of bioprotective cultures for preventing cold-smoked salmon spoilage. Int J Food Microbiol 213:79–87. <https://doi.org/10.1016/j.ijfoodmicro.2015.05.005>.
- Fall PA, Leroi F, Chevalier F, Guérin C, Pilet M-F. 2010. Protective effect of a non-bacteriocinogenic *Lactococcus piscium* CNCM I-4031 strain against *Listeria monocytogenes* in sterilized tropical cooked peeled shrimp. J Aquat Food Prod Technol 19:84–92. <https://doi.org/10.1080/10498850.2010.486910>.
- Saraoui T, Fall PA, Leroi F, Antignac JP, Chéreau S, Pilet MF. 2016. Inhibition mechanism of *Listeria monocytogenes* by a bioprotective bacterium *Lactococcus piscium* CNCM I-4031. Food Microbiol 53:70–78. <https://doi.org/10.1016/j.fm.2015.01.002>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. Genome Res 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fèvre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AA, Weiman M, Médigue C. 2013. MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. Nucleic Acids Res 41:D636–D647. <https://doi.org/10.1093/nar/gks1194>.
- Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S, Lajus A, Pascal G, Scarpelli C, Médigue C. 2006. MaGe: a microbial genome annotation system supported by synteny results. Nucleic Acids Res 34:53–65. <https://doi.org/10.1093/nar/gkj406>.
- Taga ME, Semmelhack JL, Bassler BL. 2001. The LuxS-dependent auto-inducer AI-2 controls the expression of an ABC transporter that functions in AI-2 uptake in *Salmonella* Typhimurium. Mol Microbiol 42:777–793. <https://doi.org/10.1046/j.1365-2958.2001.02669.x>.
- Armbruster CE, Pang B, Murrain K, Juneau RA, Perez AC, Weimer KED, Swords WE. 2011. RbsB (NTHI_0632) mediates quorum signal uptake in nontypeable *Haemophilus influenzae* strain 86–028NP. Mol Microbiol 82:836–850. <https://doi.org/10.1111/j.1365-2958.2011.07831.x>.
- Andreevskaya M, Johansson P, Laine P, Smolander OP, Sonck M, Rahkila R, Jääskeläinen E, Paulin L, Auvinen P, Björkroth J. 2015. Genome sequence and transcriptome analysis of meat spoilage lactic acid bacterium *Lactococcus piscium* MKFS47. Appl Environ Microbiol 81: 3800–3811. <https://doi.org/10.1128/AEM.00320-15>.