

MEETING ABSTRACT

Open Access

# OR9-001 - Exome sequencing in monogenic Behçet-like disease

Q Zhou<sup>1\*</sup>, R Laxer<sup>2</sup>, M Pelletier<sup>3</sup>, M Ramaswamy<sup>3</sup>, H-Y Wang<sup>4</sup>, D Chin<sup>1</sup>, A Güll<sup>5</sup>, C Sibley<sup>6</sup>, M Barat-Houari<sup>7</sup>, R Siegel<sup>3</sup>, DL Kastner<sup>1</sup>, I Aksentijevich<sup>1</sup>

From 7th Congress of International Society of Systemic Auto-Inflammatory Diseases (ISSAID)  
Lausanne, Switzerland. 22-26 May 2013

## Introduction

A Caucasian family with an affected mother and 2 affected daughters presented with early onset Behçet-like disease, manifesting with arthralgia/arthritis, mouth and genital ulcers, and uveitis. They are negative for *HLA\*B51*. Their symptoms are significantly ameliorated with TNF-inhibitors.

## Objectives

To identify the causative mutation(s) in this family we sequenced the exomes of the 3 affected patients.

## Methods

Exome data were filtered for novel variants not present in dbSNP, the 1000 Genomes Pilot Project, NHLBI Exome Sequencing Project (ESP5400), and 124 exomes from in-house data.

## Results

We identified 21 putative candidate variants that are both novel and consistent with dominant inheritance. Sanger sequencing validated all 21 variants. Three variants identified in *TNFRSF9*, *MGEA5*, and *TNFAIP3* genes were confirmed to have arisen *de novo* in the affected mother based on the genotyping of the healthy maternal grandmother, the maternal unaffected brother, and the unaffected paternal aunt. The candidate variant in *MGEA5* was predicted as benign by PolyPhen-2. The 2 candidate variants that remained for consideration are p.C78W (*TNFRSF9*; CD137; 4-1BB) and p.L227X (*TNFAIP3*; A20). Haplotype analysis showed that p.C78W occurred *de novo* on the haplotype inherited from the grandmother. The p.L227X mutation-associated haplotype was found on the

grandfather's haplotype; his sample is not available for analysis. Because we reached the limit for further analysis of candidate variants in the family, we studied 6 Turkish familial Behçet cases and 56 sporadic Caucasian cases. All of these individuals were negative for mutations in either candidate gene. Both 4-1BB and A20 are strong candidates and potentially in the same signaling pathway. 4-1BB is a TNF-family receptor that costimulates T cell responses and promotes survival of lymphocytes and dendritic cells; A20 is a negative regulator of NF-κB activation by TNF and TLR family receptors. Mutant 4-1BB expressed in Jurkat cells was associated with reduced expression on the plasma membrane, and activated T cells from all 3 patients had a marked decrease in surface expression, especially in CD8<sup>+</sup> T cells. Despite reduced surface expression, the C78W TNFRSF9 mutation could have a gain-of-function phenotype like TNFR1 mutations in TRAPS, leading to intracellular retention of mutant protein and spontaneous signaling, which in this case could amplify immune responses by T cells and other cell types expressing 4-1BB. Patient peripheral blood cells also had lower total A20 protein levels relative to controls and, consistent with A20 lack of function, increased I-κB degradation after cellular activation.

## Conclusion

Behçet-like disease in this family is associated with mutations in 2 genes that affect immune cell survival and production of inflammatory cytokines. Additional functional studies will determine whether 1 or both mutations may contribute to this dominantly-inherited phenotype.

## Competing interests

None declared.

<sup>1</sup>Inflammatory Disease Section, National Human Genome Research Institute (NHGRI/NIH), Bethesda, USA

Full list of author information is available at the end of the article

**Authors' details**

<sup>1</sup>Inflammatory Disease Section, National Human Genome Research Institute (NHGRI/NIH), Bethesda, USA. <sup>2</sup>Hospital for Sick Children, Toronto, Canada.

<sup>3</sup>Immunoregulation Section, National Institute of Arthritis and Musculoskeletal and Skin (NIAMS/NIH), USA. <sup>4</sup>Laboratory of Host Defenses, National Institute of Allergy and Infectious Diseases (NIAID/NIH), Bethesda, USA. <sup>5</sup>Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey.

<sup>6</sup>Office of Clinical Director, National Institute of Arthritis and Musculoskeletal and Skin (NIAMS/NIH), Bethesda, USA. <sup>7</sup>Département de Génétique, LBM -CHU de Montpellier, Montpellier, France.

Published: 8 November 2013

doi:10.1186/1546-0096-11-S1-A184

**Cite this article as:** Zhou et al.: OR9-001 - Exome sequencing in monogenic Behçet-like disease. *Pediatric Rheumatology* 2013 **11**(Suppl 1): A184.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

