

Draft Genome Sequences of Two Strains of *Serratia* spp. from the Midgut of the Malaria Mosquito *Anopheles gambiae*

Dong Pei,^a Casey Hill-Clemons,^a Guillaume Carissimo,^b Wanqin Yu,^a Kenneth D. Vernick,^{b,c} Jiannong Xu^a

Biology Department, Molecular Biology Program, New Mexico State University, Las Cruces, New Mexico, USA^a; Pasteur Institute, Paris, France^b; Department of Microbiology, University of Minnesota, Minneapolis, Minnesota, USA^c

Here, we report the annotated draft genome sequences of two strains of *Serratia* spp., Ag1 and Ag2, isolated from the midgut of two different strains of *Anopheles gambiae*. The genomes of these two strains are almost identical.

Received 23 January 2015 Accepted 4 February 2015 Published 12 March 2015

Citation Pei D, Hill-Clemons C, Carissimo G, Yu W, Vernick KD, Xu J. 2015. Draft genome sequences of two strains of *Serratia* spp. from the midgut of the malaria mosquito *Anopheles gambiae*. *Genome Announc* 3(2):e00090-15. doi:10.1128/genomeA.00090-15.

Copyright © 2015 Pei et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Jiannong Xu, jxu@nmsu.edu.

The genus *Serratia* consists of Gram-negative rod-shaped facultative anaerobes. Members of this genus share a variety of habitats, such as freshwater, soil, plants, and the gut of invertebrates and vertebrates. The genus includes many human pathogenic strains (1). In addition, other strains have been investigated, for example, a strain from the gut of water flea *Daphnia magna* (2), and a strain that is associated with leaf-cutter ant fungus gardens (3). The mosquito gut ecosystem accommodates a dynamic microbiome (4, 5). *Serratia odorifera* could significantly enhance susceptibility to dengue virus DENV-2 in *Aedes aegypti* (6). In anopheline mosquitoes, *Serratia marcescens* has been found to be able to inhibit *Plasmodium* development in *Anopheles gambiae* (7, 8).

The *Serratia* sp. strain Ag1 was isolated from the Ngousso strain of *A. gambiae* that was originally collected in Cameroon in 2006, and maintained in the Vernick lab, the Institut Pasteur, Paris, France. The *Serratia* sp. strain Ag2 was isolated from the *A. gambiae* G3 strain in the Xu lab at New Mexico State University, Las Cruces, NM, USA. The G3 strain was derived from a collection originally in MacCarthy Island, the Gambia, and colonized in 1975. The genome was determined by Illumina paired-end technology at Genewiz, Inc. A total of 6.1 M reads for strain Ag1 and 4.5 M reads for strain Ag2 were generated. *De novo* assembly was performed using CLC genomic workbench (v7.0.4). The assembly of strain Ag1 contains 110 contigs totaling 5.35 Mbp. The assembly of strain Ag2 contains 109 contigs totaling 5.32 Mbp. The genomes were annotated by the NCBI Prokaryotic Genome Automatic Annotation Pipeline. The Ag1 genome includes 4,773 coding sequences (CDS) and 85 RNA genes. The Ag2 genome includes 4,706 CDS and 85 RNA genes. Genome comparison indicated the common origins of the isolates. Approximately 98.2% of the Ag1 genome has identical counterparts in Ag2, whereas 98.7% of the Ag2 genome has identical counterparts in Ag1. The differences between the two strains were minor. For example, contig 54 (9,245 bp) in Ag1 has no counterpart in Ag2, and contig 12 (18,403 bp) and contig 78 (15,537 bp) from Ag2 are not present in Ag1. Intriguingly, the two almost identical strains are associated with different mosquito colonies originally from different parts of

Africa, and the two colonies were never maintained in the same insectary. This strongly suggests that the bacteria are persistently associated with the natural mosquito microbial flora in nature. As a member of the gut flora of Ngousso mosquitoes, strain Ag1 may be involved in the susceptibility to *Plasmodium* and O'nyong-nyong virus (9).

Recently, several genomes of mosquito associated bacteria have been sequenced, including *Elizabethkingia anophelis* (10, 11), *Asaia* spp. (12), *Enterobacter* spp. (13), *Pseudomonas* spp. (14), and *Spiroplasma diminutum* and *S. taiwanense* (15, 16). The available genomes will facilitate to characterize the structure and function of the mosquito microbiome.

Nucleotide sequence accession numbers. The genome sequences have been deposited at GenBank under the accession numbers JQEI00000000 (*Serratia* sp. Ag1) and JQJ00000000 (*Serratia* sp. Ag2).

ACKNOWLEDGMENTS

This work was supported by NIH grant SC1AI112786 and NSF grant DMS-1222592 to J.X. and European Research Council grant 323173 to K.D.V.

REFERENCES

- Mahlen SD. 2011. *Serratia* infections: from military experiments to current practice. *Clin Microbiol Rev* 24:755–791. <http://dx.doi.org/10.1128/CMR.00017-11>.
- Poehlein A, Freese HM, Daniel R, Simeonova DD. 2014. Draft genome sequence of *Serratia* sp. strain DD3, isolated from the guts of *Daphnia magna*. *Genome Announc* 2(5):e00903-14. <http://dx.doi.org/10.1128/genomeA.00903-14>.
- Aylward FO, Tremmel DM, Starrett GJ, Bruce DC, Chain P, Chen A, Davenport KW, Detter C, Han CS, Han J, Huntemann M, Ivanova NN, Kyrpides NC, Markowitz V, Mavrommatis K, Nolan M, Pagani I, Pati A, Pitluck S, Teshima H, Deshpande S, Goodwin L, Woyke T, Currie CR. 2013. Complete genome of *Serratia* sp. strain FGI 94, a strain associated with leaf-cutter ant fungus gardens. *Genome Announc* 1(2):e00239-12. <http://dx.doi.org/10.1128/genomeA.00239-12>.
- Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, Nsango SE, Shahbazkia HR, Awono-Ambene PH, Levashina EA, Christen R, Morlais I. 2012. Midgut microbiota of the malaria mosquito vector *Anopheles gambiae* and interactions with *Plasmodium falciparum* infection. *PLoS Pathog* 8:e1002742. <http://dx.doi.org/10.1371/journal.ppat.1002742>.

5. Wang Y, Gilbreath TM III, Kukutla P, Yan G, Xu J. 2011. Dynamic gut microbiome across life history of the malaria mosquito *Anopheles gambiae* in Kenya. *PLoS One* 6:e24767. <http://dx.doi.org/10.1371/journal.pone.0024767>.
6. Apte-Deshpande A, Paingankar M, Gokhale MD, Deobagkar DN. 2012. *Serratia odorifera* a midgut inhabitant of *Aedes aegypti* mosquito enhances its susceptibility to dengue-2 virus. *PLoS One* 7:e40401. <http://dx.doi.org/10.1371/journal.pone.0040401>.
7. Bahia AC, Dong Y, Blumberg BJ, Mlambo G, Tripathi A, Benmarzouk-Hidalgo OJ, Chandra R, Dimopoulos G. 2014. Exploring *Anopheles* gut bacteria for *Plasmodium* blocking activity. *Environ Microbiol* 16: 2980–2994. <http://dx.doi.org/10.1111/1462-2920.12381>.
8. Bando H, Okado K, Guelbeogo WM, Badolo A, Aonuma H, Nelson B, Fukumoto S, Xuan X, Sagnon N, Kanuka H. 2013. Intra-specific diversity of *Serratia marcescens* in *Anopheles* mosquito midgut defines *Plasmodium* transmission capacity. *Sci Rep* 3:1641. <http://dx.doi.org/10.1038/srep01641>.
9. Carissimo G, Pondeville E, McFarlane M, Dietrich I, Mitri C, Bischoff E, Antoniewski C, Bourgouin C, Failloux A-B, Kohl A, Vernick KD. 2015. Antiviral immunity of *Anopheles gambiae* is highly compartmentalized, with distinct roles for RNA interference and gut microbiota. *Proc Natl Acad Sci U S A* 112:E176–E185. <http://dx.doi.org/10.1073/pnas.1412984112>.
10. Kukutla P, Lindberg BG, Pei D, Rayl M, Yu W, Steritz M, Faye I, Xu J. 2013. Draft genome sequences of *Elizabethkingia anophelis* strains R26T and Ag1 from the midgut of the malaria mosquito *Anopheles gambiae*. *Genome Announc* 1(6):e01030-13. <http://dx.doi.org/10.1128/genomeA.01030-13>.
11. Kukutla P, Lindberg BG, Pei D, Rayl M, Yu W, Steritz M, Faye I, Xu J. 2014. Insights from the genome annotation of *Elizabethkingia anophelis* from the malaria vector *Anopheles gambiae*. *PLoS One* 9:e97715. <http://dx.doi.org/10.1371/journal.pone.0097715>.
12. Shane JL, Bongio NJ, Favia G, Lampe DJ. 2014. Draft genome sequence of *Asaia* sp. strain SF2.1, an important member of the microbiome of *Anopheles* mosquitoes. *Genome Announc* 2(1):e01202-13. <http://dx.doi.org/10.1128/genomeA.01202-13>.
13. Jiang J, Alvarez C, Kukutla P, Yu W, Xu J. 2012. Draft genome sequences of *Enterobacter* sp. isolate Ag1 from the midgut of the malaria mosquito *Anopheles gambiae*. *J Bacteriol* 194:5481. <http://dx.doi.org/10.1128/JB.01275-12>.
14. Alvarez C, Kukutla P, Jiang J, Yu W, Xu J. 2012. Draft genome sequence of *Pseudomonas* sp. strain ag1, isolated from the midgut of the malaria mosquito *Anopheles gambiae*. *J Bacteriol* 194:5449. <http://dx.doi.org/10.1128/JB.01173-12>.
15. Chang TH, Lo WS, Ku C, Chen LL, Kuo CH. 2014. Molecular evolution of the substrate utilization strategies and putative virulence factors in mosquito-associated *Spiroplasma* species. *Genome Biol Evol* 6:500–509. <http://dx.doi.org/10.1093/gbe/evu033>.
16. Lo WS, Ku C, Chen LL, Chang TH, Kuo CH. 2013. Comparison of metabolic capacities and inference of gene content evolution in mosquito-associated *Spiroplasma diminutum* and *S. Taiwanense*. *Genome Biol Evol* 5:1512–1523. <http://dx.doi.org/10.1093/gbe/evt108>.