



WALKING POSTER PRESENTATION

Open Access

Analysis of RV components after reoperation of the right ventricular outflow tract in patients with Tetralogy of Fallot

Abdullah M AL Bulushi^{1*}, Felicitas Frettlöh¹, Dominik Daniel Gabbert¹, Ines Kristo¹, Andreas Entenmann¹, Christopher Hart¹, Inga Voges¹, Minh H Pham¹, Michael Jerosch-Herold², Philip Wegner¹, Ana C Andrade¹, Hans-Heiner Kramer¹, Carsten Rickers¹

From 18th Annual SCMR Scientific Sessions
Nice, France. 4-7 February 2015

Background

In patients with repaired Tetralogy of Fallot (TOF), reoperation of the right ventricular outflow tract (RVOT) due to pulmonary regurgitation and volume overload is often necessary. This study aims to analyse the functional recovery of the right ventricle (RV) and the RVOT after a reoperation using volumetrical data from cine MRI.

Methods

31 patients (20.6 ± 10.3 years) underwent an MRI examination before and after (20 ± 18 months) pulmonary valve replacement or reconstruction (PVR) for the volume overload of the RV. For segmental analysis of the RV, a custom made analysis software was developed. The method provides an automatic contour detection algorithm for the determination of the RV blood volume and enables the user to segregate the blood volume of the RVOT and the corpus based on anatomic landmarks and to generate volume-time-curves over a cardiac cycle. The analysis of volume-time-curves of the RV allowed calculation of Peak-Filling-Rate (PFR) and Time-to-Peak-Filling-Rate (TPFR), which are surrogate parameters for diastolic dysfunction.

Results

A significant reduction in blood volumes of the total RV was present after a PVR (RV-ESV -18.4 ± 23.6 ml/m², p=0.018, RV-EDV -28.7 ± 26.3 ml/m², p<0.01). Pulmonary insufficiency was also significantly reduced ($-48.7 \pm 48.9\%$,

p<0.001). In a subgroup of 22 pts the RVOT was separately analysed and showed a significant reduction after PVR (RVOT-ESV -5.5 ± 6.1 ml/m², p<0.001, RVOT-EDV -7.3 ± 7.9 ml/m², p<0.001). The PFR decreased (-149.2 ± 261.8 ml/s, p=0.001) and the TPFR increased (0.06 ± 0.12 s, p=0.041), indicating improvement of RV diastolic function. Additionally, late gadolinium imaging showed that non-viable tissue (patch material and scar tissue) was present in $5.7 \pm 3.8\%$ of the RV mass and in $11.9 \pm 7.2\%$ of the RVOT mass before operation and did not change on follow-up.

Conclusions

After reoperation of the RVOT in TOF patients the systolic and diastolic function of the right ventricular components (RVOT and corpus) improved within less than a year. The end-diastolic and end-systolic blood volumes decreased significantly in the RVOT despite its burden of the non-viable tissue and the total RV. The dedicated analysis software enables the user to assess the components of the RV in a time-efficient and precise manner.

Funding

N/A.

Authors' details

¹Congenital Heart Disease, Pediatric Cardiology, University Hospital Schleswig Holstein, Kiel, Germany, Kiel, Germany. ²Department of Radiology, Harvard Medical School, Women's Hospital Boston, Boston, MA, USA.

Published: 3 February 2015

¹Congenital Heart Disease, Pediatric Cardiology, University Hospital Schleswig Holstein, Kiel, Germany, Kiel, Germany

Full list of author information is available at the end of the article

doi:10.1186/1532-429X-17-S1-Q91

Cite this article as: AL Bulushi *et al.*: Analysis of RV components after reoperation of the right ventricular outflow tract in patients with Tetralogy of Fallot. *Journal of Cardiovascular Magnetic Resonance* 2015 **17** (Suppl 1):Q91.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

