

Research Article

Caffeine Effect on Cognitive Function during a Stroop Task: fNIRS Study

Yafei Yuan, Guanghao Li, Haoran Ren, and Wei Chen 

Department of Electronic Engineering, Center for Intelligent Medical Electronics, Fudan University, 200433 Shanghai, China

Correspondence should be addressed to Wei Chen; w_chen@fudan.edu.cn

Received 21 September 2020; Revised 28 October 2020; Accepted 29 October 2020; Published 21 November 2020

Academic Editor: Marco Sandrini

Copyright © 2020 Yafei Yuan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Acting as a brain stimulant, coffee resulted in heightening alertness, keeping arousal, improving executive speed, maintaining vigilance, and promoting memory, which are associated with attention, mood, and cognitive function. Functional near-infrared spectroscopy (fNIRS) is a noninvasive optical method to monitor brain activity by measuring the absorption of the near-infrared light through the intact skull. This study is aimed at acquiring brain activation during executing task performance. The aim is to explore the effect of coffee on cognitive function by the fNIRS neuroimaging method, particularly on the prefrontal cortex regions. The behavioral experimental results on 31 healthy subjects with a Stroop task indicate that coffee can easily and effectively modulate the execute task performance by feedback information of the response time and accuracy rate. The findings of fNIRS showed that apparent hemodynamic changes were detected in the bilateral VLPFC regions and the brain activation regions varied with different coffee conditions.

1. Introduction

Caffeine, primarily obtained from coffee in daily life, is the most widely and frequently used psychoactive drug over worldwide [1–4]. Coffee is a highly-concentrated source of caffeine about 2% [5]. Coffee consumption is associated with some health benefits including reduced risk of cardiovascular disease (CVD) [6], lower incidence of diabetes mellitus [7, 8], less Alzheimer's disease (AD) [9, 10], and decreased death from inflammatory diseases [11, 12]. Acting as a brain stimulant, coffee resulted in heightening alertness, keeping arousal, improving executive speed, maintaining vigilance, and promoting memory, which are associated with attention, mood, and cognitive function [13–15]. The most profound function is improving cognition which involves human activities of daily living. The benefits of cognition from coffee consumption are typically correlated with caffeine that exerts its effects through nonselective antagonism of adenosine A_1 receptors (A_1R) and A_{2A} receptors (A_{2AR}) [10, 16]. Studies indicate that caffeine absorption is from the gastrointestinal tract rapid and reaches 99% after about 45 min ingestion [17]. Due to hydrophobic properties, the caffeine allows its

passage through all biological membranes, including the blood-brain barrier. Peak plasma caffeine concentrations appeared between 15 and 120 min after oral ingestion [18]. Moreover, caffeine half-life ranges from 3 to 5 hours in humans [19]. The majority of the research studies have focused on cognitive function and generally showed consistent findings [20]. However, with respect to studies on the higher-order complex brain function which relies on frontal lobe function, there is no consistent result. Such functions include the ability to plan, conceptualize, initiate, and sustain behavioral with aims [1]. The cognitive function constructs which are known as executive functioning are sensitive to the ingestion of caffeine. Hence, knowing the underlying interaction between caffeine and human cognition is valuable for human health and production.

The functional neuroimaging method has been more inclined to be adapted in cognitive neuroscience [21, 22], which explores the constituent parts of the human cognitive system might be and how the human cognition works with other mental processes [23]. As a new functional neuroimaging technique, the functional near-infrared spectroscopy (fNIRS) is the noninvasive optical method to monitor brain

activity by measuring the absorption of the near-infrared light between 650 nm and 950 nm through the intact skull [24]. What is more, the absorption spectra of oxygenated hemoglobin (Oxy-Hb) and deoxygenated hemoglobin (Deoxy-Hb) are distinct in this region [25]. The concentrations of Oxy-Hb and Deoxy-Hb can be calculated separately using the modified Beer–Lambert law [26, 27]. Thus, Jöbsis first proposed the application of near-infrared spectroscopy (NIRS) in vivo to acquire the concentration changes of Oxy-Hb and Deoxy-Hb by the diffusely scattered light measurements [28].

Compared with other neuroimaging technique as positron emission tomography (PET), functional magnetic resonance imaging (fMRI), and magnetoencephalography (MEG), the fNIRS has many advantages [29]. The main advantage is the ability to directly measure the functional contrasts such as Oxy-Hb, Deoxy-Hb, and total hemoglobin with very high temporal resolution which enables the studies on the hemodynamic response to neural activation [24]. fNIRS is an inexpensive, relatively portable, and tolerant of motion artifacts neuroimaging method compare with other measurements which is completely noninvasive and could be used in daily life [22, 30]. The fNIRS is widely employed to investigate human cognition [31–33]. The frontal areas of the brain, specifically the prefrontal cortex (PFC), are associated with cognition [34–36]. The PFC regions carry out executive functions including higher-order cognitive functions which are essential for planning and executing complex motor control actions [37–39]. Hence, in order to investigate the underlying mechanism relation between light and cognition, selecting the PFC as the region of interest (ROI) is a reasonable and effective research route method [40]. Furthermore, studies about how the coffee modulates the cognition is seldom reported which is important for human health.

In this work, the aim is to explore the effect of coffee on cognitive function by fNIRS neuroimaging method, particularly on PFC regions. As focusing on how the coffee affects the cognitive function, 31 participants were arranged to ingest coffee and carried on the Stroop task before and after coffee ingestion. Moreover, the variation in ROIs with modulation by coffee is inspected by fNIRS.

2. Materials and Methods

2.1. Participants. Thirty-one right-handed healthy participants (15 males and 16 females) were selected for the study purpose from the age group of 21 to 24 years. All participants had normal or corrected-to-normal vision and normal color vision. No participants had a history of neurological or psychiatric disorder and were under any psychotropic medication. The research experiment strictly followed the ethics regulation of Fudan University and obtained the consent of each participant.

2.2. Procedures and Materials. All participants performed three Stroop tasks under three different light conditions. The modified color-word matching Stroop task was used as an event-related cognition task in this research [41, 42].

The functional neuroimaging method used the Stroop task for exploring human cognition [38, 42–44]. The rule of the single trial for the congruent, incongruent, and neutral conditions of color-word matching Stroop task is depicted in Figure 1(a). The single trial consisted of 12 words that were printed in black ink and appeared randomly. The complete task included 11 trials performed in one light mode. Participants were instructed to press the relevant response key during the executive task by the designed rules. The correct response action was to press the number keys “1,” “2,” and “3” for red, green, and blue color words, respectively. The response time and the response were recorded using a computer for further analysis.

The coffee effect task procedure is designed with two Stroop cycles as shown in Figure 1(b). Participants first performed the practice experiment and then carried out an actual procedure with over 92% practice accuracy rate. Participants intake a cup of Nescafe coffee about 210 ml after normal/Stroop task. The task process time is arranged by rest 4 minutes → normal/Stroop task → rest 20 minutes → coffee/Stroop task → end.

2.3. Data Acquisition by fNIRS. The fNIRS data were recorded using a multichannel continuous-wave fNIRS system instrument (NIRx Medical Technologies LLC-NIR-Scout, USA), which consists of eight LED light sources and eight photodetectors [38]. The distance between the detector and source is approximately 3 cm. The detector record relative changes in Oxy-Hb and Deoxy-Hb at a sample rate of 7.81 Hz at two wavelengths (760 and 850 nm) [45]. The location of probe placement and arrangement of specific brain regions was according to previous studies [46]. In this study, we focused on the PFC region for investigating cognition. Figure 1(c) shows the set-up of fNIRS channels. Four different ROIs were selected: the right ventrolateral prefrontal cortex (R-VLPFC) (channels 17-20), the right dorsolateral prefrontal cortex (R-DLPFC) (channels 13-16), the left L-DLPFC (channels 5-7 and 11), and the posterior left L-VLPFC (channels 1-4). The fNIRS transmitters were specially designed to wrap with a tightly black bandage to ensure that there was no extraneous light interference during the Stroop cognitive task.

Participants were instructed to sit comfortably in a chair and maintain a calm and relaxed position. They were asked to focus on the screen with their minds blank. The visual of the task was presented on a 21-inch thin-film transistor (TFT) screen.

2.4. Data Processing. The fNIRS raw data analysis was executed based on SPM with additional modules for ANOVAs. First, a low-frequency band-pass filter (0.01-0.2 Hz) was applied to eliminate the baseline drift, artifact, and physiological noise. fNIRS records the changes in Oxy-Hb and Deoxy-Hb concentration simultaneously. However, there are some scientific problems in the selection of signals to analyze brain activation. In this research, it mainly focused on the Oxy-Hb signal changes, as it was normally observed to own higher amplitude than the Deoxy-Hb signal [45]. Furthermore, the signal-to-noise




FIGURE 1: Task procedures and materials. (a) The color-word matching Stroop task rule. (b) The task process time. (c) The setting up of fNIRS channels.

TABLE 1: Stroop task performance results.

	Response time (ms)			Accuracy (%)		
	Congruent	Incongruent	Neutral	Congruent	Incongruent	Neutral
Normal	541.5 ± 76.9	601.4 ± 103.0	553.9 ± 84.0	97.9 ± 3.1	97.2 ± 2.0	97.4 ± 3.2
Coffee	523.3 ± 66.0	566.5 ± 89.2	536.7 ± 84.1	98.0 ± 2.4	97.2 ± 2.4	98.0 ± 2.5

(S/N) ratio of Oxy-Hb is better, and the signal is more sensitive to process task response [47]. To explore whether those channels activated during the Stroop task, we analyzed channel-by-channel. If the light intensity of channels fell below 400 mV or exceeded 4,000 mV at any point during the session, then they were rejected [35]. The behavioral performance and Oxy-Hb data were analyzed in Vision 22.0 SPSS using a 3 (stimulus conditions) × 2 (coffee condition) repeated-measure ANOVAs.

3. Results

3.1. Behavioral Results: Stroop Interference. The results of Stroop task performance are shown in Table 1, including the response time and accuracy rate. Figure 2 depicts the behavioral response time (RT) and accuracy rate results. Concerning

the response time shown in Figure 2(a), by using the repeated-measure stimulus condition (congruent vs. incongruent vs. neutral) × coffee condition (normal vs. coffee), ANOVA demonstrated a significant effect for stimulus condition ($F = 28.81$, $df = 2$, $P < 0.0005$) and coffee conditions ($F = 10.25$, $df = 1$, $P = 0.003$). No significant interaction effect of the stimulus condition × coffee condition was observed ($F = 2.13$, $df = 2$, $P = 0.128$). Three stimulus condition under normal state ($F = 26.52$, $df = 2$, $P < 0.0005$) and coffee state ($F = 13.76$, $df = 2$, $P < 0.0005$) showed an obvious Stroop interference. Further, with regard to the coffee condition with different stimulus condition, the congruent ($F = 5.10$, $df = 1$, $P = 0.031$) and incongruent ($F = 13.82$, $df = 1$, $P = 0.001$) show significant difference, and neutral ($F = 2.93$, $df = 1$, $P = 0.097$) shows marginal significant difference. The RT of the incongruent stimulus condition was longer than the




FIGURE 2: The results of Stroop task performance. (a) Response time with two conditions. (b) Accuracy rate with two conditions.




FIGURE 3: The brain activation under different conditions. (a) The brain activation under normal/rest condition. (b) The brain activation under normal/Stroop condition. (c) The brain activation under coffee/rest condition. (d) The brain activation under coffee/Stroop condition.

congruent and neutral stimulus condition in all normal and coffee states. Participants responded quickly after coffee ingestion condition than normal condition.

With respect to the accuracy rate shown in Figure 2(b), by using the repeated-measure stimulus condition (congruent vs .incongruent vs .neutral) × coffee condition (normal vs .coffee) , ANOVA results demonstrated a marginally significant effect for stimulus condition ($F = 2.45$, $df = 2$, $P = 0.095$) and no significant effect for coffee condition. No main significant interaction effect of the stimulus condition × coffee conditions was observed. It is clear that the accuracy rate is higher under the coffee ingestion condition compared with normal condition in Figure 2(b). Furthermore, the accuracy rates were significantly lower in the incongruent condition compared with the congruent and neutral ones. As the behavioral results, it is clearly shown that the participants exhibit the best behavioral performance under the coffee ingestion condi-




FIGURE 4: The significant difference in Oxy-Hb levels of the channels compared. (a) The comparison of Oxy-Hb levels with coffee/rest vs. normal/rest. (b) The comparison of Oxy-Hb levels with coffee/Stroop vs. normal/Stroop.




FIGURE 5: Mean Oxy-Hb concentration in PFC regions. (a) Mean Oxy-Hb concentration in PFC regions under normal/rest condition. (b) Mean Oxy-Hb concentration in PFC regions under normal/Stroop condition. (c) Mean Oxy-Hb concentration in PFC regions under coffee/rest condition. (d) Mean Oxy-Hb concentration in PFC regions under coffee/Stroop condition.




FIGURE 6: The results of the coffee effect on behavioral performance.

tion than normal condition, not only due to faster response time but also more accuracy rate. In this research, coffee ingestion is regarded as the high-efficiency method to improve human performance.

3.2. *fNIRS Results: Brain Activation.* The brain activation of the *t*-statistic map for Oxy-Hb is shown in Figure 3 under four conditions which were normal/rest, normal/Stroop, coffee/rest, and coffee/Stroop. Figure 3(a) under normal/rest condition showed that there were no channels with brain activation of the Oxy-Hb level. Figure 3(b) under normal/Stroop condition showed brain activation of Oxy-Hb level in several channels (1, 2, 5, and 8) mainly locating at L-VLPFC. Figure 3(c) under coffee/rest condition showed brain activation of Oxy-Hb level in several channels (4, 6, 11, 12, and 13) mainly locating at L-VLPFC and L-DLPFC. Figure 3(d) under the coffee/Stroop condition showed brain activation of Oxy-Hb level in several channels (1, 2, 3, 4, 5, 7, 9, 10, 15, and 18) mainly locating at VLPFC. It is clear to see that the significantly different exhibited brain activation in PFC under normal vs. coffee conditions. The lowest mean Oxy-Hb level happened in normal/rest condition. While in normal/Stroop condition, it is required that a high mean Oxy-Hb level completes the task resulting in activating the channels (1, 2, 5, and 8). Furthermore, channel 12 was significantly activated under coffee/rest condition demonstrating the strong correlation between coffee effects in L-DLPFC. Moreover, compared with the Stroop task modes, it is found that the relevant ROIs change following the coffee intake.




FIGURE 7: The brain activation under coffee/rest condition. (a) The brain activation after coffee ingestion 10 min. (b) The brain activation after coffee ingestion 15 min. (c) The brain activation after coffee ingestion 20 min.

Compared to the normal/Stroop condition, more channels were involved in completing tasks under coffee/Stroop mainly located in R-VLPFC and L-VLPFC.

In order to locate the exact channel variates, we used a paired *t*-test to statistically compare the PFC activation under the coffee condition. The results of all channels with significant differences ($P < 0.05$) are shown in Figure 4. Figure 4(a) shows that the channels (4, 13, and 19) present significant differences between the normal/rest and normal/Stroop conditions. And Figure 4(b) shows that the channels (4 and 13) present significant differences between coffee/rest and coffee/Stroop conditions. In order to investigate the variate and hemodynamic response accompanying with getting worse light condition, we focus on the mean Oxy-Hb level of these ROIs such as L-VLPFC, L-DLPFC, R-VLPFC, and R-DLPFC. Figure 5 illustrates the mean Oxy-Hb concentrations in PFC regions under four conditions. Significant differences were observed in normal and coffee conditions. It is clear to see the apparent changes in ROIs between the different coffee conditions. The mean Oxy-Hb concentration in L-VLPFC, L-DLPFC, R-VLPFC, and R-DLPFC increases with coffee intake. Under rest conditions, the coffee intake makes Oxy-Hb concentration of the L-DLPFC and R-VLPFC regions significantly rising. Under Stroop task conditions, the coffee intake makes Oxy-Hb concentration of the L-VLPFC, L-DLPFC, and R-DLPFC regions significantly rising.

4. Discussion

The present study is aimed at investigating how coffee intake on the cognitive function and PFC brain activation effect during the color-word matching Stroop task. In this research, the Stroop interference results existed in normal and coffee conditions were in accordance with the literature [41]. The results indicated that coffee condition could improve the participant's behavioral performance during executive tasks while comparing the normal condition which is consistent with work [20]. The findings were concluded not only based on the response time results but also the accuracy rate results. Participants under the coffee/Stroop condition perform faster executive speed with a higher accuracy rate compared to under normal/Stroop. That is to say, people could keep efficient working by coffee intake condition. As shown in

Figure 6, it depicted the coffee effect on behavioral performance by using coffee/Stroop condition subtracting normal/Stroop concerning to response time. Clearly, the coffee condition is higher than the normal condition in all kinds of stimulus condition, which have more facilitative effects on the executive functions resulting in fast response and high accuracy. Specially, coffee could improve more performance in the complex task (incongruent) compared to easy task (congruent and neutral). Interestingly, there is some significant difference between males and females on the coffee effect under all stimuli. The coffee takes a stronger effect on behavioral performance with males compared to females, which proves males are more sensitive to coffee.

Additionally, the neuroimaging results by fNIRS demonstrated the Oxy-Hb levels in response to the Stroop task in the bilateral VLPFC and DLPFC. The apparent hemodynamic changes were detected in the bilateral VLPFC regions under different conditions which are consistent with previous studies on cognition by event-related neuroimaging [38, 43, 44]. Under the efficiency work light condition related with coffee ingestion, the VLPFC was activated by mainly increasing several channel Oxy-Hb levels (1, 2, 3, 4, 7, 10, 15, and 18) mainly located in L-VLPFC. The VLPFC started to be activated due to coffee ingestion to execute tasks. Previous studies concluded that the activation of the right DLPFC may be due to the sensitivity in task difficulty [48]. A similar phenomenon was observed in our study, as the Oxy-Hb level of DLPFC increases under the coffee/Stroop condition shown in Figure 6(d). In order to explore the process of coffee effect, Figure 7 shows the process of coffee ingestion within 20 min, which clearly describes the process of variations in PFC. The mean Oxy-Hb level of channels (4 and 6) increase after coffee ingestion.

According to the behavioral performance and fNIRS results, it was concluded that coffee intake could effectively improve human cognitive function. It was proposed that humans show better performance with accomplishing high-efficiency work within coffee condition. This research data demonstrates that coffee intake condition makes people efficiently and easily undergoes the mission. In summary, this study confirmed that coffee can be regarded as a safe, effective, less expensive, and accessible tool to modulate human cognitive function. Additionally, the fNIRS is a reliable method to explore the mechanisms of brain activation for cognitive performance in relation to relevant factors.

5. Conclusion

In conclusion, the present observations confirm that the fNIRS is a noninvasive, less expensive, and accessible neuroimaging method. It is well-suited for monitoring PFC changes associated with cognitive functions. This study has focused on the effect of coffee intake on human cognition and explored the underlying mechanism. The behavioral results indicate that coffee can easily and effectively modulate the execution task performance by the means of feedback information, such as response time and accuracy. The findings of fNIRS show that the apparent hemodynamic changes were detected in the bilateral VLPFC regions, and the brain activation regions variate with different coffee condition. Furthermore, it reveals that coffee is a sensitive modulator for cortical hemodynamic correlated with cognitive function. By combining the two results, human behavioral performance benefit from coffee intake and show high-efficiency work. Moreover, coffee could be regarded as a safe, effective, inexpensive, and accessible method to modulate human cognitive function, which is of significant importance for human daily life.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We acknowledge the financial supports by the National Key R&D Program of China (Grant No. 2017YFE0112000) and the Shanghai Municipal Science and Technology Major Project (Grant No. 2017SHZDZX01). Furthermore, the authors wish to thank professor H. Du from China academy of space technology. Furthermore, the authors would like to express sincere thanks for YC. Yuan.

References

- [1] K. Soar, E. Chapman, N. Lavan, A. S. Jansari, and J. J. Turner, "Investigating the effects of caffeine on executive functions using traditional Stroop and a new ecologically-valid virtual reality task, the Jansari assessment of Executive Functions (JEF ©)," *Appetite*, vol. 105, pp. 156–163, 2016.
- [2] S. Ullrich, Y. C. de Vries, S. Kuhn, D. Repantis, M. Dresler, and K. Ohla, "Feeling smart: effects of caffeine and glucose on cognition, mood and self-judgment," *Physiology & Behavior*, vol. 151, pp. 629–637, 2015.
- [3] M. Rivera-Oliver and M. Diaz-Rios, "Using caffeine and other adenosine receptor antagonists and agonists as therapeutic tools against neurodegenerative diseases: a review," *Life Sciences*, vol. 101, no. 1–2, pp. 1–9, 2014.
- [4] A. Zhou, A. E. Taylor, V. Karhunen et al., "Habitual coffee consumption and cognitive function: a Mendelian randomization meta-analysis in up to 415,530 participants," *Scientific Reports*, vol. 8, no. 1, p. 7526, 2018.
- [5] D. D. Church, J. R. Hoffman, M. B. LaMonica et al., "The effect of an acute ingestion of Turkish coffee on reaction time and time trial performance," *Journal of the International Society of Sports Nutrition*, vol. 12, no. 1, p. 37, 2015.
- [6] F. Rodriguez-Artalejo and E. Lopez-Garcia, "Coffee consumption and cardiovascular disease: a condensed review of epidemiological evidence and mechanisms," *Journal of Agricultural and Food Chemistry*, vol. 66, no. 21, pp. 5257–5263, 2018.
- [7] X. Jiang, D. Zhang, and W. Jiang, "Coffee and caffeine intake and incidence of type 2 diabetes mellitus: a meta-analysis of prospective studies," *European Journal of Nutrition*, vol. 53, no. 1, pp. 25–38, 2014.
- [8] R. K. West, R. Ravona-Springer, A. Livny et al., "Age modulates the association of caffeine intake with cognition and with gray matter in elderly diabetics," *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, vol. 74, no. 5, pp. 683–688, 2019.
- [9] C. F. Haskell-Ramsay, P. A. Jackson, J. S. Forster, F. L. Dodd, S. L. Bowerbank, and D. O. Kennedy, "The acute effects of caffeinated black coffee on cognition and mood in healthy young and older adults," *Nutrients*, vol. 10, no. 10, p. 1386, 2018.
- [10] A. Rosso, J. Mossey, and C. F. Lippa, "Review: caffeine: neuroprotective functions in cognition and Alzheimer's disease," *American Journal of Alzheimer's Disease & Other Dementias*, vol. 23, no. 5, pp. 417–422, 2008.
- [11] L. F. Andersen, D. R. Jacobs Jr., M. H. Carlsen, and R. Blomhoff, "Consumption of coffee is associated with reduced risk of death attributed to inflammatory and cardiovascular diseases in the Iowa Women's Health Study," *The American Journal of Clinical Nutrition*, vol. 83, no. 5, pp. 1039–1046, 2006.
- [12] J. A. Greenberg, C. C. Dunbar, R. Schnoll, R. Kokolis, S. Kokolis, and J. Kassotis, "Caffeinated beverage intake and the risk of heart disease mortality in the elderly: a prospective analysis," *The American Journal of Clinical Nutrition*, vol. 85, no. 2, pp. 392–398, 2007.
- [13] S. Haller, M. L. Montandon, C. Rodriguez, F. R. Herrmann, and P. Giannakopoulos, "Impact of coffee, wine, and chocolate consumption on cognitive outcome and MRI parameters in old age," *Nutrients*, vol. 10, no. 10, p. 1391, 2018.
- [14] C. F. Haskell, D. O. Kennedy, K. A. Wesnes, and A. B. Scholey, "Cognitive and mood improvements of caffeine in habitual consumers and habitual non-consumers of caffeine," *Psychopharmacology*, vol. 179, no. 4, pp. 813–825, 2005.
- [15] T. Garcia-Blanco, A. Davalos, and F. Visioli, "Tea, cocoa, coffee, and affective disorders: vicious or virtuous cycle?," *Journal of Affective Disorders*, vol. 224, pp. 61–68, 2017.
- [16] A. Kerkhofs, A. C. Xavier, B. S. da Silva et al., "Caffeine controls glutamatergic synaptic transmission and pyramidal neuron excitability in human neocortex," *Frontiers in Pharmacology*, vol. 8, p. 899, 2017.
- [17] A. Nehlig, "Is caffeine a cognitive enhancer?," *Journal of Alzheimer's disease : JAD*, vol. 20, pp. 85–94, 2010.
- [18] B. B. Fredholm, K. Bättig, J. Holmén, A. Nehlig, and E. E. Zvartau, "Actions of caffeine in the brain with special reference to factors that contribute to its widespread use," *Pharmacological Reviews*, vol. 51, no. 1, pp. 83–133, 1999.
- [19] P. J. Rogers, "Caffeine, mood and mental performance in everyday life," *Nutrition Bulletin*, vol. 32, no. s1, pp. 84–89, 2007.

- [20] A. Shabir, A. Hooton, J. Tallis, and M. F. Higgins, "The influence of caffeine expectancies on sport, exercise, and cognitive performance," *Nutrients*, vol. 10, pp. 1–11, 2018.
- [21] J. P. Mitchell, "Contributions of functional neuroimaging to the study of social cognition," *Current Directions in Psychological Science*, vol. 17, no. 2, pp. 142–146, 2008.
- [22] M. Coltheart, "How can functional neuroimaging inform cognitive theories?," *Perspectives on Psychological Science*, vol. 8, no. 1, pp. 98–103, 2013.
- [23] H. Fujiwara, W. Yassin, and T. Murai, "Neuroimaging studies of social cognition in schizophrenia," *Psychiatry and Clinical Neurosciences*, vol. 69, no. 5, pp. 259–267, 2015.
- [24] J. YE, S. TAK, K. JANG, J. JUNG, and J. JANG, "NIRS-SPM: statistical parametric mapping for near-infrared spectroscopy," *NeuroImage*, vol. 44, no. 2, pp. 428–447, 2009.
- [25] S. Lloyd-Fox, A. Blasi, and C. E. Elwell, "Illuminating the developing brain: the past, present and future of functional near infrared spectroscopy," *Neuroscience and Biobehavioral Reviews*, vol. 34, no. 3, pp. 269–284, 2010.
- [26] V. Quaresima and M. Ferrari, "A mini-review on functional near-infrared spectroscopy (fNIRS): where do we stand, and where should we go?," *Photonics*, vol. 6, no. 3, p. 87, 2019.
- [27] N. K. Schaal, P. Hepp, A. Schweda, O. T. Wolf, and C. Krampe, "A functional near-infrared spectroscopy study on the cortical haemodynamic responses during the Maastricht acute stress test," *Scientific Reports*, vol. 9, no. 1, article 13459, 2019.
- [28] F. F. Jöbsis, "Noninvasive infrared monitoring of cerebral and myocardial oxygen sufficiency and circulatory parameters," *Science*, vol. 198, no. 4323, pp. 1264–1267, 1977.
- [29] S. Perrey, "Non-invasive NIR spectroscopy of human brain function during exercise," *Methods*, vol. 45, no. 4, pp. 289–299, 2008.
- [30] Y. Hoshi, "Functional near-infrared spectroscopy: current status and future prospects," *Journal of Biomedical Optics*, vol. 12, no. 6, article 062106, 2007.
- [31] A. Curtin, H. Ayaz, Y. Tang, J. Sun, J. Wang, and S. Tong, "Enhancing neural efficiency of cognitive processing speed via training and neurostimulation: an fNIRS and TMS study," *NeuroImage*, vol. 198, pp. 73–82, 2019.
- [32] Z. Hu, K. F. Lam, and Z. Yuan, "Effective connectivity of the fronto-parietal network during the tangram task in a natural environment," *Neuroscience*, vol. 422, pp. 202–211, 2019.
- [33] E. Holmes, D. W. Barrett, C. L. Saucedo, P. O'Connor, H. Liu, and F. Gonzalez-Lima, "Cognitive enhancement by transcranial photobiomodulation is associated with cerebrovascular oxygenation of the prefrontal cortex," *Frontiers in Neuroscience*, vol. 13, p. 1129, 2019.
- [34] S. Y. Xu, F. M. Lu, M. Y. Wang et al., "Altered functional connectivity in the motor and prefrontal cortex for children with Down's syndrome: an fNIRS study," *Frontiers in Human Neuroscience*, vol. 14, p. 6, 2020.
- [35] J. Rovetti, H. Goy, M. K. Pichora-Fuller, and F. A. Russo, "Functional near-infrared spectroscopy as a measure of listening effort in older adults who use hearing aids," *Trends in Hearing*, vol. 23, pp. 1–22, 2019.
- [36] T. Moriarty, K. Bourbeau, B. Bellovary, and M. N. Zuhl, "Exercise intensity influences prefrontal cortex oxygenation during cognitive testing," *Behavioral Sciences*, vol. 9, no. 8, p. 83, 2019.
- [37] C. Udina, S. Avtzi, T. Durduran et al., "Functional near-infrared spectroscopy to study cerebral hemodynamics in older adults during cognitive and motor tasks: a review," *Frontiers in Aging Neuroscience*, vol. 11, p. 367, 2019.
- [38] Z. Ji, T. Feng, L. Mei, A. Li, and C. Zhang, "Influence of acute combined physical and cognitive exercise on cognitive function: an NIRS study," *PeerJ*, vol. 7, article e7418, 2019.
- [39] M. L. Schroeter, S. Zysset, M. Wahl, and D. Y. von Cramon, "Prefrontal activation due to Stroop interference increases during development—an event-related fNIRS study," *NeuroImage*, vol. 23, no. 4, pp. 1317–1325, 2004.
- [40] D. Yang, K. S. Hong, S. H. Yoo, and C. S. Kim, "Evaluation of neural degeneration biomarkers in the prefrontal cortex for early identification of patients with mild cognitive impairment: an fNIRS study," *Frontiers in Human Neuroscience*, vol. 13, p. 317, 2019.
- [41] C. M. MacLeod and P. A. MacDonald, "Interdimensional interference in the Stroop effect: uncovering the cognitive and neural anatomy of attention," *Trends in Cognitive Science*, vol. 4, no. 10, pp. 383–391, 2000.
- [42] H. Yanagisawa, I. Dan, D. Tsuzuki et al., "Acute moderate exercise elicits increased dorsolateral prefrontal activation and improves cognitive performance with Stroop test," *NeuroImage*, vol. 50, no. 4, pp. 1702–1710, 2010.
- [43] J. León-Carrion, J. Damas-López, J. F. Martín-Rodríguez et al., "The hemodynamics of cognitive control: the level of concentration of oxygenated hemoglobin in the superior prefrontal cortex varies as a function of performance in a modified Stroop task," *Behavioural Brain Research*, vol. 193, no. 2, pp. 248–256, 2008.
- [44] K. Byun, K. Hyodo, K. Suwabe et al., "Positive effect of acute mild exercise on executive function via arousal-related prefrontal activations: an fNIRS study," *NeuroImage*, vol. 98, pp. 336–345, 2014.
- [45] Z. Xu, M. Hu, Z. R. Wang, J. Li, X. H. Hou, and M. Q. Xiang, "The positive effect of moderate-intensity exercise on the mirror neuron system: an fNIRS study," *Frontiers in Psychology*, vol. 10, p. 986, 2019.
- [46] D. Tsuzuki, V. Jurcak, A. K. Singh, M. Okamoto, E. Watanabe, and I. Dan, "Virtual spatial registration of stand-alone fNIRS data to MNI space," *NeuroImage*, vol. 34, no. 4, pp. 1506–1518, 2007.
- [47] X. Cheng, X. Li, and Y. Hu, "Synchronous brain activity during cooperative exchange depends on gender of partner: a fNIRS-based hyperscanning study," *Human Brain Mapping*, vol. 36, pp. 2039–2048, 2015.
- [48] S. D. Newman, P. A. Carpenter, S. Varma, and M. A. Just, "Frontal and parietal participation in problem solving in the Tower of London: fMRI and computational modeling of planning and high-level perception," *Neuropsychologia*, vol. 41, no. 12, pp. 1668–1682, 2003.