








Association of Allergic Sensitivity and Pollination in Allergic Respiratory Disease: The Role of Pollution

Gandhi Fernando Pavón-Romero ¹, María del Carmen Calderón-Ezquerro ²,
Michelle Alejandra Rodríguez-Cervantes ³, David Fernández-Villanueva ¹, Esmeralda Melgoza-Ruiz ¹,
Fernando Ramírez-Jiménez ¹, Luis M Teran ¹

¹Department of Immunogenetics and Allergy, Instituto Nacional de Enfermedades Respiratorias “Ismael Cosío Villegas”, Mexico City, Mexico; ²Grupo Bioaerosoles atmosféricos, Instituto de Ciencias de la Atmósfera y Cambio Climático, UNAM, Mexico City, Mexico; ³Department of Pediatrics, Instituto Nacional de Pediatría, Mexico City, Mexico

Correspondence: Luis M Teran, Department of Immunogenetics and Allergy, Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas, Mexico City, 14080, Mexico, Tel/Fax +52 5557481740, Email teranjlm@gmail.com

Purpose: To evaluate the association between allergic sensitivity and pollen counts in patients with allergic respiratory disease (ARD) and its relationship with atmospheric pollutants.

Methods: From 2012 to 2018, we evaluated the sensitivity by skin prick test in ARD patients. The pollen counts were analyzed according to international guidelines (2014–2018). The pollutant and meteorological data were obtained at the same time from AIRE-CDMX websites. We analyzed the association between allergic sensitivity and pollen counts using the χ^2 test and stratified by disease allergic rhinitis (AR) and AR with asthma (ARwA), periods (before/after 2015), and pollination seasons (S1:2014–2015), (S2:2015–2016), (S3:2016–2017), (S4:2017–2018). Likewise, we correlated the pollen counts with the concentrations of pollutants using Pearson’s correlation. For all analyses, we used SPSS v.21 software, and a p -value <0.05 was considered significant.

Results: A total of 520 patients were enrolled, of whom 67.3% had ARwA and 33.7% had AR ($p<0.05$). The frequency of patients allergic to at least one pollen was higher compared with patients sensitive to indoor allergens (55.3% vs 44.6%, $p<0.001$). A total of 46.8% of the patients were only sensitive to trees in comparison to other outdoor allergens ($p<0.001$). The *Fraxinus* sp. and the Cupressaceae family allergens were approximately two times more frequent than the other tree allergens in both diseases ($p<0.05$). These pollens doubled their counts since 2015 ($p<0.001$), which was associated with increases in sensitivity for *Fraxinus* sp. and the Cupressaceae family compared to previous years ($p<0.001$). Regarding pollutants, the most significant correlations were with PM₁₀, NO₂, PM_{CO} for *Fraxinus* sp. pollen concentrations in all seasons ($p\leq 0.02$).

Conclusion: The high increases in pollen counts of the *Fraxinus* sp. and Cupressaceae family were associated with increases in the frequency of sensitization to these species, and this phenomenon correlated with increases in PM₁₀, NO₂, and PM_{CO}.

Keywords: air pollutants, airborne particulate matter, pollen, allergy, asthma, rhinitis allergic seasonal

Introduction

The prevalence and incidence of allergic respiratory diseases (ARDs), such as allergic rhinitis (AR) and asthma,¹ are increasing worldwide. AR affects approximately 40% of the world’s population² and is considered a risk factor for the development of allergic asthma (ARwA),³ with this being the main phenotype (~80%).⁴ It is estimated that 330 million suffer from asthma. Both diseases are health problems due to their impact on the quality of life (sleep quality, outdoor activities, education, occupational performance, etc.).^{2,5} House dust mites (HDMs) are the major indoor allergen,⁶ while pollen is the predominant cause of outdoor respiratory allergies.⁷ Pollen monitoring provides relevant information about the role of this organic pollutant in ARD patients. For example, the description of specific periods of the year when there is an increase or decrease in pollen counts seated in a specific city or region could help the physician to predict the increase in the severity of ARD patients susceptible to pollen and therefore would aid in the prescription of preventive or therapeutic measures for improving the clinical symptoms.⁸ Moreover, there is an increase in the prevalence of ARD




Figure 1 The interaction of pollutants such as $PM_{2.5}$ and O_3 with pollen induces: (A) an increase in the production of pollen concentrations, (B) a fragile pollen membrane, and (C) post-translational modifications in the allergenic protein. In all cases, there are increased levels of allergen proteins, which can be processed by antigen-presenting cells, resulting in a type I hypersensitivity inflammation (mechanism mediated by IL-4, IL-5, and IL-13, and an increase in immunoglobulin E), whose main objective is the degranulation of preformed mediators (histamine or tryptase) contained in mast cells, which promotes mucus secretion by goblet cells and bronchospasm, main symptom of asthma. **Abbreviations:** $PM_{2.5}$, particulate matter of 2.5 micrometers or less; O_3 , ozone; IL, interleukin.

worldwide, which is related to changes in the concentrations of environmental pollutants, specifically to the increases in pollen counts of various species.⁹

It is well established that pollution facilitates the development of allergic respiratory diseases. Nitrogen dioxide (NO_2), ozone (O_3), and particulate matter (PM_{10} or $PM_{2.5}$) have been associated with asthma exacerbations.¹⁰ This observation is supported by a recent report from the American Thoracic Society Workshop Report showing that air pollution is associated with the onset of both asthma and an increased risk of allergic sensitization.¹¹ In this context, some pollutants such as $PM_{2.5}$ and O_3 increases pollen counts, leading to a higher concentration of allergenic proteins in the environment.¹² Additionally, $PM_{2.5}$ can induce greater fragility in the exine, causing a collapse of the pollen membrane and a subsequent release of epitopes.¹³ These proteins can be up taken and processed by antigen-presenting cells in the respiratory tract mucosa, initiating the type I hypersensitivity-mediated inflammation, whose main objective is the release of preformed mediators (histamine) from mast cells by immunoglobulin E, increasing mucus secretion, vascular permeability and bronchospasm¹⁴ (Figure 1). In the present study, we investigated the association of the allergic sensitivity of ARD patients with the presence of both outdoor aeroallergens and pollution in Mexico City.

Materials and Methods

Geographic Area

Mexico City is the smallest state in Mexico; it represents 7.3% of the national territory and has 9,209,944 inhabitants. It comprises 16 municipalities, Tlalpan being the greatest, with an area of 306.52 km², and is located at 19° 09' 57" north latitude and 99° 09' 57" west longitude (<http://www.inafed.gob.mx/work/enciclopedia/EMM09DF/delegaciones/09012a.html>). This demarcation is the fourth most populated (~700,000 inhabitants) in Mexico City (<http://cuentame.inegi.org.mx/monografias/informacion/df/poblacion/>). Most of the year, it has a subhumid climate (87%), rain occurs in summer, the total annual precipitation can reach up to 1200 mm per year, and its temperature ranges between 5 °C and 25 °C. (<http://www.cuentame.inegi.org.mx/monografias/informacion/df/territorio/clima.aspx?tema=meande=09>). This region is the location of the *Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas*. On its roof (15 m high), a Hirst-type volumetric spore trap (Burkard Manufacturing Co., Ltd., Rickmansworth, UK) is installed.

Subjects of Study

Ethics Statement

All participants provided written informed consent. This study was performed in accordance with the principles stated in the Declaration of Helsinki, and the study protocol was approved by the science and bioethics committee for research of the Instituto Nacional de Enfermedades Respiratorias “Ismael Cosío Villegas” (Approval number: E03-13), before the start of the study (February 20th, 2013).

Patients

The admission criteria to select medical records included: a) the clinical history had to refer to whether the patients lived and carried out their daily activities most of the time in the south of Mexico City, b) AR was diagnosed according to Allergic Rhinitis and its Impact on Asthma-ARIA 2008 guidelines (clinical antecedent of sneezing, itchiness, nasal congestion, and rhinorrhea and evidence of allergic sensitivity by any diagnostic method as skin prick test),¹⁵ c) the asthma was diagnosed according to Global Initiative for Asthma-GINA 2012 guidelines,⁵ the presence of suggestive lung symptoms (cough, wheezing, chest tightness, dyspnea) plus reversibility test (an increase of Forced expiratory volume in the first second) by spirometry; being this criterion only apply to the asthma group. From 3520 medical records from AR or ARwA patients attended in our department, only 520 medical records completed these criteria.

Allergen Sensitivity

The allergen sensitivity was evaluated by skin prick test (SPT) using 42 allergens ALK-Abelló (*Port Washington, NY, United States*), distributed as 16 trees (*Acacia* sp., *Morus rubra*, *Olea europaea*, *Eucalyptus* sp., *Juniperus californica*, *Populus tremuloides*, *Populus alba*, *Liquidambar styraciflua*, *Cupressus arizonica*, *Alnus glutinosa*, *Ligustrum vulgare*, *Juniperus virginiana*, *Schinus molle*, *Quercus alba*, *Prosopis* sp., and *Casuarina* sp.); 11 types of grass (*Holcus lanatus*, *Sorghum halapense*, *Orchard grass*, *Lolium perenne*, *Phleum pratense*, *Agrostis alba*, *Anthoxanthum odoratum*, *Triticum aestivum*, *Cynodon dactylon*, *Hordeum vulgare*, and *Bromus pratensis*); 7 weeds (*Salsola kali*, *Taraxacum officinale*, *Artemisia vulgaris*, *Ambrosia trifida*, *Amaranthus retroflexus*, *Rumex crispus*, and *Chenopodium album*); 5 epithelia (*Felis domesticus*, *Oryctolagus cuniculus*, *Bos taurus*, *Canis lupus familiaris*, and *Equus caballus*), 2 mites (*Dermatophagoides pteronyssinus* and *Dermatophagoides farinae*), and *Blatella germanica*. We used 0.9% saline solution as a negative control and histamine hydrochloride (1:1000) as a positive control. All allergens were applied with a disposable polypropylene duo tip. The diameter of the wheal was recorded in millimeters, and a positive result measured 3 mm more in diameter than the negative control.¹⁶ For pollen counts and pollutants comparisons, we group those allergens that belong to the same family and cannot be distinguished from each other (For example, *Juniperus californica* and *Juniperus virginiana*, conformed to the Cupressaceae family).

Meteorological and Pollutants Data

The meteorological data (wind speed-WSP, temperature-TMP, relative humidity-RH) and information of pollutants (particulate matter of ten micrometers or less-PM₁₀, particulate matter of two point five micrometers or less-PM_{2.5}, carbon monoxide-CO, nitrogen monoxide-NO, NO₂, nitrogen oxides-NO_x, O₃, particulate matter coarse fraction-PM_{CO}, sulfur dioxide-SO₂) were extracted from the *Red de Meteorología y Radiación Solar-REDMET*, the *Red Automática de Monitoreo Atmosférico-RAMA*, and the *Red Manual de Monitoreo Atmosférico-REDMA* on the AIRE-CDMX websites.¹⁷ We calculated the averages of these data per day and by week (from 2014 to 2018) for subsequent comparison to the pollen counts.

Pollen Monitoring

Pollen count was measured with a Hirst-type volumetric spore trap-HS (*Burkard Manufacturing Co. Ltd., UK*), which is an impact suction sampler located at *Instituto Nacional de Enfermedades Respiratorias, Ismael Cosío Villegas*, in Tlalpan Mexico City placed outdoors at a height of 15 m from the ground. The HS mechanism consists of a constant incoming flow of air that impacts the receiving surface continuously and draws in a constant flow of 10 L/min that is arranged tightly around a cylindrical part called a drum that rotates continuously at 2 mm/h. The particles are deposited

sequentially on a Melinex tape impregnated with silicone fluid, which is cut into 24 fragments (48 mm) that are mounted on slides using glycerin jelly stained with fuchsine and fixed with phenol, which makes it possible to obtain accurate results with time (gr of pollens-gP/m³). The tape was analyzed every day by 40x optical microscopy (*Olympus CH 30, Tokyo, Japan*) by an aerobiology technician validated by the Aerobiology Network of *Red Mexicana de Aerobiología-REMA*. The pollen data were registered at the REMA website with PollenCntAdm software.

Pollen Data

We calculated the average of the daily concentrations over seven consecutive days to obtain the weekly mean concentrations to demonstrate the weekly variability in pollination (gP/m³ per week). Finally, we calculated the annual pollen integral (APIn: sum of the mean daily concentrations of the whole year) of each species to analyze the magnitudes and interannual differences in pollination. For the report of pollination dynamics, we defined the start of the pollination season as the time when the pollen concentration exceeded 2.5% and the finish as when 97.5% of the total annual count had been accumulated.¹⁸ Due to the peculiarities of most tree species in our region, where the flowering phase is in the winter, the start of pollination occurs at the end of the year, and the finish occurs at the beginning of the following year.¹⁹

Statistical Analysis

We analyzed the general pattern of allergic sensitization by age and disease, and then we compared these results with pollen monitoring (from 2014 to 2018). The association analysis between allergic sensitivity and pollen counts, both general and by period (before and after 2015), was performed with Epi Info v 7.0 (*Division of Health Informatics & Surveillance, DHIS, USA*) and was considered statistically significant with a *p*-value <0.05 and an odds ratio (OR) >1. Clinical quantitative variables and correlations between weather and pollutant variables were performed with SPSS software v.21 (*SPSS software, IBM, New York, USA*). Additionally, the pollen counts and pollutant concentrations were analyzed both per year and during the pollination season [season 1 (S1: 2014–2015), season 2 (S2: 2015–2016), season 3 (S3: 2016–2017), and season 4 (S4: 2017–2018)].

Results

Demographic Data and Allergic Sensitivity

A total of 520 patients without a predominance of sex were analyzed, with a median age of 20 years. 67.3% and 32.7% suffered from ARwA, and AR, respectively. A total of 63.6% had at least 2 allergens in the SPT (**Figure 2A**), 44.6% were exclusively sensitive to indoor allergens, 15.7% were sensitive to any pollen, and 39.6% were sensitive to both. The frequency of patients who were sensitive to at least one pollen was higher than that of patients who were exclusively indoors (55.3% vs 44.6%, *p*<0.001; OR=1.5) (**Figure 2B**). A total of 46.8% of the patients were only sensitive to trees, 8.6% to grasses, and 3.8% to weeds. The main intersections were the sensitivities to trees-grasses-weeds (16.3%) and trees-grasses (12.1%) (**Figure 2C**).

The most frequent pollen allergens were the trees of the Oleaceae family: *Fraxinus excelsior* (18%), *Olea europaea* (15%), and *Ligustrum lucidum* (14.2%), followed by *Betula verrucosa* (14.2%), *Quercus alba* (14%), *Quercus rubra* (14%), *Juniperus californica* (12.1%), *Juniperus virginiana* (10.1%), *Cupressus arizonica* (9.8%), and *Alnus verrucosa* (9.6%). These frequencies did not change when analyzed by disease. When we evaluated the sensitivity of the main tree allergens by specie and family, the frequency did not modify. *Fraxinus excelsior* was the principal allergen, followed by *Quercus* sp. -*Quercus alba* and *Quercus rubra*-, Cupressaceae family -*Juniperus californica*, *Juniperus virginiana*, and *Cupressus arizonica*-, and *Alnus* sp. However, indoor allergens were ranked first (**Figure 3**).

Allergic Sensitivity by Disease

We found that patients with allergic respiratory disease were more sensitive to Dermatophagoides or HDMS than to pollens independently if they suffered AR (89.1% vs 53.6%, *p*<0.001, OR=7.1) or ARwA (71.7% vs 50%, *p*<0.001, OR=2.5). In the same context, patients with AR were almost four times more sensitive to trees than to grasses (46.9% vs 18.5%, *p*<0.001, OR=3.8), and patients with ARwA were two times more sensitive (42.3% vs 20.6%, *p*<0.001, OR=2.8). The *Fraxinus* sp. and allergens of the Cupressaceae family were approximately two times more frequent than the other tree allergens in both diseases (**Figure 4** and **Supplementary Table 1**).




Figure 2 Distribution of allergy sensitivity. **(A)** Venn diagram of patients sensitive to only one allergen (monosensitive) and patients sensitive to two or more (polysensitive), **(B)** Venn diagram of patients sensitive to indoor allergens and/or pollen, **(C)** Venn diagram of patients sensitive to and/or different of pollen (trees and/or grasses and/or weeds).




Figure 3 The main sensitization to aeroallergens in skin prick test.

Pollen Monitoring

The main pollens identified during the five years by APIn corresponded to trees, mainly belonging to *Fraxinus* sp. (32.5%), Cupressaceae family (31.1%), *Alnus* sp. (16.1%), *Casuarina* sp. (9.5%), *Pinus* sp. (3.7%), *Quercus* sp. (2.7%), Myrtaceae family (2.6%), Salicilacea family (2.9%), Moraceae family (1.5%), and *Schinus* sp. (1.2%) (Figure 5A, Supplementary Table 2). Interestingly, after 2015, there were increases in the pollen counts of most species, reaching the




Figure 4 Allergic sensitivity by disease.

Notes: * $p < 0.001$; † $p < 0.01$; ‡ $p < 0.05$. Patients were more sensitive to HDM than to pollens. Both groups were more sensitive to trees than grasses ($p < 0.001$). Likewise, *Fraxinus* sp. and Cupressaceae family allergens were twice more prevalent than other tree allergens ($p < 0.05$).

Abbreviations: AR, allergic rhinitis; ARwA, allergic rhinitis with asthma; HDM, house dust mite. OR, odds ratio.




Figure 5 Pollen monitoring (2014–2018). (A) Pollen frequency, (B) Pollen count measured in annual pollen Integral (APIn), (C) Pollen counts by pollination season.

Abbreviations: S1, season 1; S2, season 2; S3, season 3; S4, season 4.

maximum values in 2016, except for *Casuarina* sp. (Figure 5B, Supplementary Table 2). However, when analyzing the main counts by the pollination season of each species (*Fraxinus* sp., Cupressaceae family, and *Alnus* sp.), the first season (2014–2015) had higher concentrations of pollen and was very similar in comparison to the second season (2015–2016), except for *Quercus* sp. However, these pollen gradually decreased in the third (2016–2017) and fourth seasons (2017–2018), reaching the most statistical significance when comparing S1 vs S4 ($p<0.001$). (Figure 5C and Table 1)

Allergic Sensitivity per Period

When comparing the sensitivity to pollen allergens, taking into consideration the start of the increase in pollen counts (2015), we analyzed the prevalence of sensitization by comparing two periods, the first period (2012 to 2014) and the second period (2015 to 2018). We identified that pollen sensitization significantly increased in the second period two times in AR and four times in ARwA ($p<0.01$ for each). Regarding specific sensitization by disease, there was an increase in the main sensitizer allergens in the AR group (*Quercus* sp. and *Alnus* sp. OR>3.0, $p<0.05$; by each), except for the Cupressaceae family. Instead, the ARwA group had increased sensitization frequencies for all allergens (*Quercus* sp. OR=5, Cupressaceae family OR=4, and *Alnus* sp. OR=3; $p<0.001$ by each). However, the most significant increase was observed for *Fraxinus* sp. in both the AR (OR=5, $p<0.001$) and ARwA (OR=9, $p<0.001$) groups. Moreover, HDMs were still the main allergens identified in both periods, and no increase in their frequency was detected (Figure 6 and Supplementary Table 3).

Meteorological Variables and Atmospheric Pollutants

When comparing the weather variables that occurred during the pollination season vs the no-pollination season, the temperature was lower in S1 and S2 ($p<0.04$), although it decreased in S3 and S4, even though it did not reach statistical significance. The WSP tended to be lower at S1, S2, and S4 but only showed significance at S3 ($p=0.01$); otherwise, the RH was lower in these seasons ($p<0.001$), excluding S3.

Regarding the pollutants, PM₁₀, PM_{2.5}, PM_{CO}, and NO_x were higher during the four pollination seasons ($p<0.01$ for each season). NO₂ also had a greater concentration in the pollination seasons ($p<0.001$); however, it showed a statistical tendency in S4. NO ($p=0.001$) and SO₂ ($p<0.02$) were higher in S1, S2, and S4. Meanwhile, CO had increased concentrations in S1 and S2 ($p<0.004$) (Table 2).

Global Correlations of Weather and Pollutant Correlations with Pollen Counts

When analyzing the counts from 2014 to 2018 of each pollinic season with both meteorological and pollution variables of the same time, we identified that temperature and WSP were the weather variables that correlated with the pollen of *Fraxinus* sp. and Cupressaceae family ($r=-0.30$, $p<0.01$ / $r=-0.18$, $p=0.03$), respectively. The pollutants (PM₁₀, PM_{2.5}, CO, NO, NO₂, NO_x, and PM_{CO}) correlated positively with *Fraxinus* sp., Cupressaceae family, and *Alnus* sp. ($r>0.18$ and $p<0.05$ for each). Conversely, *Quercus* sp. pollen only presented significant correlations with PM_{2.5} ($r=0.22$ and $p=0.04$), CO ($r=0.25$ and $p=0.01$), NO ($r=0.2$, $p=0.03$), and NO_x ($r=0.20$, $p=0.03$), and it was the only pollen that correlated with SO₂ ($r=0.17$ and $p=0.05$). (Table 3 and Supplementary Figure 1).

Correlations by Pollination Season

We analyzed the correlations between air pollution and/or meteorological variables and airborne pollen during the four pollen seasons. The most constant and significant correlations were identified among air pollutants, and pollen concentrations were observed for PM₁₀, NO₂, and PM_{CO} with *Fraxinus* sp. in all seasons ($p\leq 0.02$ for each season). However, NO and NO_x concentrations were also related to *Fraxinus* sp. during S2, S3, and S4 ($p\leq 0.02$). NO_x was positively correlated with the Cupressaceae family in S2 and S3 ($p\leq 0.01$), and NO was positively correlated with the pollen of *Quercus* sp. in S1, S2, and S3 ($p<0.04$). Additionally, O₃ was related to *Alnus* sp. in S2, S3, and S4 ($p\leq 0.01$).

Interestingly, we found a notable increase from S2 in the number of significant correlations with the pollution variables that gradually decrease in the following seasons. Likewise, the contaminants mentioned above, in addition to CO and SO₂ and other meteorological variables (TMP, WSP, and RH), had significant correlations with at least one pollen in some pollination seasons (Figure 7, Supplementary Table 4, and Supplementary Figure 2).

Table I Seasons Comparison

Variable	C1			C2			C3		
	Season 1	Season 2	p	Season 1	Season 3	p	Season 1	Season 4	p
<i>Fraxinus</i> sp.	1161.50 (428–2372.25)	930.5 (292–1384.75)	0.06	1161.50 (428–2372.25)	634 (269–1095)	0.01	1161.50 (428–2372.25)	459 (377.75–644.75)	0.001
<i>Alnus</i> sp.	497 (238.5–1294)	342.5 (148.75–809.25)	0.1	497 (238.5–1294)	56 (8–633)	0.001	497 (238.5–1294)	58 (0–510.5)	<0.001
Cupressaceae	971.5 (489–1371.75)	585.50 (403.75–799.5)	0.01	971.5 (489–1371.75)	525 (359–864)	0.009	971.5 (489–1371.75)	368 (159.25–470)	<0.001
<i>Quercus</i> sp.	6.50 (0–34)	15 (0–36.25)	0.43	6.50 (0–34)	5 (0–50)	0.77	6.50 (0–34)	6 (1.25–22.75)	0.59
TMP	15.67 (14.20–16.78)	16.51 (14.93–19.58)	0.13	15.67 (14.20–16.78)	16.34 (15.22–19.21)	0.071	15.67 (14.20–16.78)	17.09 (14.02–18.22)	0.23
WSP	1.90 (1.61–2.11)	1.86 (1.67–2.11)	0.89	1.90 (1.61–2.11)	1.90 (1.74–2.08)	0.81	1.90 (1.61–2.11)	1.93 (1.70–2.22)	0.53
RH	50.26 (43.77–55.77)	43.55 (34.08–52.95)	0.037	50.26 (43.77–55.77)	59.65 (49.39–69.59)	0.001	50.26 (43.77–55.77)	43.69 (41.26–49.49)	0.02
PM ₁₀	57.57 (44.50–66.16)	66.71 (53.54–76.64)	0.04	57.57 (44.50–66.16)	56.95 (48.33–67.52)	0.66	57.57 (44.50–66.16)	53.21 (46.33–60.15)	0.39
PM _{2.5}	27.85 (22.48–29.71)	33 (27.75–38.35)	0.009	27.85 (22.48–29.71)	26.80 (22.20–31.31)	0.85	27.85 (22.48–29.71)	26.27 (22.59–29.31)	0.54
CO	0.96 (0.89–1.09)	0.92 (0.85–1.07)	0.6	0.96 (0.89–1.09)	0.65 (0.52–0.91)	<0.001	0.96 (0.89–1.09)	0.54 (0.46–0.64)	<0.001
NO	36.21 (30.28–40.02)	31.96 (24.14–41.83)	0.37	36.21 (30.28–40.02)	25.26 (17.31–32.04)	<0.001	36.21 (30.28–40.02)	24.72 (22.29–31.89)	0.001
NO ₂	34.99 (31.84–38.41)	37.36 (33.81–41.89)	0.04	34.99 (31.84–38.41)	32.28 (28.73–34.98)	0.04	34.99 (31.84–38.41)	33.91 (29.28–37.65)	0.39
NO _x	70.47 (62.70–77.71)	68.34 (60.32–83.82)	0.98	70.47 (62.70–77.71)	57.76 (45.98–68.97)	0.001	70.47 (62.70–77.71)	58.64 (53.83–70.83)	0.007
O ₃	20.11 (16.55–26.26)	23.49 (19.05–36.41)	0.02	20.11 (16.55–26.26)	27.76 (18.69–32.82)	0.006	20.11 (16.55–26.26)	23.63 (20–29.35)	0.02
PM _{CO}	30.77 (19.60–32.63)	32.75 (26.63–38.25)	0.1	30.77 (19.60–32.63)	29.90 (24.06–36.78)	0.67	30.77 (19.60–32.63)	27.87 (23.76–31.82)	0.33
SO ₂	8.22 (5.55–10.16)	7.79 (5.31–17.49)	0.69	8.22 (5.55–10.16)	6.95 (4.43–8.53)	0.63	8.22 (5.55–10.16)	9.69 (6.83–13.57)	0.18
Variable	C4			C5			C6		
	Season 2	Season 3	p	Season 2	Season 4	p	Season 3	Season 4	p
<i>Fraxinus</i> sp.	930.5 (292–1384.75)	634 (269–1095)	0.25	930.5 (292–1384.75)	459 (377.75–644.75)	0.021	634 (269–1095)	459 (377.75–644.75)	0.13
<i>Alnus</i> sp.	342.5 (148.75–809.25)	56 (8–633)	0.001	342.5 (148.75–809.25)	58 (0–510.5)	0.001	56 (8–633)	58 (0–510.5)	0.01
Cupressaceae	585.50 (403.75–799.5)	525 (359–864)	0.83	585.50 (403.75–799.5)	368 (159.25–470)	<0.001	525 (359–864)	368 (159.25–470)	<0.001
<i>Quercus</i> sp.	15 (0–36.25)	5 (0–50)	0.82	15 (0–36.25)	6 (1.25–22.75)	0.74	5 (0–50)	6 (1.25–22.75)	0.65
TMP	16.51 (14.93–19.58)	16.34 (15.22–19.21)	0.92	16.51 (14.93–19.58)	17.09 (14.02–18.22)	0.84	16.34 (15.22–19.21)	17.09 (14.02–18.22)	0.65

WSP	1.86 (1.67–2.11)	1.90 (1.74–2.08)	0.67	1.86 (1.67–2.11)	1.93 (1.70–2.22)	0.35	1.90 (1.74–2.08)	1.93 (1.70–2.22)	0.64
RH	43.55 (34.08–52.95)	59.65 (49.39–69.59)	<0.001	43.55 (34.08–52.95)	43.69 (41.26–49.49)	0.59	59.65 (49.39–69.59)	43.69 (41.26–49.49)	<0.001
PM₁₀	66.71 (53.54–76.64)	56.95 (48.33–67.52)	0.022	66.71 (53.54–76.64)	53.21 (46.33–60.15)	<0.001	56.95 (48.33–67.52)	53.21 (46.33–60.15)	0.13
PM_{2.5}	33 (27.75–38.35)	26.80 (22.20–31.31)	0.002	33 (27.75–38.35)	26.27 (22.59–29.31)	<0.001	26.80 (22.20–31.31)	26.27 (22.59–29.31)	0.64
CO	0.92 (0.85–1.07)	0.65 (0.52–0.91)	<0.001	0.92 (0.85–1.07)	0.54 (0.46–0.64)	<0.001	0.65 (0.52–0.91)	0.54 (0.46–0.64)	0.004
NO	31.96 (24.14–41.83)	25.26 (17.31–32.04)	0.003	31.96 (24.14–41.83)	24.72 (22.29–31.89)	0.01	25.26 (17.31–32.04)	24.72 (22.29–31.89)	0.46
NO₂	37.36 (33.81–41.89)	32.28 (28.73–34.98)	<0.001	37.36 (33.81–41.89)	33.91 (29.28–37.65)	0.01	32.28 (28.73–34.98)	33.91 (29.28–37.65)	0.23
NO_x	68.34 (60.32–83.82)	57.76 (45.98–68.97)	0.001	68.34 (60.32–83.82)	58.64 (53.83–70.83)	0.006	57.76 (45.98–68.97)	58.64 (53.83–70.83)	0.31
O₃	23.49 (19.05–36.41)	27.76 (18.69–32.82)	0.54	23.49 (19.05–36.41)	23.63 (20–29.35)	0.8	27.76 (18.69–32.82)	23.63 (20–29.35)	0.18
PM_{CO}	32.75 (26.63–38.25)	29.90 (24.06–36.78)	0.21	32.75 (26.63–38.25)	27.87 (23.76–31.82)	0.003	29.90 (24.06–36.78)	27.87 (23.76–31.82)	0.12
SO₂	7.79 (5.31–17.49)	6.95 (4.43–8.53)	0.08	7.79 (5.31–17.49)	9.69 (6.83–13.57)	0.76	6.95 (4.43–8.53)	9.69 (6.83–13.57)	0.003

Abbreviations: TMP, temperature; WSP, wind speed; RH, relative humidity; PM₁₀, particulate matter of 10 micrometers or less; PM_{2.5}, particulate matter of 2.5 micrometers or less; CO, carbon monoxide; NO, nitrogen monoxide; NO₂, nitrogen dioxide; NO_x, nitrogen oxides; O₃, ozone; PM_{CO}, coarse fraction of PM₁₀ and PM_{2.5} particles; SO₂, sulfur dioxide; C1, comparison 1 (2014–2015 vs 2015–2016); C2, comparison 2 (2014–2015 vs 2016–2017); C3, comparison 3 (2014–2015 vs 2017–2018); C4, comparison 4 (2015–2016 vs 2016–2017); C5, comparison 5 (2015–2016 vs 2017–2018); C6, comparison 6 (2016–2017 vs 2017–2018).




Figure 6 Allergic sensitivity per period by disease.

Notes: * $p < 0.001$; † $p < 0.01$; ‡ $p < 0.05$. In the second period (2015–2018), pollen sensitization increased significantly, twice in AR and four times in ARwA ($p < 0.01$). *Quercus* sp. and *Alnus* sp. allergens sensitization raised in the AR group ($p < 0.05$), whereas the ARwA group increased sensitization frequencies for *Quercus* sp., Cupressaceae family, and *Alnus* sp. ($p < 0.001$). However, *Fraxinus* sp. sensitization was the one that increased the most in both groups ($p < 0.001$).

Abbreviations: AR, allergic rhinitis; ARwA, allergic rhinitis with asthma; HDM, house dust mite; NS, non-significant; NA, not applicable.

Discussion

In the present study, we associated the sensitivity to aeroallergens in ARD patients with the pollen counts and the possible relationships with atmospheric pollutants. Since 2015, the pollination of trees has augmented for most taxa, especially for *Fraxinus* sp., *Quercus* sp., and the Cupressaceae family, which was associated with increases in sensitization to these species. In particular, some pollutants, such as PM₁₀, NO₂, and PM_{CO}, were directly related to *Fraxinus* sp. pollination.

Pollinosis is the inflammation of the conjunctival, nasal, and/or bronchial mucosa induced by allergens from pollen grains through a type I hypersensitivity mechanism.²⁰ There is evidence that shows that half of AR patients are sensitive to any kind of pollen,²¹ other groups have reported that its prevalence has almost doubled in recent decades.²² Due to its clinical importance and tendency to increase, some studies have analyzed the factors that could be associated with this phenomenon.²³

Consistent with previous reports, we identified that our population has a polysensitized pattern,²⁴ with the most prevalent allergen sensitizer being *Dermatophagoides* sp., plus at least one pollen (members of the Lamiales, Fagales, and Cupressales orders). Interestingly, this study showed tree pollen allergies relegate the grasses to less relevant positions, which indicates that the pattern of allergic sensitivity to pollen in Mexican patients is different from that described in Europe, where grasses are the main sensitizers.²¹

There is evidence that a population is sensitized to the pollen species to which they are more exposed in its geographic localization.²⁵ In our case, the species that induced greater levels of sensitization were *Fraxinus* sp., *Quercus* sp., and the Cupressaceae family. Some reports refer to the pollens of the Fagales, Cupressales, and Lamiales orders as being among the main sensitizers.^{7,21} These observations are supported by other palynological studies developed in Mexico City, which described that the high counts of these pollens are identified during the dry season (from November to May);^{19,26} (Supplementary Figure 3). Since 1977, *Pinus* sp. has been reported as the most frequent pollen in Mexico City and traditionally has been considered a non-allergenic pollen²⁷ for this reason, it was not included in the SPT by our group. However, its sensitivity is increased in populations that live near forested areas.²⁸ Trees are the dominant species of the forest surrounding this city, particularly in southern Mexico City, which may favor the allergic respiratory diseases in susceptible patients.

Table 2 Meteorological and Pollutants Analysis Between Pollination and No Pollination Seasons

Variables	Season 1			Season 2			Season 3			Season 4		
	Pollination Season	No Pollination Season	p	Pollination Season	No Pollination Season	p	Pollination Season	No Pollination Season	p	Pollination Season	No Pollination Season	p
<i>Fraxinus</i> sp.	1161.5 (428–2372.25)	8.50 (1–24.75)	<0.001	930.5 (292–1384)	37 (9.5–37)	<0.001	634 (269–1095)	22 (7.5–44.75)	<0.001	459 (377.75–644.75)	4.5 (2.75–9)	<0.001
<i>Alnus</i> sp.	497 (238.5–1294)	0.000 (0–1)	<0.001	342.5 (148.75–809.25)	18 (1.5–18)	<0.001	56 (8–633)	4 (2–13.5)	<0.001	58 (0–510.5)	0 (0–0)	<0.001
Cupressaceae	971.50 (489–1371.75)	30.50 (22–69.5)	<0.001	585.5 (403.75–799.5)	196 (37.5–196)	<0.001	525 (359–864)	95 (55.75–212)	<0.001	368 (159.25–470)	214.5 (151–366.25)	0.14
<i>Quercus</i> sp.	6.50 (0–34)	0 (0–0)	<0.001	15 (0.0–36.25)	0.0 (0.0–0.0)	<0.001	5 (0.0–50)	0.0 (0.0–0.0)	<0.001	6 (1.25–22.75)	0 (0–0)	<0.001
TMP	15.67 (14.20–16.78)	17.16 (16.31–17.98)	0.003	16.51 (14.93–19.58)	17.8 (17.01–18.31)	0.042	16.34 (15.22–19.21)	17.46 (17.3–18.07)	0.104	17.09 (14.02–18.22)	17.20 (16.60–18.83)	0.24
WSP	1.90 (1.61–2.11)	2.06 (1.83–2.20)	0.095	1.86 (1.67–2.11)	2.02 (1.80–2.28)	0.067	1.90 (1.74–2.08)	2.11 (1.90–2.45)	0.012	1.93 (1.70–2.22)	2.06 (1.91–2.4)	0.09
RH	50.26 (43.77–55.77)	63.83 (59.99–67.05)	<0.001	43.55 (34.08–52.95)	58.83 (55.75–61.96)	<0.001	59.65 (49.39–69.59)	61.25 (57.85–63.45)	0.54	43.69 (41.26–49.49)	61.98 (56.72–68.24)	<0.001
PM ₁₀	57.57 (44.50–66.16)	38.57 (38.80–42.64)	<0.001	66.71 (53.64–76.64)	47.92 (44.32–50.92)	<0.001	56.95 (48.33–67.52)	41 (37.41–46.14)	<0.001	53.21 (46.33–60.15)	37.59 (29.46–42.8)	<0.001
PM _{2.5}	27.85 (22.48–29.71)	18.16 (17.35–22.21)	0.001	33 (27.75–38.35)	26.57 (23.87–29.89)	0.001	26.8 (22.2–31.31)	20.71 (16.4–21.83)	<0.001	26.27 (22.59–29.31)	19.16 (13.8–23.06)	0.002
CO	0.96 (0.89–1.09)	0.87 (0.77–0.93)	0.004	0.92 (0.85–1.07)	0.81 (0.73–0.91)	0.001	0.65 (0.52–0.91)	0.73 (0.65–0.82)	0.128	0.54 (0.46–0.64)	0.54 (0.48–0.68)	0.71
NO	36.21 (30.28–40.02)	23.64 (21.41–29.57)	<0.001	31.96 (24.14–41.83)	23.62 (20.05–28.66)	0.001	25.26 (17.31–32.04)	20.19 (19.10–22.75)	0.078	24.72 (22.29–31.89)	17.58 (15.41–21.55)	0.005
NO ₂	34.99 (31.84–38.41)	26.68 (24.66–29.74)	<0.001	37.36 (33.81–41.89)	29.85 (27.90–33.91)	<0.001	32.28 (28.73–34.98)	26.96 (24.6–29.82)	0.001	33.91 (29.28–37.65)	30.33 (28.07–31.79)	0.06
NO _x	70.47 (62.70–77.61)	51.95 (47.23–57.33)	<0.001	68.34 (60.32–83.82)	55.41 (48.51–59.96)	<0.001	57.76 (45.98–68.97)	49.27 (44.35–53.04)	0.006	58.64 (53.83–70.83)	46.89 (45.77–57.09)	0.007
O ₃	20.11 (16.55–26.26)	20.44 (17.17–25.42)	0.905	23.49 (19.05–36.41)	26.42 (22.03–27.83)	0.773	27.76 (18.69–32.82)	25.02 (20.32–28.22)	0.327	23.63 (20–29.35)	21.13 (16.82–24.31)	0.04
PM _{co}	30.77 (19.60–32.63)	17.76 (15.52–21.08)	0.001	32.75 (26.63–38.25)	20.70 (18.61–23.73)	<0.001	29.90 (24.06–36.78)	22.18 (18.25–23.81)	<0.001	27.87 (23.76–31.82)	17.73 (13.67–19.67)	<0.001
SO ₂	8.22 (5.55–10.16)	4.74 (4.20–5.90)	0.002	7.79 (5.31–17.49)	5.89 (5.01–7.49)	0.027	6.95 (4.43–8.53)	5.21 (4.39–7.06)	0.163	9.69 (6.83–13.57)	5.84 (5.15–7.13)	0.001

Abbreviations: TMP, temperature; WSP, wind speed; RH, relative humidity; PM₁₀, particulate matter of 10 micrometers or less; PM_{2.5}, particulate matter of 2.5 micrometers or less; CO, carbon monoxide; NO, nitrogen monoxide; NO₂, nitrogen dioxide; NO_x, nitrogen oxides; O₃, ozone; PM_{co}, coarse fraction of PM₁₀ and PM_{2.5} particles; SO₂, sulfur dioxide; Season 1 (2014–2015); Season 2 (2015–2016); Season 3 (2016–2017); Season 4 (2017–2018).

Table 3 Global Correlations Between Meteorological Variables and Pollution with Pollen

Variables	<i>Fraxinus</i> sp.		Cupressaceae		<i>Alnus</i> sp.		<i>Quercus</i> sp.	
	Pearson r	p	Pearson r	p	Pearson r	p	Pearson r	p
TMP	-0.31	<0.001	-0.13	0.16	-0.11	0.24	0.11	0.24
WSP	-0.14	0.11	-0.18	0.03	-0.14	0.12	-0.15	0.1
RH	-0.05	0.56	0.05	0.59	-0.06	0.52	0.03	0.75
PM₁₀	0.50	<0.001	0.44	<0.001	0.31	0.001	0.14	0.12
PM_{2.5}	0.34	<0.001	0.38	<0.001	0.34	<0.001	0.21	0.02
CO	0.37	<0.001	0.38	<0.001	0.25	0.006	0.25	0.01
NO	0.35	<0.001	0.26	0.004	0.29	0.001	0.20	0.03
NO₂	0.34	<0.001	0.24	0.008	0.18	0.05	0.15	0.11
NO_x	0.37	<0.001	0.27	0.003	0.27	0.003	0.20	0.03
O₃	-0.18	0.06	-0.03	0.78	-0.07	0.45	0.03	0.76
PM_{CO}	0.56	<0.001	0.41	<0.001	0.23	0.01	0.07	0.48
SO₂	0.14	0.13	0.003	0.97	0.01	0.88	0.17	0.05

Abbreviations: TMP, temperature; WSP, wind speed; RH, relative humidity; PM₁₀, particulate matter of 10 micrometers or less; PM_{2.5}, particulate matter of 2.5 micrometers or less; CO, carbon monoxide; NO, nitrogen monoxide; NO₂, nitrogen dioxide; NO_x, nitrogen oxides; O₃, ozone; PM_{CO}, coarse fraction of PM₁₀ and PM_{2.5} particles; SO₂, sulfur dioxide.

The intensity of pollen exposure has been defined by expert consensus.²⁹ Currently, there is no specific pollen count associated with sensitization.³⁰ However, this can also be due to the quantification method used in some studies. Bousquet described that the pollen counts differ if they are analyzed by gP/m³ or weight (mg/m³). For example, in France, Cupressaceae were the most abundant by count. However, grasses were the most abundant when the measurements were calculated in mg/m³ despite its lower counts.³¹ In addition, the traditional pollen count also differs from the allergenic proteins suspended in the atmosphere.³² Therefore, some species can cause symptoms with lower counts.³¹ All the above mentioned observations are justified because more than one-third of our population is sensitized to grasses or *Quercus* sp., which is the second sensitizer in our population despite their low counts. Interestingly, the pollens responsible for the main sensitizations have doubled in their counts since 2015. This phenomenon has been reported at other stations of pollen monitoring in Mexico City.²⁶ Likewise, some reports have described great augmentations of the interannual pollen counts for certain species, such as every three years for *Cryptomeria* sp., from two to four years for the Fagales order (*Betula* sp., *Corylus* sp., and *Alnus* sp.), and every three years for *Fraxinus* sp.³³⁻³⁵ It is probable that 2015 corresponds to the zenith of pollen due to synchronic mass seeding; however, we need a longer period for identifying the flowering rhythms for the main species in our area. This might be due to the natural phenological behavior of the long-lived plants, which had a synchronic production of seeds at long intervals, where resource matching first had to gain the substrates needed for flowering readiness, and mass seeding reduced levels of loss to seed predators.³⁶

The increase in pollen production has been described as a factor involved in the augmentation of sensitization frequency. For example, Lee KS described a positive correlation between the duration of the pollen season and the rates of sensitization to tree pollens in Korea (~0.28% per year, a total of 8295 new patients sensitized to Fagales species over 22 years).²³ Similar cases have been reported in Germany with *Betula* sp. and in Japan with *Cryptomeria japonica*.^{37,38} In the same context, Switzerland has reported this phenomenon with the most prevalent species (birch and grasses). In contrast, the prevalence did not increase when there were no changes in pollen counts.³⁷ In our case, the great




Figure 7 Seasons correlations between significant meteorological variables and pollutants with pollen.

Notes: The number of significant correlations with pollution variables increased significantly from S1 to S2. The most consistent and significant correlations were found between air pollutants and pollen concentrations in all seasons for PM₁₀, NO₂, and PM_{CO} with *Fraxinus* sp. ($p < 0.02$).

Abbreviations: PM₁₀, particulate matter of 10 micrometers or less; NO₂, nitrogen dioxide; PM_{CO}, coarse fraction of PM₁₀ and PM_{2.5} particles; SO₂, sulfur dioxide; Season 1 (2014–2015); Season 2 (2015–2016); Season 3 (2016–2017); Season 4 (2017–2018).

augmentation in pollen counts was associated with respective increases in sensitization frequency for the main tree species (*Fraxinus* sp., Cupressaceae family, and *Alnus* sp.).

However, as previously mentioned, some pollens (*Quercus* sp. and *Casuarina* sp.) did not show significant increases in their counts, although we reported increases in sensitization to these pollens in our population. This phenomenon has been reported in other countries. Hirsch T. in Germany analyzed two populations located in Dresden and Munich exposed to different pollen amounts. The author described a higher prevalence of sensitization in patients with less exposure and vice versa, suggesting that some environmental factors, such as pollutants, are involved in this phenomenon.³⁹

In this sense, we evaluated the role of pollutants in our geographical area. There is evidence that the variation in the concentrations of some pollutants, such as PM₁₀, PM_{2.5}, CO, NO, NO₂, NO_X, SO₂, and O₃, is associated with changes in the pollen counts of *Alnus* sp. and *Fraxinus* sp. due to these molecules modifying the flowering and pollination periods.^{12,40} In our case, positive correlations of PM₁₀, NO₂, and PM_{CO} were found, mainly to *Fraxinus* sp. in all seasons. NO₂ exposure induces posttranslational modification (S-nitrosylation) of ragweed plants, resulting in an increase in Amb a 1 isoforms and increasing the allergenicity of ragweed pollen.⁴¹ Additionally, the increase

in NO₂ and SO₂ during the *Quercus* sp. pollen season can facilitate the bioavailability of airborne pollen allergens.⁴² These events may increase the incidence of allergic diseases in contaminated areas. Similarly, ultrafine particles such as PM₁₀ bind to pollen, altering its allergic properties, enhancing allergen release, and ultimately acting as an adjuvant, precipitating allergic disease.¹³ Likewise, PM_{CO} was related to the variations in the concentrations of some particular pollens in our study. This is likely due to this molecule being a nonspecific classification molecule between PM_{2.5} and PM₁₀. The mechanisms by which pollutants may alter *Fraxinus* sp. pollen allergenicity remain to be shown.

Nevertheless, reforestation has increased pollen counts. Patients become sensitive to at least one species of tree involved in reforestation programs.⁴³ Although Mexican laws guarantee a healthy environment and population well-being, Mexico City lacks a census of trees utilized in reforestation.⁴⁴ Mexico has the highest density and diversity of oaks,⁴⁵ which, alongside ash and cedar, are the main species used in reforestation in many cities of Mexico.⁴⁶

In our study, the sensitivity to pollen was not specific to any one particular entity (AR or ARwA). There is evidence that a particular allergic disease could be sensitive to a specific allergen. For example, asthma without AR patients had a greater sensitivity to ragweed,⁴⁷ and most of our patients had asthma with AR.

We performed phenological analysis following current international recommendations for aerobiological sampling using standardized instruments and trained professionals. The sample corresponded to patients who undertook their daily activities in the vicinity of the aerobiological sampling area, and the sensitization assessments were carried out following the recommendations of allergic diagnosis using just one brand of allergen kit, avoiding allergenic potency bias. Among the limitations of the present study is that we did not analyze other environmental factors, such as biotic stress (infection), abiotic stress (nutritional deficiency), and other climate data. Interestingly, in 2015, there were increases in regional temperature and precipitation intensity compared to other years.⁴⁸ However, is necessary to evaluate these interactions additionally, it is important to describe the morphologic changes or particles adhered to the pollen grains induced by contaminants through scanning electron microscope¹³ or analyze isoforms from pollen exposed to pollutants by proteomic technology.⁴⁹ Unfortunately, the SARS-CoV-2 pandemic prevented the monitoring of both phenological records and those data inherent to allergic sensitization. Knowing the pollination and the prevalence of pollinosis in patients with ARD is very important in health policy. Pollination can negatively affect lung function.⁵⁰ Sensitization to the Oleaceae and Fagaceae families is associated with an asthmatic crisis.^{51,52} This phenomenon can be precipitated by meteorological phenomena different from those already analyzed, such as electrical storms, which can interact with the development of clinical outcomes.⁵¹ Finally, we expect these results to help clarify the relationship between pollination and allergic sensitization and to be considered for the development of reforestation policies in both national and international regions.

Conclusion

The high increases in pollen counts of *Fraxinus* sp., Cupressaceae family, and *Alnus* sp. are associated with increases in the frequencies of sensitization to these species in ARD patients. This phenomenon is more related to the correlations of atmospheric pollutants such as PM₁₀, PM_{CO}, and NO₂ with *Fraxinus* sp. pollen.

Abbreviations

ARD, allergic respiratory disease; ARDs; allergic respiratory diseases; AR, allergic rhinitis; ARwA, AR with asthma; APIn, annual pollen integral; CO, carbon monoxide; HS, Hirst-type volumetric spore trap; HDMS, House dust mites; NO₂, Nitrogen dioxide; NO, nitrogen monoxide; NO_x, nitrogen oxides; OR, odds ratio; O₃, ozone; PM_{CO}, particulate matter coarse fraction; PM_{2.5}, particulate matter of two point five micrometers or less; PM₁₀, particulate matter of ten micrometers or less; RAMA, Red Automática de Monitoreo Atmosférico; REDMET, Red de Meteorología y Radiación Solar; REDMA, Red Manual de Monitoreo Atmosférico; REMA, Red Mexicana de Aerobiología; RH, relative humidity; S1, season 1; S2, season 2; S3, season 3; season 4; SPT, skin prick test; SO₂, sulfur dioxide; TMP, temperature; WSP, wind speed.

Data Sharing Statement

The pollutant and meteorological data were obtained from AIRE-CDMX websites (<http://www.aire.cdmx.gob.mx/default.php>), including RAMA, Red Automática de Monitoreo Atmosférico; REDMET, Red de Meteorología y Radiación Solar; REDMA, Red Manual de Monitoreo Atmosférico; REMA, Red Mexicana de Aerobiología.

Ethics Approval and Informed Consent

This study was performed in accordance with the principles stated in the Declaration of Helsinki, and the study protocol was approved by the science and bioethics committee for research of the Instituto Nacional de Enfermedades Respiratorias “Ismael Cosío Villegas” (Approval number: E03-13).

Consent for Publication

All authors have reviewed and approved the present manuscript.

Acknowledgments

We thank Felipe Fuentes-Arenas, Christian Rubén Merino-Camacho, Nancy Haydée Serrano-Pérez, Maria Itzel Vargas-Parra, Katia Vanessa Gutiérrezz-Quiroz, and César Guerrero-Guerra for their support and technical assistance.

Author Contributions

All authors have contributed significantly to this work, either in conception, study design, implementation, data acquisition, analysis, and interpretation or in all of these areas. All authors participated in drafting, revising, or critically reviewing the article; gave final approval of the version to be published; agreed on the journal to which the article has been submitted; and agreed to be responsible for all aspects of the work.

Funding

There are no funding sources to disclose.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Kuo CRW, Chan R, Lipworth B. Does unified allergic airway disease impact on lung function and type 2 biomarkers? *Allergy Asthma Clin Immunol.* 2019;15(1):4–7. doi:10.1186/s13223-019-0388-4
2. Brożek JL, Bousquet J, Agache I, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines-2016 revision. *J Allergy Clin Immunol.* 2017;140(4):950–958. doi:10.1016/J.JACI.2017.03.050
3. Guerra S, Sherrill DL, Martínez FD, Barbee RA. Rhinitis as an independent risk factor for adult-onset asthma. *J Allergy Clin Immunol.* 2002;109(3):419–425. doi:10.1067/MAI.2002.121701
4. Padem N, Saltoun C. Classification of asthma. *Allergy Asthma Proc.* 2019;40(6):385–388. doi:10.2500/AAP.2019.40.4253
5. Global Strategy For Asthma Management and Prevention. Global initiative for asthma (updated 2012); 2012. Available from: <https://ginasthma.org/archived-%0areports/>. Accessed February 16, 2022.
6. Miller JD. The role of dust mites in allergy. *Clin Rev Allergy Immunol.* 2019;57(3):312–329. doi:10.1007/S12016-018-8693-0
7. Guide US, Matricardi PM, Kleine-tebbe J, et al. User’s guide EAACI molecular allergology; 2016.
8. Geller-Bernstein C, Portnoy JM. The clinical utility of pollen counts. *Clin Rev Allergy Immunol.* 2019;57(3):340–349. doi:10.1007/S12016-018-8698-8
9. Negrini AC, Negrini S, Giunta V, Quaglini S, Ciprandi G. Thirty-year survey on airborne pollen concentrations in Genoa, Italy: relationship with sensitizations, meteorological data, and air pollution. *Am J Rhinol Allergy.* 2011;25(6):232–241. doi:10.2500/ajra.2011.25.3729
10. Sierra-Vargas MP, Teran LM. Air pollution: impact and prevention. *Respirology.* 2012;17(7):1031–1038. doi:10.1111/J.1440-1843.2012.02213.X
11. Thurston GD, Balmes JR, Garcia E, et al. Outdoor air pollution and new-onset airway disease. an official American thoracic society workshop report. *Ann Am Thorac Soc.* 2020;17(4):387. doi:10.1513/ANNALSATS.202001-046ST
12. Rahman A, Luo C, Khan MHR, Ke J, Thilakanayaka V, Kumar S. Influence of atmospheric PM2.5, PM10, O3, CO, NO2, SO2, and meteorological factors on the concentration of airborne pollen in Guangzhou, China. *Atmos Environ.* 2019;212:290–304. doi:10.1016/J.ATMOENV.2019.05.049
13. Ortega-Rosas CI, Meza-Figueroa D, Vidal-Solano JR, González-Grijalva B, Schiavo B. Association of airborne particulate matter with pollen, fungal spores, and allergic symptoms in an arid urbanized area. *Environ Geochem Health.* 2021;43(5):1761–1782. doi:10.1007/S10653-020-00752-7
14. Pavón-Romero GF, Serrano-Pérez NH, García-Sánchez L, Ramírez-Jiménez F, Terán LM. Neuroimmune pathophysiology in asthma. *Front Cell Dev Biol.* 2021;9. doi:10.3389/FCELL.2021.663535

15. Bousquet J, Khaltaev N, Cruz AA, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy*. 2008;63(SUPPL.86):8–160. doi:10.1111/J.1398-9995.2007.01620.X
16. Heinzerling L, Mari A, Bergmann KC, et al. The skin prick test - European standards. *Clin Transl Allergy*. 2013;3(1):1–10. doi:10.1186/2045-7022-3-3
17. Dirección de Monitoreo Atmosférico. Informes anuales de calidad del aire; 2018. Available from: <http://www.aire.cdmx.gob.mx/default.php?opc=%27akbi%27>. Accessed February 9, 2022.
18. Andersen TB. A model to predict the beginning of the pollen season. *Grana*. 2009;30:269–275. doi:10.1080/00173139109427810
19. Calderón-Ezquerro MC, Guerrero-Guerra C, Martínez-López B, et al. First airborne pollen calendar for Mexico City and its relationship with bioclimatic factors. *Aerobiologia*. 2016;32(2):225–244. doi:10.1007/s10453-015-9392-4
20. Taketomi EA, Sopenete MC, Moreira PFDS, Vieira FDAM. Pollen allergic disease: pollens and its major allergens. *Braz J Otorhinolaryngol*. 2006;72(4):562–567.
21. Burbach GJ, Heinzerling LM, Edenharter G, et al. GA2LEN skin test study II: clinical relevance of inhalant allergen sensitizations in Europe. *Allergy Eur J Allergy Clin Immunol*. 2009;64(10):1507–1515. doi:10.1111/j.1398-9995.2009.02089.x
22. Bousquet J, Anto JM, Bachert C, et al. Allergic rhinitis. *Nat Rev Dis Prim*. 2020;6(1):95. doi:10.1038/s41572-020-00227-0
23. Lee KS, Kim K, Choi YJ, et al. Increased sensitization rates to tree pollens in allergic children and adolescents and a change in the pollen season in the metropolitan area of Seoul, Korea. *Pediatr Allergy Immunol*. 2021;32(5):872–879. doi:10.1111/PAI.13472
24. Ciprandi G, Incorvaia C, Puccinelli P, Soffia S, Scurati S, Frati F. Polysensitization as a challenge for the allergist: the suggestions provided by the polysensitization impact on allergen immunotherapy studies. *Expert Opin Biol Ther*. 2011;11(6):715–722. doi:10.1517/14712598.2011.576246
25. Wang XY, Ma TT, Wang XY, et al. Prevalence of pollen-induced allergic rhinitis with high pollen exposure in grasslands of northern China. *Allergy*. 2018;73(6):1232–1243. doi:10.1111/ALL.13388
26. Calderon-Ezquerro MC, Guerrero-Guerra C, Galán C, et al. Pollen in the atmosphere of Mexico City and its impact on the health of the pediatric population. *Atmos Environ*. 2018;186:198–208. doi:10.1016/j.atmosenv.2018.05.006
27. Gastaminza G, Lombardero M, Bernaola G, et al. Allergenicity and cross-reactivity of pine pollen. *Clin Exp Allergy*. 2009;39(9):1438–1446. doi:10.1111/J.1365-2222.2009.03308.X
28. Domínguez-Ortega J, López-Matas MÁ, Alonso MD, et al. Prevalence of allergic sensitization to conifer pollen in a high cypress exposure area. *Allergy Rhinol*. 2016;7(4):200–206. doi:10.2500/AR.2016.7.0183
29. Pollen.com. National allergy forecast & info about allergies; 2022. Available from: <https://www.pollen.com/>. Accessed February 9, 2022.
30. Pfaar O, Bastl K, Berger U, et al. Defining pollen exposure times for clinical trials of allergen immunotherapy for pollen-induced rhinoconjunctivitis – an EAACI position paper. *Allergy Eur J Allergy Clin Immunol*. 2017;72(5):713–722. doi:10.1111/all.13092
31. Bousquet J, Cour P, Guerin B, Michel FB. Allergy in the mediterranean area I. pollen counts and pollinosis of montpellier. *Clin Exp Allergy*. 1984;14(3):249–258. doi:10.1111/j.1365-2222.1984.tb02204.x
32. Buters JTM, Kasche A, Weichenmeier I, et al. Year-to-year variation in release of Bet v 1 allergen from birch pollen: evidence for geographical differences between West and South Germany. *Int Arch Allergy Immunol*. 2008;145(2):122–130. doi:10.1159/000108137
33. Emberlin J, Smith M, Close R, Adams-Groom B. Changes in the pollen seasons of the early flowering trees *Alnus* spp. and *Corylus* spp. in Worcester, United Kingdom, 1996–2005. *Int J Biometeorol*. 2007;51(3):181–191. doi:10.1007/S00484-006-0059-2
34. Kubik-Komar A, Piotrowska-Weryszko K, Weryszko-Chmielewska E, Kaszewski BM. Analysis of *Fraxinus* pollen seasons and forecast models based on meteorological factors. *Ann Agric Environ Med*. 2018;25(2):285–291. doi:10.26444/AAEM/80909
35. Teranishi H, Kenda Y, Katoh T, Kasuya M, Oura E, Taira H. Possible role of climate change in the pollen scatter of Japanese cedar *Cryptomeria japonica* in Japan. *Clim Res*. 2000;14(1):65–70. doi:10.3354/cr014065
36. Parmenter RR, Zlotin RI, Moore DI, Myers OB. Environmental and endogenous drivers of tree mast production and synchrony in piñon–juniper–oak woodlands of New Mexico. *Ecosphere*. 2018;9(8):e02360. doi:10.1002/ECS2.2360
37. Frei T, Gassner E. Trends in prevalence of allergic rhinitis and correlation with pollen counts in Switzerland. *Int J Biometeorol*. 2008;52(8):841–847. doi:10.1007/S00484-008-0178-Z
38. Shida T, Akiyama K, Hasegawa M, et al. Change in skin reactivity to common allergens in allergic patients over a 30-year period. Association with aeroallergen load. *Aerugi*. 2000;49(11):1074–1086. doi:10.15036/arerugi.49.1074
39. Weiland SK, Von Mutius E, Hirsch T, et al. Prevalence of respiratory and atopic disorders among children in the East and West of Germany five years after unification. *Eur Respir J*. 1999;14(4):862–870. doi:10.1034/J.1399-3003.1999.14D23.X
40. Oduber F, Calvo AI, Blanco-Alegre C, et al. Links between recent trends in airborne pollen concentration, meteorological parameters and air pollutants. *Agric for Meteorol*. 2019;264:16–26. doi:10.1016/J.AGRFORMET.2018.09.023
41. Zhao F, Elkesh A, Durner J, et al. Common ragweed (*Ambrosia artemisiifolia* L.): allergenicity and molecular characterization of pollen after plant exposure to elevated NO₂. *Plant Cell Environ*. 2016;39(1):147–164. doi:10.1111/PCE.12601
42. Ouyang Y, Xu Z, Fan E, Li Y, Zhang L. Effect of nitrogen dioxide and sulfur dioxide on viability and morphology of oak pollen. *Int Forum Allergy Rhinol*. 2016;6(1):95–100. doi:10.1002/ALR.21632
43. Urashima M, Asaka D, Endo T, et al. Japanese cedar pollinosis in Tokyo residents born after massive national afforestation policy. *Allergy*. 2018;73(12):2395–2397. doi:10.1111/ALL.13575
44. Cámara de diputados del H. Ley general de desarrollo forestal sustentable. Diario Oficial de la Federación; 2018. Available from: https://www.gob.mx/cms/uploads/attachment/file/493529/ley_general_de_desarrollo_forestal_sustentable.pdf. Accessed February 16, 2022.
45. Rodríguez-Acosta M, Coombes AJ. Manual para la propagación de *Quercus*: una guía fácil y rápida para cultivar encinos en México y América Central; 2020:79. Available from: <https://www.bgci.org/wp/wp-content/uploads/2021/01/manual-para-la-propagacion-de-quercus.pdf>. Accessed February 16, 2022.
46. CONABIO. Reforestación | biodiversidad Mexicana; 2020. Available from: <https://www.biodiversidad.gob.mx/diversidad/reforestacion>. Accessed February 16, 2022.
47. Ariano R, Berra D, Chiodini E, et al. Ragweed allergy: pollen count and sensitization and allergy prevalence in Two Italian allergy centers. *Allergy Rhinol*. 2015;6(3):177–183. doi:10.2500/ar.2015.6.0141
48. Dirección de Monitoreo Atmosférico. Informes anuales de calidad del aire; 2018. Available from: <http://www.aire.cdmx.gob.mx/default.php?opc=z6bhmi>. Accessed February 9, 2022.

49. Teran LM, Montes-Vizuet R, Li X, Franz T. Respiratory proteomics: from descriptive studies to personalized medicine. *J Proteome Res.* 2015;14(1):38–50. doi:10.1021/PR500935S
50. Lambert KA, Lodge C, Lowe AJ, et al. Pollen exposure at birth and adolescent lung function, and modification by residential greenness. *Allergy Eur J Allergy Clin Immunol.* 2019;74(10):1977–1984. doi:10.1111/all.13803
51. Losappio L, Heffler E, Contento F, Cannito C, Rolla G. Thunderstorm-related asthma epidemic owing to *Olea Europaea* pollen sensitization. *Allergy Eur J Allergy Clin Immunol.* 2011;66(11):1508–1510. doi:10.1111/j.1398-9995.2011.02698.x
52. Darrow LA, Hess J, Rogers CA, et al. Ambient pollen concentrations and emergency department visits for asthma and wheeze. *J Allergy Clin Immunol.* 2012;130(3):630–638.e4. doi:10.1016/j.jaci.2012.06.020

Journal of Asthma and Allergy

Dovepress

Publish your work in this journal

The Journal of Asthma and Allergy is an international, peer-reviewed open-access journal publishing original research, reports, editorials and commentaries on the following topics: Asthma; Pulmonary physiology; Asthma related clinical health; Clinical immunology and the immunological basis of disease; Pharmacological interventions and new therapies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-asthma-and-allergy-journal>