

# Migration dynamics of an important rice pest: The brown planthopper (*Nilaparvata lugens*) across Asia—Insights from population genomics

James P. Hereward<sup>1</sup>  | Xuhong Cai<sup>2</sup> | Ambrocio Melvin A. Matias<sup>1,3</sup>  |  
Gimme H. Walter<sup>1</sup>  | Chenxi Xu<sup>2</sup> | Yongmo Wang<sup>2</sup> 

<sup>1</sup>School of Biological Sciences, The University of Queensland, Brisbane, Queensland, Australia

<sup>2</sup>Hubei Insect Resources Utilization and Sustainable Pest Management Key Laboratory, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, Hubei, China

<sup>3</sup>Institute of Biology, University of the Philippines Diliman, Quezon City, Philippines

## Correspondence

Yongmo Wang, College of Plant Science and Technology, Huazhong Agricultural University, Hubei Insect Resources Utilization and Sustainable Pest Management Key Laboratory, Wuhan, Hubei, China.  
Email: ymwang@mail.hzau.edu.cn

## Funding information

This work was supported by the National Key R&D Program of China (2017YFD0200602), the National Natural Science Foundation of China (31101430) and the Fundamental Research Funds for the Central Universities of China (2662017JC006).

## Abstract

**Introduction:** Brown planthoppers (*Nilaparvata lugens*) are the most serious insect pests of rice, one of the world's most important staple crops. They reproduce year-round in the tropical parts of their distribution, but cannot overwinter in the temperate areas where they occur, and invade seasonally from elsewhere. Decades of research have not revealed their source unambiguously.

**Methods and Results:** We sequenced the genomes of brown planthopper populations from across temperate and tropical parts of their distribution and show that the Indochinese peninsula is the major source of migration into temperate China. The Philippines, once considered a key source, is not significant, with little evidence for their migration into China. We find support for immigration from the west of China contributing to these regional dynamics.

**Discussion:** The lack of connectivity between the Philippine population and the mainland Chinese populations explains the different evolution of Imidacloprid resistance in these populations. This study highlights the promise of whole-genome sequence data to understand migration when gene flow is high—a situation that has been difficult to resolve using traditional genetic markers.

## KEYWORDS

agrochemicals, biochemical phenomena, biological evolution, genomics, insect control, insecta

## 1 | INTRODUCTION

Understanding the migration dynamics and connectivity of populations is crucial to setting management priorities for pest insects. This is due to the central role population connectivity plays in the evolution of traits such as insecticide resistance and biological traits that enhance the pest status of an insect, such as the ability to overcome crops bred to be resistant to insect damage (Hendry

et al., 2011). Migration is often the pathway by which insecticide resistance spreads (Denholm, Devine, & Williamson, 2002; Pasteur & Raymond, 1996; Raymond, Callaghan, Fort, & Pasteur, 1991). Genetics provides a way to assess migration dynamics in pest insects that cannot be tracked over large distances, but this has been difficult in the past when rates of gene flow (migration) between populations are high. The use of only a few genetic markers has often failed to provide the necessary resolution in high gene flow systems

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Evolutionary Applications* published by John Wiley & Sons Ltd

(Reitzel, Herrera, Layden, Martindale, & Shank, 2013). Perhaps the ability to generate whole-genome sequence data can provide the necessary resolution in these high gene flow pest systems.


Rice is one of the world's central food crops and is essential to global food security (Long-ping, 2014). One of its most serious threats, the brown planthopper (*Nilaparvata lugens*) (Stål) (Hemiptera: Delphacidae) (Matteson, 2000; Sogawa & Cheng, 1979), physically damages rice plants with its phloem feeding and also causes indirect damage through virus transmission (Nault & Ammar, 1989). Despite an extraordinary amount of research, the migratory routes and population connectivity of this pest remain poorly understood (Bao, Cheng, Cheng, Chu, & Shen, 2000; Cheng et al., 1979; Dung, 1981; Jiang, Tan, Shen, Cheng, & Chen, 1982; Kisimoto, 1976; Kisimoto & Sogawa, 1995; Riley et al., 1991; Rosenberg & Magor, 1983; Seino, Shiotsuki, Oya, & Hirai, 1987; Sogawa & Cheng, 1979). A clear understanding of these aspects of migration is essential to managing this important pest species effectively and has been the core focus of decades of research. The general consensus is that *N. lugens* travels from the tropical part of their distribution into the northern regions of Vietnam and the southern, central and eastern parts of China, migration from these areas follows, apparently associated with the spring monsoon into south Korea and Japan; in addition, a reverse migration to the tropics, with the autumn monsoon, has been proposed (Cheng et al., 1979; GAABPG, 1979; Jiang et al., 1982; Kisimoto, 1976). The primary aim of this study was to address this important question using whole-genome population genetics and to determine the connectivity of potential southern sources and Chinese populations of brown planthoppers.

Brown planthoppers are present throughout the year in the tropical parts of their geographical distribution, which includes the Indochinese peninsula, South-East Asian islands and parts

of India (Figure 1). Here, conditions are ideal for *N. lugens*, where they generally complete one generation every 30 days (Li, 1984; Zhu et al., 2000), resulting in 12 to 13 generations per year. These bugs cannot, however, overwinter on rice in the northern parts of their distribution where the coldest monthly temperature is below 12°C, so seasonal appearances in their northern range are a result of migration (Cheng et al., 1979). Migration of brown planthoppers commonly starts in March, culminating in a peak immigration into the Yangtze river basin in June to July, where the insects may persist for up to five generations (Cheng et al., 1979; Ding, 1979; Hu et al., 2014).

Numerous studies have investigated seasonal long-distance movement of this species through an extensive assortment of techniques: catching individuals at sea, in the air with airplanes and on mountaintops (Cheng et al., 1979; Dung, 1981; Kisimoto, 1976; NCRGBP, 1981). Meteorological analyses, radar observations and mark-recapture experiments have also been conducted (Riley et al., 1991; Rosenberg & Magor, 1983; Seino et al., 1987). Based on these studies, there is a widely held belief that migration occurs from southern parts of their distribution to the Chinese mainland, South Korea and Japan (Bao et al., 2000; Kisimoto & Sogawa, 1995; Sogawa & Cheng, 1979).

Three putative sources of planthopper migration into China, South Korea and Japan, have been identified, and these are not seen as mutually exclusive. The Indochinese peninsula is generally considered the primary source (Otuka, Matsumura, Watanabe, & Dinh, 2008; Wu, Ping, & Yong, 1997; Zhu, Lu, Hang, & Xu, 1994). In this region, the Red River and Mekong River Deltas are planted extensively to rice. The Mekong Delta alone contains roughly 10,000 square kilometres of rice and supports up to three crops annually (Nguyen, 2007). The smaller Red River Delta is limited to



**FIGURE 1** Map showing the geographical range of brown planthoppers and the extent of the overwintering and nonoverwintering parts of its range (see key, bottom left). Populations sampled in this study are shown, and the size of the circles represents the number of population-specific (private) alleles in each population

two crops per year, presumably supporting fewer brown planthoppers. A putative migration route has origins in the Mekong River Delta, moving to the Red River Delta in February and March, and then to south China, east China and Korea and Japan in the months that follow (Wu et al., 1997). However, independent analyses of the seasonal monsoon suggest that brown planthopper populations in the Mekong River Delta cannot migrate into the Red River Delta and then to China because the seasonal winds in February to April are not suitable (Hu, Lu, Tuan, & Liu, 2017; Zhai & Cheng, 2006).

The second putative source is the Philippines where rice is also the main food crop. Weather simulations indicate that brown planthoppers could migrate from the Philippines into Taiwan and Japan, but also to southern China, sometimes in association with typhoons (Otuka et al., 2005). The Philippines populations possibly mix with Indochinese populations in southern China before arriving in Japan (Otuka et al., 2005; Otuka et al., 2012), but this has not been tested further. Various features of *N. lugens* from the Philippines suggest that they are different from the mainland populations. They are nearly one hundred times more susceptible to the pesticide imidacloprid than populations in Taiwan, northern Vietnam, China and Japan (Matsumura et al., 2008). In addition, the Philippines and other east Asian populations of brown planthopper differ from one another in their ability to overcome planthopper-resistant rice varieties (Sogawa, 1992).

The third putative source is Myanmar and Bangladesh, but these bugs are likely to extend only into the west of Yunnan province (China) and it is thought that they would not be able to migrate into central and eastern China because of mountainous barriers and unsuitable monsoon conditions (Wu et al., 1997). These three putative sources of brown planthopper migration have found varying levels of support from empirical studies, but have not been tested directly.

Studies based on mitochondrial DNA (mtDNA) indicated high gene flow across the distribution of brown planthopper (Mun, Song, Heong, & Roderick, 1999). One haplotype was found in every country sampled across South and East Asia, and since only three haplotypes were recovered from the 71 individuals investigated, little resolution of the migration dynamics was possible. Understanding population connectivity in this pest is key to understanding how undesirable traits like insecticide resistance will spread. The objective of this study was to investigate population connectivity of populations of brown planthopper at a broad geographical scale, throughout China and South-East Asian, to better understand the migration routes of this key insect pest of rice.

## 2 | MATERIALS AND METHODS

### 2.1 | Sample collection and sequencing

*Nilaparvata lugens* adults were collected in rice fields from 10 locations in China, Vietnam, Thailand and Philippines from April to August 2010 (Table S1). The population from Los Baños, Philippines, was collected in 2009. Samples were collected directly into 95% ethanol

and then stored at  $-20^{\circ}\text{C}$ , prior to DNA extraction. In Tengchong, Nanjing (Fujian province) and Changde (all China), brown planthoppers cannot overwinter at any life stage of development. In Menghai, Wenshan and Longzhou (also all in mainland China), they can sometimes be found in rice volunteers in winter. In the tropics (Vietnam, Thailand and Philippines), they survive throughout the year.

DNA was extracted from six individuals from each population using TaKaRa MiniBEST Universal Genomic DNA extraction Kit Ver.5.0 (U-Me Biotech, Wuhan). Library preparation and sequencing was conducted by Total Genomics Solution (Shenzhen, China). Sequencing was performed on Illumina HiSeq X Ten returning 150 bp paired end reads with an insert size of approximately 280 bp. Sequence data quality metrics were estimated with Fastqc (Andrews, 2018) and Fastqsceen (Winget, 2018). The sequence data of three of the individuals from Vietnam did not contain sufficient data that mapped to brown planthopper, after quality filtering, and were discarded from further analysis. Six additional individuals from that site (Vietnam) were sequenced with BGI genomics (Shenzhen, China) with the same insert size, read length, library preparation and sequencing technology. We discarded the three Vietnamese samples that failed quality control but kept the three from the original sequencing that had passed, resulting in nine individuals from this population. All other samples had high-quality sequences and we used all six individuals in downstream analysis, resulting in full genome sequences of 63 individuals in total. We removed adaptors and quality-trimmed the end of the sequences of all samples to q10 using bbdduk from the bbtools package (version 36), with a kmer of 8 (Bushnell, 2018).

### 2.2 | Data assembly

We mapped all reads to the brown planthopper reference genome, assembly GCA\_000757685.1 of bioproject PRJNA177647 in GenBank (NCBI), which is approximately 1.2 Gbp (Xue et al., 2014), using BWA mem v0.7 (Li & Durbin, 2009). We followed GATK (Genome Analysis Tool Kit) best practices (Van der Auwera et al., 2013) and called variants using HaplotypeCaller (gatk v 4.0.10.1), and this was made parallel by splitting the reference into windows, each containing complete contigs. For each window, we called individual g.vcf files and later combined these and performed joint genotyping using GenotypeGVFs (gatk v 4.0.10.1).

We filtered *snps* (single nucleotide polymorphisms) to quality Q30 using vcftools, then to a minimum depth of three (as studies have showed that with appropriate filtering genotypes can be called from as few as three sequence reads) and a maximum depth of 30 (three times our highest coverages). Following this, we removed any marker that was missing more than 30% of the data (30% of individuals were not genotyped), to remove the markers most affected by missing data. We then removed any locus with a minor allele count below three (e.g. one homozygote and one heterozygote). Applying hard filters for minor allele frequency can cause weak structure to be missed (Linck & Battey, 2019), but the vast majority of singletons are

errors, and also confound analyses. We used the minor allele count (=3) instead, because it is the most conservative way of removing singletons, but not rare alleles. We repeat-masked the reference genome using RepeatMasker 4.0.8 using the Dfam and RepBase databases—the repeat-masked reference was converted to a bed file, and this was used to remove *snps* in repetitive regions of the genome with vcftools (Danecek et al., 2011). We then created a bed file with all the contigs below 5 kbp and removed all of these *snps* in the same way, assuming that some of the smaller contigs were likely to be less reliable. All indels and nonbiallelic *snps* were removed leaving only biallelic *snps*, due to the requirements of downstream analyses. We filtered out markers that were out of Hardy–Weinberg equilibrium because most of these are likely errors, and we were interested in the neutral process of migration, rather than investigating any effects of selection. Deviations from HWE were calculated separately using vcftools in the tropical populations Philippines, and Indochinese Peninsular (Thailand and Vietnam combined, see below), and then, all markers with deviations from HWE in these populations were removed from the total data set (using a threshold of  $p = .05$ ). This final data set contained 762,576 *snps* and is referred to as the *snp* genotype data. An alternative approach to calling *snps* and filtering them is to infer genotype likelihoods, with some analyses methods then being able to use the likelihoods rather than the raw genotypes themselves, preserving the uncertainty in the inferences. We also called genotype likelihoods with the program ANGSD (Korneliussen, Albrechtsen, & Nielsen, 2014), using the Samtools likelihood model; this data set is referred to as the genotype likelihood data.

## 2.3 | Mitogenome analysis

We mapped the sequence data to the mitochondrial genome of *N. lugens* using BWA (Burrows–Wheeler Aligner, Li & Durbin, 2009). There are six mitochondrial genomes for brown planthopper on GenBank; however, only one has the control region and repeats (JX880069). This mitogenome also had a large insertion between *ND2* (NADH dehydrogenase two) and *COI* (cytochrome oxidase subunit one) that was not supported by read mapping, so we removed this region (based on alignment to the other five mitogenomes) and used this edited sequence as the reference. We used a pipeline to call *snps* and prepared a consensus sequence for each individual. Briefly, after mapping, we removed the unmapped reads using Samtools (Li et al., 2009), sorted, marked duplicates and indexed the bam file using GATK4 (Van der Auwera et al., 2013), called *snps* using HaplotypeCaller from GATK4 in ploidy 1 mode, filtered the vcf file to q30 with vcftools (Danecek et al., 2011), made a consensus sequence with bcftools (Li et al., 2009), masked regions with no mapping using bedtools (Quinlan & Hall, 2010) and then renamed the fasta file with the name of the sequenced individual in a perl wrapper script.

We manually checked the resulting consensus sequences by mapping back the raw reads, and had to manually edit a small portion of the *ND2* gene because read mapping in that region indicated

some kind of duplication, so we edited based on the reads that did not disrupt the coding sequence of the gene. The mapping was also unreliable around the repetitive control region in the reference, so we deleted this from the overall alignment. Sequences were aligned with MAFFT (Katoh & Standley, 2013), and we made a TCS haplotype network with Popart (Leigh & Bryant, 2015).

## 2.4 | Genetic structure

The distribution of brown planthoppers was plotted on a map, and areas with mean monthly temperature of the coldest month below 13.5°C were designated as nonoverwintering, between 13.5°C and 15°C as intermediate, and over 15°C as overwintering. The temperature data were obtained from WorldClim global climate layers (<https://www.worldclim.org/>) accessed using the R (R Core Team, 2017) package raster.

We first examined the overall population structure by calculating  $F_{ST}$  (Weir & Cockerham, 1984) using the full *snp* genotype data described above. Overall  $F_{ST}$  was calculated in hierfstat (Goudet, 2004), as were pairwise  $F_{ST}$ s, performing 1,000 bootstrap replicates. We used the genotype likelihoods to infer admixture proportions using NGSadmix (Skotte, Korneliussen, & Albrechtsen, 2013), which is designed to infer admixture proportions in low coverage sequencing data.

A discriminant analysis of principal components (DAPC) (Jombart, Devillard, & Balloux, 2010) was performed on the full *snp* genotype data to assess the genetic relationship between the populations. DAPC maximizes differences between defined populations across multiple principle components. Cross-validation indicated that 30 components should be retained. The DAPC analysis was performed using the adegenet package in R (Jombart, 2008).

To assess population structure without predefining populations, we performed a principal component analysis (PCA) on the full *snp* genotype data. Principal component analysis cannot be performed on data with missing values, so for each population we replaced missing data with the population-specific mode using custom R scripts (see data availability).

## 2.5 | Assignment testing

We used assignment testing to infer the source of the immigrants into the nonoverwintering locations. Based on the PCA and admixture analyses, we designated the “core-overwintering” populations (including Tengchong), which we assume to be the putative sources, into three groups: Philippines, Tengchong and Indochinese peninsular. Indochinese peninsular comprised Thailand and Vietnam. We generated linear discriminant functions based on these three groups, using DPAC (Jombart et al., 2010) and retaining 10 axes. We then assigned individuals from the nonoverwintering sites to these three groups using DPAC (Jombart et al., 2010).

## 2.6 | Genetic diversity and population-specific alleles

The number of population-specific alleles in each population was calculated using a custom R script, which calculated allele frequency and then identified *snps* that had a value other than zero in only one population, and these were represented as the size of the circle for each population on the map. We then tested whether the population-specific alleles were driving genetic structure by running PCA on the full *snp* data and on a data set with all population-specific (private) alleles removed.

The genetic diversity summary statistics  $\pi$  and  $\theta$  were calculated from the genotype likelihood data for every 10 kb sliding window with 2.5 kb overlap across the genome in ANGSD (Korneliusson et al., 2014). This analysis aimed to capture the genomic variance—that is the variation in patterns across contigs or chromosomes. These data were subsampled by selecting one out of every three windows. Windows that were not available across all 10 populations were then discarded. Windows in which the effective number of sites was less than 100 bp were also removed. The remaining windows were then used to plot the genomic diversity. There was no variation across populations when considering the raw distributions. The estimate for each window was standardized by the mean and standard deviation across the population ( $(x - \bar{x}) / (sd(x))$ ), with the direction (sign) of the numerator indicating higher or lower genomic diversity.


## 3 | RESULTS

The mitogenome analysis indicated little spatial structuring of haplotypes. We recovered 53 unique haplotypes from the 63 different individuals, based on 296 variable sites in the 14,380bp mitogenome alignment. Of the few shared haplotypes, one was found in Wenshan, Menghai and the Philippines, and another in Menghai, Longzhou and Thailand (Figure 2). When we restricted the mtDNA analysis to COI, the same major haplotype was present across


all populations, but several unique haplotypes were found in the Philippines (Figure 2).

We achieved a mean coverage of 10.5× across the filtered *snps* and around 6× coverage across the whole genome in our initial mapping (Table S2). Genetic differentiation was low across all populations, with an overall  $F_{ST}$  of 0.0093 ( $\pm$  95% CI 0.0092–0.0094). The highest pairwise  $F_{ST}$ s were from comparisons of the Philippines to other populations (0.024, Table S3), and then from comparisons of Tengchong to other populations (0.015, Table S3). Despite the low overall genetic differentiation, whole-genome sequence data revealed major insights into the migration dynamics of brown planthopper. Admixture analysis (NGSadmix), DAPC and the PCA indicated that the Philippines insects represent a distinct population with very little evidence of admixture from the Philippines into mainland populations (Figure 3, Figure 4). Tengchong, in the far west of China, was also separated from other populations in the admixture analysis, with evidence of admixture across the rest of the mainland sites (Figure 3). The PCA separated some individuals from the Tengchong population, but others were placed in the main cluster, indicating a mixture between common genotypes and an unidentified source population.

The Philippines and Tengchong samples had more unique, population-specific alleles than the other populations (Figure 1), even though the PCA indicated that Tengchong is a mixed population (Figure 4). This likely indicates a distinct source of genetic diversity (presumably outside the geographical scope of this study) that contributes to the migration into Tengchong. We tested whether the population-specific alleles were driving our inferences of population structure by removing all of them and conducting a separate PCA analysis. The PCA remained qualitatively the same when all population-specific alleles had been removed from the data set (Figure 4, bottom), indicating that allele frequencies, rather than population-specific alleles, are driving our inference of genetic structure. The higher number of population-specific alleles in these two populations was not the result of higher overall genomic diversity as these two populations had lower genomic diversity than other populations (Figure 5). The nonoverwintering sites (Changde and Nanjing) did not



**FIGURE 2** Mitochondrial genome analyses. Haplotype networks based on the complete mitochondrial genome (16,794 bp, a), and the COI gene only (2,795 bp, b) the size of each circle represents the number of individuals that have the same sequence. For localities, see Figure 1



**FIGURE 3** (a) Discriminant analysis of principal components (DAPC) of the *snp* genotype data maximizing genetic differences across defined populations. (b) Admixture analysis based on the genotype likelihood data using NGSadmix to infer admixture proportions. Each bar is one individual, and the colours represent the posterior probability of assignment to each of three putative “population” clusters

have lower genomic diversity than the sites in tropical and subtropical regions that have year-round production of brown planthoppers (Figure 5).

All individuals from the nonoverwintering sites were assigned to the Indochinese peninsular group (comprising Thailand and Vietnam) at close to 100% probability. When we plotted the linear discriminant scores for the nonoverwintering individuals and the three “core-overwintering” groups, the nonoverwintering individuals were placed between the Indochinese peninsular group and Tengchong (but closer to Indochinese peninsular). This is likely a result of the admixture between Tengchong and the other mainland populations (Figure S1).

## 4 | DISCUSSION


Our results show that a population genomics approach contributes considerably to our understanding of the migration dynamics of brown planthoppers. The Indochinese peninsular is indicated as a major source of brown planthoppers in their migration to the temperate regions of China (and most likely on to Japan). The Philippines population was genetically distinct from the other populations in all analyses, and little evidence of gene flow from the Philippines into temperate China was found. Tengchong (in western temperate China) also formed a separate group in the admixture analysis and DAPC, but the PCA indicated a mixed population (Figures 3 and 4). The high prevalence of population-specific alleles in this population likely indicates immigration from more western regions

(potentially Myanmar, Bangladesh and India, which were not sampled in this study). The populations in subtropical China and the tropical Indochinese peninsular show admixture with the Tengchong population, indicating that planthoppers from these putative western sources may contribute to the overall migration dynamics, despite the mountainous barriers and unsuitable monsoon conditions, and contrary to previous assertions (Wu et al., 1997).

Our mitogenome analysis is broadly in line with the results of Mun et al. (1999) who found only three haplotypes, one of which was present across all sites that they sampled in China, the Philippines, Bangladesh, Malaysia, Vietnam, Thailand and Korea. By using the whole mitogenome, we recovered many more haplotypes, but still no geographical clustering of the haplotypes (Figure 2). This indicates no long-term separation of the various populations and is consistent with an interpretation that the insects are highly migratory and populations from all parts of its distribution have been connected through migration and gene flow in the relatively recent past. Leveraging genome-wide nuclear *snp* data, we were, however, able to make inferences into the more recent migratory dynamics of this species (as outlined above and below).

Physiological comparisons of bugs from temperate, subtropical and tropical regions have led to suggestions that tropical populations of brown planthopper do not contribute to the migratory dynamics of this pest (Iwanaga, Nakasuji, & Tojo, 1987; Nagata & Masuda, 1980; Wada, Ito, Takahashi, & Tang, 2007, 2009). Our analysis shows, by contrast, that the tropical populations in the Indochinese peninsular, the subtropical populations in China, and the temperate populations in China are connected genetically, all


**FIGURE 4** Effect of population-specific alleles on genetic structure. (a) Principal components analysis (PCA) of the full *snp* genotype data. (b) PCA of *snp* genotype data with all population-specific alleles removed. The genetic structure was not driven by private alleles



of which indicates that the tropical populations in the Indochinese peninsular (Vietnam and Thailand) are major sources of the migration into China. Planthoppers from these locations essentially form a single panmictic population, as revealed in the PCA, and they were separated only in the DAPC (which maximizes differences across designated “populations”). We expected that the nonoverwintering populations (Changde and Nanjing) might have reduced genetic diversity if they had experienced a founder effect following immigration in the spring. This was not the case (Figure 5), and the same amount of genetic diversity was present in these two populations as in the Indochinese peninsular populations. This indicates that sufficiently large numbers of insects make the northern migration that bottlenecks or founder effects do not occur.

Population genetics approaches to estimating migration rates enable inferences to be made about small insects over the large

distances that separate populations, and that are not possible with ecological approaches to measuring dispersal such as mark-recapture techniques. Our study indicates that using the whole genome provides greater power to make these inferences than previous molecular methods based on relatively few markers (such as the mitochondrial analysis of Mun et al. (1999)), as suggested by Luikart, England, Tallmon, Jordan, and Taberlet (2003). It is not yet clear what the power difference is between empirical microsatellite data sets and whole-genome analyses to detect migration because there are not many data sets to compare. Studies that compare microsatellites and *snps* generally find that moderate numbers of *snps* can detect the same genetic structure as microsatellite data, and that the resolution increases as the number of *snps* increase (Fischer et al., 2017; Gärke et al., 2012). Generating whole-genome sequence data also allows further analysis of



**FIGURE 5** Genomic diversity. Plots of the normalized deviation from mean diversity measures  $\pi$  (a), and  $\theta$  (b), raw histograms shown

endosymbionts and tests of selection in the future, although this was outside the scope of the current study.

This study highlights that it is still difficult to make precise estimates of migration using classic population genetic methods when gene flow is very high (high migration systems like brown planthopper). The number of individuals that could be included in this whole-genome analysis was limited by cost. Increasing the number of individuals per population may have provided greater power in our assignment testing. The real promise of whole-genome sequencing in this kind of study, however, likely lies in the potential to perform not only parentage analysis, but to leverage the information from linkage and recombination to infer deeper pedigree information and estimate migration across multiple generations. This approach would require many individuals to be sampled from each putative population and sequenced with sufficient depth, and is currently cost prohibitive, but will likely become feasible with further advances in sequencing technology.

## 4.1 | Implications for management

The Philippines has been reported as a major source of the northward migration of brown planthoppers into temperate regions of China and Japan, based on trap catches, backwards telemetry and modelling (Otuka et al., 2005; Otuka et al., 2012). In our analysis, the Philippine population of brown planthoppers formed a highly distinct cluster in the admixture analysis (Figure 1) and the PCA (Figure 4). There is very little evidence of admixture of the Philippine population into the temperate or subtropical Chinese populations. This indicates very little gene flow and migration between the Philippine populations and mainland Chinese temperate populations. Our results explain the difference reported between the Philippines and other populations in their resistance to imidacloprid (Matsumura et al., 2008), as well as their relatively different responses to new rice varieties (Sogawa, 1992). Recently, a distinct strain of the planthopper endosymbiont, *Arsenophonus*, has been associated with imidacloprid resistance (Pang et al., 2018). All susceptible insects in that study had a distinct strain of *Arsenophonus*, and when this strain was transplanted into resistant individuals, they became susceptible. This strain is found predominantly in the Philippines, although it was also detected at a lower frequency in the Chinese mainland province of Guangxi (Pang et al., 2018). By showing that the Philippines population is distinct, with little evidence for migration between the Philippines and the Chinese mainland, our data explain the distribution of the susceptible and resistance-related *Arsenophonus* strains, although the presence of a susceptible-related endosymbiont in Guanxi (China) in that study is most likely the result of rare immigration of Philippines planthoppers into mainland China. This highlights the key role of migration in the evolution and spread of insecticide resistance (Denholm et al., 2002; Pasteur & Raymond, 1996; Raymond et al., 1991).

Our results show that the Philippines is distinct, in population genetics terms, from the mainland populations and there is very little migration between them. Insecticide resistance traits that evolve in the Philippines are less likely to spread to the mainland and vice versa, as seems to be the case for imidacloprid (Pang et al., 2018). Populations in mainland South-East Asia and mainland China are highly connected by gene flow, however, and insecticide resistance that evolves in Vietnam or Thailand is highly likely to spread to China, and could quite possibly spread back the other way. This creates a situation where insecticide resistance management has to occur across borders. Attempts to manage insecticide resistance in China that do not consider the populations and insecticide selection pressures south of the border would be futile.

Our data also indicate that there may be a somewhat distinct population to the west of Tengchong, perhaps in India. Future studies should investigate populations further to the west of the insect's distribution, from Japan, and more sites in the South-East Asian part of its distribution to further establish the migratory dynamics of this insect and the routes by which insecticide resistance traits travel.

## 5 | CONCLUSIONS

In summary, we used whole genomes to make new inferences about brown planthopper migration. Chinese and Indochinese peninsular populations should be considered the same in terms of management and insecticide resistance. The Philippines population is distinct from these mainland populations, and this explains the prevalence of imidacloprid susceptibility there compared to elsewhere. Our results also indicate the potential for immigration from the west into the Chinese and Indochinese peninsular populations. This route was previously discounted but may present a means for new insecticide resistance traits to enter the Chinese and Indochinese peninsular populations. More generally, our results highlight the promise that population genomics holds for understanding the connectivity of pest insect populations characterized by high rates of gene flow.

### ACKNOWLEDGMENTS

We thank Dr. Z. LV for providing some insect samples from Philippines.

### CONFLICT OF INTEREST

There are no competing interests to declare.

### AUTHOR CONTRIBUTIONS

J.P.H and A.M.M analysed the data and designed the figures. J.P.H and Y.M.W wrote the manuscript, with revisions and input from A.M.M, and G.H.W. and X.C prepared libraries for sequencing. C.X and Y.M.W collected the samples. Y.M.W designed the research.

### DATA AVAILABILITY STATEMENT

Raw data (Bam files), intermediate data (the vcf file) and sample metadata can be found at UQ espace, Brown planthopper genomics, Hereward, Matias & Wang, (2020). <https://doi.org/10.14264/a371d45>.

### ORCID

James P. Hereward  <https://orcid.org/0000-0002-6468-6342>

Ambrocio Melvin A. Matias  <https://orcid.org/0000-0002-1329-4285>

[org/0000-0002-1329-4285](https://orcid.org/0000-0002-1329-4285)

Gimme H. Walter  <https://orcid.org/0000-0003-1025-7101>

Yongmo Wang  <https://orcid.org/0000-0001-9904-1352>

### REFERENCES

- Andrews, S. (2018). *FastQC: A quality control tool for high throughput sequence data*. [online]. Retrieved from <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Bao, Y. X., Cheng, J. Y., Cheng, X. N., Chu, C. S., & Shen, T. L. (2000). Dynamical and numerical simulations on the processes of *Nilaparvata lugens* long-distance migration northward during the midsummer in China. *Acta Entomologica Sinica*, 43, 176–183. (Chinese with English summary).
- Bushnell, B. (2018). *Bbmap - a suite of bioinformatics tools*. [online]. Retrieved from <https://sourceforge.net/projects/bbmap/>
- Cheng, S. N., Chen, J. C., Si, H., Yan, L. M., Chu, T. L., Wu, C. T., ... Yan, C. S. (1979). Studies on the migrations of brown planthopper *Nilaparvata lugens*. *Acta Entomologica Sinica*, 22, 1–21. (in Chinese with English summary).
- Danecek, P., Auton, A., Abecasis, G., Albers, G. A., Banks, E., DePristo, M. A., ... Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, 27(15), 2156–2158.
- Denholm, I., Devine, G. J., & Williamson, M. S. (2002). Insecticide Resistance on the Move. *Science*, 297(5590), 2222–2223.
- Ding, D. (1979). Temperature and the brown planthopper *Nilaparvata lugens* (Stål) distribution. *Jiangsu Agricultural Sciences*, 7(2), 40–43.
- Dung, W. S. (1981). A general survey on seasonal migrations of *Nilaparvata lugens* (Stal) and *Sogatella furcifera* (Horvath) (Homoptera: Delphacidae) by means of airplane collections. *Acta Phytophylacila Sin*, 8, 74–82.
- Fischer, M. C., Rellstab, C., Leuzinger, M., Roumet, M., Gugerli, F., Shimizu, K. K., Holderegger, R., & Widmer, A. (2017). Estimating genomic diversity and population differentiation - an empirical comparison of microsatellite and SNP variation in *Arabidopsis halleri*. *BMC genomics*, 18(1), 69. <https://doi.org/10.1186/s12864-016-3459-7>
- GAABPG - Guangdong Agricultural Academy Brown Planthopper Group. (1979). Studies on the regularity of migration of the brown planthopper. *Guangdong Nongye Kexue*, 3, 43–48. (in Chinese).
- Gärke, C., Ytournal, F., Bed'hom, B., Gut, I., Lathrop, M., Weigend, S., & Simianer, H. (2012). Comparison of SNPs and microsatellites for assessing the genetic structure of chicken populations. *Animal Genetics*, 43(4), 419–428. <https://doi.org/10.1111/j.1365-2052.2011.02284.x>
- Goudet, J. (2004). Hierfstat, a package for r to compute and test hierarchical F-statistics. *Molecular Ecology Notes*, 5(1), 184–186.
- Hendry, A. P., Kinnison, M. T., Heino, M., Day, T., Smith, T. B., Fitt, G., ... Carroll, S. P. (2011). Evolutionary principles and their practical application. *Evolutionary Applications*, 4(2), 159–183. <https://doi.org/10.1111/j.1752-4571.2010.00165.x>
- Hereward, J., Matias, D., & Wang, Y. (2010). Brown planthopper genomics. The University of Queensland. Data Collection. <https://doi.org/10.14264/a371d45>
- Hu, G., Lu, F., Zhai, B.-P., Lu, M.-H., Liu, W.-C., Zhu, F., ... Zhang, X.-X. (2014). Outbreaks of the Brown Planthopper *Nilaparvata lugens* (Stål) in the Yangtze River Delta: Immigration or Local reproduction? *PLoS One*, 9(2), e88973. <https://doi.org/10.1371/journal.pone.0088973>
- Hu, G., Lu, M. H., Tuan, H. A., & Liu, W. C. (2017). Population dynamics of rice planthoppers, *Nilaparvata lugens* and *Sogatella furcifera* (Hemiptera, Delphacidae) in Central Vietnam and its effects on their spring migration to China. *Bulletin of Entomological Research*, 107(3), 369–381.
- Iwanaga, K., Nakasuji, F., & Tojo, S. (1987). Wing polymorphism in Japanese and foreign strains of the brown planthopper, *Nilaparvata lugens*. *Entomologia Experimentalis Et Applicata*, 43, 3–10. <https://doi.org/10.1111/j.1570-7458.1987.tb02194.x>
- Jiang, G. H., Tan, H. Q., Shen, W. Z., Cheng, X. N., & Chen, R. C. (1982). The relation between long-distance southward migration of the brown planthopper (*Nilaparvata lugens* Stål.) on synoptic weather conditions. *Acta Entomologica Sinica*, 25, 147–155. (in Chinese).
- Jombart, T. (2008). adegenet: An R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24, 1403–1405.
- Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components: A new method for the analysis of genetically structured populations. *BMC Genetics*, 11(1), 94.
- Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, 30(4), 772–780.
- Kisimoto, R. (1976). Synoptic weather conditions inducing long-distance immigration of the planthopper. *Ecological Entomology*, 1(2), 95–109.

- Kisimoto, R., & Sogawa, K. (1995). Migration of the brown planthopper, *Nilaparvata lugens* and the white-backed planthopper *Sogatella furcifera* in East Asia: The role of weather and climate. In V. A. Drake, & A. G. Gatehouse (Eds.), *Insect migration: Tracking resources through space and time* (pp. 67–91). Cambridge, UK: Cambridge University Press.
- Korneliusson, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics*, 15, 1471–2105.
- Leigh, J. W., & Bryant, D. (2015). POPART: Full-feature software for haplotype network construction. *Methods in Ecology and Evolution*, 6(9), 1110–1116.
- Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 25(14), 1754–1760.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*, 25(16), 2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>
- Li, R. (1984). Population growth of the brown planthopper, *Nilaparvata lugens* Stål, as influenced by temperature. *Acta Phytophylacica Sinica*, 11(2), 101–107.
- Linck, E., & Battey, C. J. (2019). Minor allele frequency thresholds strongly affect population structure inference with genomic data sets. *Molecular Ecology Resources*, 19(3), 639–647. <https://doi.org/10.1111/1755-0998.12995>
- Long-ping, Y. (2014). Development of hybrid rice to ensure food security. *Rice Science*, 21(1), 1–2. [https://doi.org/10.1016/S1672-6308\(13\)60167-5](https://doi.org/10.1016/S1672-6308(13)60167-5)
- Luikart, G., England, P. R., Tallmon, D., Jordan, S., & Taberlet, P. (2003). The power and promise of population genomics: From genotyping to genome typing. *Nature Reviews Genetics*, 4(12), 981–994. <https://doi.org/10.1038/nrg1226>
- Matsumura, M., Takeuchi, H., Satoh, M., Sanada-Morimura, S., Otuka, A., Watanabe, T., & Van Dinh, T. (2008). Species-specific insecticide resistance to imidacloprid and fipronil in the rice planthoppers *Nilaparvata lugens* and *Sogatella furcifera* in East and South-east Asia. *Pest Management Science*, 64, 1115–1121.
- Matteson, P. C. (2000). Insect pest management in tropical Asian irrigated rice. *Annual Review of Entomology*, 45, 549–574. <https://doi.org/10.1146/annurev.ento.45.1.549>
- Mun, J. H., Song, Y. H., Heong, K. L., & Roderick, G. K. (1999). Genetic variation among Asian populations of rice planthoppers, *Nilaparvata lugens* and *Sogatella furcifera* (Hemiptera: Delphacida): Mitochondrial DNA sequences. *Bulletin of Entomological Research*, 89, 245–253.
- Nagata, T., & Masuda, T. (1980). Insecticide susceptibility and wing-form ratio of the brown planthopper, *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae) and the whitebacked planthopper, *Sogatella furcifera* (Horvath) (Hemiptera: Delphacidae) of Southeast Asia. *Applied Entomology and Zoology*, 15, 10–19.
- Nault, L. R., & Ammar, E. D. (1989). Leafhopper and planthopper transmission of plant viruses. *Annual Review of Entomology*, 34, 503–529.
- NCRGBP - National Cooperative Research Group of Brown Planthopper. (1981). The migration research of brown planthopper *Nilaparvata lugens* (Stål). *Chinese Journal of Agrometeorology, Scientific Agricultural Sinica*, 21(2), 52–59.
- Nguyen, H. N. (2007). Flooding in Mekong River Delta, Viet Nam. In UNDP. (Eds.), *Human development report 2007/2008—Fighting climate change: Human solidarity in a divided world, Ha Noi*. UNDP.
- Otuka, A., Matsumura, M., Watanabe, T., & Dinh, V. T. (2008). A migration analysis for rice planthoppers, *Sogatella furcifera* (Horvath) and *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae), emigrating from northern Vietnam from April to May. *Applied Entomology and Zoology*, 43, 527–534.
- Otuka, A., Watanabe, T., Suzuki, Y., Matsumura, M., Furuno, A., & Chino, M. (2005). A migration analysis of the rice planthopper *Nilaparvata lugens*, from the Philippines to East Asia with three-dimensional computer simulations. *Population Ecology*, 47(2), 143–150.
- Otuka, A., Zhou, Y., Lee, G. S., Matsumura, M., Zhu, Y., Park, H. H., ... Sanade-Morimura, S. (2012). Prediction of overseas migration of the small brown planthopper, *Laodelphax striatellus* (Hemiptera: Delphacidae) in East Asia. *Applied Entomology & Zoology*, 47(4), 379–388.
- Pang, R., Chen, M., Yue, L., Xing, K., Li, T., Kang, K., ... Zhang, W. (2018). A distinct strain of *Arsenophonus* symbiont decreases insecticide resistance in its insect host. *PLoS Genetics*, 14(10), e1007725. <https://doi.org/10.1371/journal.pgen.1007725>
- Pasteur, N., & Raymond, M. (1996). Insecticide resistance genes in mosquitoes: Their mutations, migration, and selection in field populations. *The Journal of Heredity*, 87(6), 444–449. <https://doi.org/10.1093/oxfordjournals.jhered.a023035>
- Quinlan, A., & Hall, M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, 26(6), 841–842.
- R Core Team. (2017). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Raymond, M., Callaghan, A., Fort, P., & Pasteur, N. (1991). Worldwide migration of amplified insecticide resistance genes in mosquitoes. *Nature*, 350(6314), 151–153. <https://doi.org/10.1038/350151a0>
- Reitzel, A. M., Herrera, S., Layden, M. J., Martindale, M. Q., & Shank, T. M. (2013). Going where traditional markers have not gone before: Utility of and promise for rad sequencing in marine invertebrate phylogeography and population genomics. *Molecular Ecology*, 22(11), 2953–2970. <https://doi.org/10.1111/mec.12228>
- Riley, J. R., Cheng, X. N., Zhang, X. X., Reynolds, D. R., Xu, G. M., Smith, A. D., ... Zhai, B. P. (1991). The long-distance migration of *Nilaparvata lugens* (Stål.) (Delphacidae) in China: Radar observations of mass return flight in the autumn. *Ecological Entomology*, 16, 471–489.
- Rosenberg, L. J., & Magor, J. I. (1983). Flight duration of the brown planthopper, *Nilaparvata lugens* (Homoptera: Delphacidae). *Ecological Entomology*, 8, 341–350.
- Seino, H., Shiotsuki, Y., Oya, S., & Hirai, Y. (1987). Prediction of long-distance migration of rice planthoppers to Northern Kyushu considering low-level jet stream. *Journal of Agricultural Meteorology*, 43, 203–208.
- Skotte, L., Korneliusson, T. S., & Albrechtsen, A. (2013). Estimating individual admixture proportions from next generation sequencing data. *Genetics*, 195(3), 693–702.
- Sogawa, K. (1992). A change in biotype property of brown planthopper populations immigrating into Japan and their probable source area. *Proceedings of the Association for Plant Protection of Kyushu*, 38, 63–68. (Japanese with English summary).
- Sogawa, K., & Cheng, C. H. (1979). Economic thresholds, nature of damage, and losses caused by the brown planthopper. *Brown Planthopper: Threat to rice production in asia* (pp. 125–142). Los Banos, Laguna, Philippines: The International Rice Research Institute.
- Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine, A., ... DePristo, M. A. (2013). From FastQ data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. *Current Protocols in Bioinformatics*, 43(1), 11.10.1–11.10.33.
- Wada, T., Ito, K., Takahashi, A., & Tang, J. (2007). Variation of pre-oviposition period in the brown planthopper, *Nilaparvata lugens*, collected in tropical, subtropical and temperate Asia. *Journal of Applied Entomology*, 131, 698–703.
- Wada, T., Ito, K., Takahashi, A., & Tang, J. (2009). Starvation tolerance of macropter brown planthopper, *Nilaparvata lugens*, from temperate, subtropical, and tropical populations in East and South-East Asia. *Entomologia Experimentalis Et Applicata*, 130, 73–80.

- Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analyses of population structure. *Evolution*, 38(6), 1358–1370.
- Winget, S. (2018). *Fastqscreen: A program for contaminant screening of fastq files*. [Online] Retrieved from [https://www.bioinformatics.babraham.ac.uk/projects/fastq\\_screen/](https://www.bioinformatics.babraham.ac.uk/projects/fastq_screen/)
- Wu, G. R., Ping, Y. X., & Yong, T. L. (1997). Long-term forecast on the outbreak of brown planthopper (*Nilaparvata lugens* Stål) and white-backed planthopper (*Sogatella furcifera* Horvath). *Journal Scientia Agricultura Sinica*, 30(4), 25–29.
- Xue, J., Zhou, X., Zhang, C. X., Yu, L. L., Fan, H. W., Wang, Z., ... Cheng, J. A. (2014). Genomes of the rice pest brown planthopper and its endosymbionts reveal complex complementary contributions for host adaptation. *Genome Biology*, 15(12), 521.
- Zhai, B. P., & Cheng, J. A. (2006). Summary of symposium about two migration of rice insects in 2006. *Chinese Bulletin of Entomology*, 43, 585–588. (in Chinese with English summary).
- Zhu, M., Song, Y. H., Uhm, K. B., Turner, R. W., Lee, J. H., & Roderick, G. K. (2000). Simulation of the long-range migration of brown planthopper, *Nilaparvata lugens*, (Stål), by using boundary layer atmospheric model and the geographic information system. *Journal of Asia-Pacific Entomology*, 3(1), 25–32.
- Zhu, S., Lu, Z., Hang, S., & Xu, H. (1994). Studies on regulative effects of temperature in the population of brown planthopper *Nilaparvata lugens* Stål. *Entomological Journal of East China*, 3(1), 53–59.

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Hereward JP, Cai X, Matias AMA, Walter GH, Xu C, Wang Y. Migration dynamics of an important rice pest: The brown planthopper (*Nilaparvata lugens*) across Asia—Insights from population genomics. *Evol Appl*. 2020;13:2449–2459. <https://doi.org/10.1111/eva.13047>