



# A novel negevirus isolated from *Aedes vexans* mosquitoes in Finland

Maija T. Suvanto<sup>1,2</sup> · Phuoc Truong Nguyen<sup>1</sup> · Ruut Uusitalo<sup>1,2,3</sup> · Essi M. Korhonen<sup>1,2</sup> · Giulia Faolotto<sup>4</sup> · Olli Vapalahti<sup>1,2,5</sup> · Eili Huhtamo<sup>1,2</sup> · Teemu Smura<sup>1,5</sup>

Received: 23 June 2020 / Accepted: 13 August 2020 / Published online: 20 September 2020  
© The Author(s) 2020

## Abstract

Negevirus are insect-specific enveloped RNA viruses that have been detected in mosquitoes and sandflies from various geographical locations. Here, we describe a new negevirus from Northern Europe, isolated from pool of *Aedes vexans* mosquitoes collected in Finland, designated as Mekrijärvi negevirus (MEJNV). MEJNV had a typical negevirus genome organization, is 9,740 nucleotides in length, and has a GC content of 47.53%. The MEJNV genome contains three ORFs, each containing the following identified conserved domains: ORF1 (7,068 nt) encodes a viral methyltransferase, an FtsJ-like methyltransferase, a viral RNA helicase, and an RNA-dependent RNA polymerase, ORF2 (1,242 nt) encodes a putative virion glycoprotein, and ORF3 (660 nt) encodes a putative virion membrane protein. A distinctive feature relative to other currently known negevirus is a 7-nucleotide-long overlap between ORF1 and ORF2. MEJNV shares the highest sequence identity with Ying Kou virus from China, with 67.71% nucleotide and 75.19% and 59.00% amino acid sequence identity in ORF 1 and ORF 2, respectively. ORF3 had the highest amino acid sequence similarity to Daeseongdong virus 1 and negevirus Nona 1, both with 77.61% identity, and to Ying Kou virus, with 71.22% identity. MEJNV is currently the northernmost negevirus described. Our report supports the view that negevirus are a globally distributed, diverse group of viruses that can be found from mosquitoes in a wide range of terrestrial biomes from tropical to boreal forests.

Negevirus are positive-sense single-stranded enveloped RNA viruses of insects. They have been isolated from mosquitoes of various genera (e.g., *Culex*, *Aedes*, *Anopheles*, and *Psorophora*) and from *Lutzomyia* sandflies around the world, including the Americas, Africa, Middle East, Europe, Asia, and Australia [1–6]. Negevirus have 9- to 10-kb genomes that contain three open reading frames (ORFs). ORF1 encodes a methyltransferase, an FtsJ-like helicase, a viral

RNA helicase, and an RNA-dependent RNA polymerase (RdRp), ORF2 encodes glycoproteins, and ORF3 encodes virion membrane proteins [1–3]. The recently proposed "Negevirus" taxon consists of two clades, named "*Nelorpi-virus*" and "*Sandewavirus*" [3, 5], and the taxon is related to plant viruses of the genera *Cilevirus*, *Higrevirus* and *Blunervirus* [1, 7]. In recent years, the "Negevirus" taxon has been expanding with several new negevirus being reported. The geographical origins of mosquito-associated negevirus include various terrestrial biomes, mainly temperate, subtropical, and tropical forest areas. Here, we report a novel negevirus isolated from *Aedes vexans* mosquitoes from a boreal coniferous forest area in Finland.

Mosquitoes were collected for an arbovirus study in August 2018 in Mekrijärvi (62°45'43"N, 30°57'32"E), Eastern Finland [8]. One mosquito pool consisting of 26 individual female mosquitoes caused a strong cytopathic effect (CPE) in *Aedes albopictus* C6/36 cells. The voucher specimen for this mosquito pool was identified morphologically [9] as *Aedes vexans*. RNA was extracted from virus isolation supernatant collected 10 days after infection, using TRIzol Reagent (Thermo Fisher). A sequencing library was prepared using an NEBNext Ultra II RNA Library Prep

Handling Editor: T. K. Frey.

✉ Maija T. Suvanto  
maiya.t.suvanto@helsinki.fi

<sup>1</sup> Department of Virology, Medicum, University of Helsinki, Helsinki, Finland

<sup>2</sup> Department of Veterinary Biosciences, University of Helsinki, Helsinki, Finland

<sup>3</sup> Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland


<sup>4</sup> Laboratory of Microbiology and Virology, University Hospital Maggiore Della Carità Di Novara, Piemonte, Novara, Italy

<sup>5</sup> Department of Virology and Immunology, Helsinki University Hospital, HUSLAB, Helsinki, Finland


Kit (New England Biolabs), quantified using an NEBNext Library Quant Kit for Illumina (New England Biolabs), and sequenced on a MiSeq platform using a MiSeq Reagent Kit v2 with 150-bp paired-end reads. Raw sequence reads were trimmed, and low-quality (quality score < 30) and short (< 50 nt) sequences were removed using Trimmomatic [10]. Thereafter, *de novo*-assembly was conducted using MegaHit, followed by re-assembly against the *de novo* assembled consensus sequence using the BWA-MEM algorithm [11] implemented in SAMTools version 1.8 [12]. A total of 357,745 reads mapped to the largest contig, with mean coverage of 7,624 (range, 146–17,546).

A genomic sequence of 9,740 nt was identified as a negevirus using an NCBI BLASTx search, and the isolate was designated as "Mekrijärvi negevirus" (MEJNV) (GenBank accession number MT522375). Three ORFs were identified in the MEJNV genome using NCBI ORFfinder (<https://www.ncbi.nlm.nih.gov/orffinder/>) (Fig. 1), flanked by untranslated regions (unverified) of 509 nt (5'UTR) and 162 nt (3'UTR) in length. MEJNV was most similar to Ying Kou virus strain YK1714 (NC\_040636.1) originating from *Culex pipiens pallens* mosquitoes in China, with 67.71% nucleotide sequence identity. The genome size and the detected ORFs were similar, and the GC content (GC%) of 47.53% was slightly higher when compared to previously described negeviruses (avg. GC% = 41.11%). The 3' terminus of ORF1 and the 5' terminus of ORF2 were found to overlap by seven nucleotides, and this is a distinct feature of the MEJNV genome. Conserved protein domains were identified using NCBI CD-search (<https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>) (Fig. 1). The identified domains and their positions within the ORFs align with those of previously described negeviruses [1] (Fig. 1). The evolutionary relationship of MEJNV to other related negeviruses was

assessed by constructing a phylogenetic tree based on ORF1 amino acid sequences of negevirus strains ( $n = 43$ ). The database sequences were aligned using MAFFT version 7.453 [13] and used to compute a midpoint-rooted maximum-likelihood tree with 1,000 bootstrap replicates in IQ-TREE version 1.6.12 [14], using the ModelFinder [15] and ultrafast bootstrap [16] algorithms (Fig. 2). ModelFinder suggested LG+F+R10 as the optimal substitution model. The final tree was visualized with FigTree version 1.4.4 (<https://tree.bio.ed.ac.uk/software/figtree/>). Nucleotide sequence alignments done with MAFFT showed the highest similarity between MEJNV and Ying Kou virus (NC\_040636.1) from China, with 67.71% nucleotide sequence identity. Similar analysis done with amino acid sequences showed 75.19% identity between ORF1, 59.00% between ORF2, and 71.22% between ORF3 sequences. However, the closest matches to ORF3 of MEJNV were those of Daeseongdong virus 1 (NC\_028487.1) and negevirus Nona 1 (AB972669.1), both with 77.61% identity. Notably, the GC% of MEJNV was similar to that of Ying Kou virus (GC% = 48.18%). Phylogenetic analysis suggested that MEJNV forms a monophyletic cluster (bootstrap value, 100) with Ying Kou virus. The closest relatives of MEJNV and Ying Kou viruses included viruses from Japan and South Korea (Daeseongdong virus 1 and negevirus Nona 1) and more distant relatives from South America, Australia, and the Philippines. Notably, the geographically closest negevirus sequence from Sweden in 2009 (Biggie virus, GenBank accession no. QGA70894) was not grouped with MEJNV. These results indicate that MEJNV is a distinct strain of negevirus related to negeviruses from distant geographical locations, and it is currently the northernmost negevirus strain to be isolated. Our results support the view that negeviruses are distributed globally in mosquitoes.



**Fig. 1** Genome organization of the novel Mekrijärvi negevirus. The lengths (in nucleotides) of ORFs 1–3 and the intergenic region are shown in parentheses. The nucleotide positions of ORFs and their putative conserved protein domains within the genome are shown in square brackets



**Fig. 2** Phylogenetic tree of negeviruses and related viruses (n=43) computed from protein sequences of ORF1 containing the RNA-dependent RNA polymerase (RdRp). The host, country of origin, and collection year are shown in square brackets. Unknown information

is indicated by a hyphen. Sequences were aligned using MAFFT, and a maximum-likelihood tree with 1,000 bootstrap replicates was built using IQ-TREE. The tree is rooted at the midpoint

**Acknowledgements** We would like to thank Megrin Matkailu Oy for providing facilities during fieldwork. We acknowledge CSC–IT Center for Science Ltd. (Espoo, Finland) for the allocation of computational resources.

**Funding** Open access funding provided by University of Helsinki including Helsinki University Central Hospital. This study was supported by grants from the Jane and Aatos Erkko Foundation, the Sigrid Jusélius Foundation, the Academy of Finland, the University of Helsinki Doctoral Programme in Interdisciplinary Environmental Sciences (DENVI), the Paulo Foundation and the Finnish Cultural Foundation the Kymenlaakso Regional Fund.

## Compliance with ethical standards

**Conflict of interest** The authors declare no conflict of interest.

**Availability of data and material** The Mekrijärvi negevirus genome sequence has been deposited in the NCBI GenBank database under accession number MT522375.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Vasilakis N, Forrester NL, Palacios G, Nasar F, Savji N, Rossi SL, Guzman H, Wood TG, Popov V, Gorchakov R, Gonzalez AV, Haddow AD, Watts DM, da Rosa APAT, Weaver SC, Lipkin WI, Tesh RB (2013) Negevirus: a proposed new taxon of insect-specific viruses with wide geographic distribution. *J Virol* 87(5):2475–2488. <https://doi.org/10.1128/jvi.00776-12>
- O'Brien CA, McLean BJ, Colmant AMG, Harrison JJ, Hall-Mendelin S, van den Hurk AF, Johansen CA, Watterson D, Bielefeldt-Ohmann H, Newton ND, Schulz BL, Hall RA, Hobson-Peters J (2017) Discovery and characterisation of castlereya virus, a new species of Negevirus isolated in Australia. *Evolut Bioinform* 13:10–12. <https://doi.org/10.1177/1176934317691269>
- Nunes MRT, Contreras-Gutierrez MA, Guzman H, Martins LC, Barbirato MF, Savit C, Balta V, Uribe S, Vivero R, Suaza JD, Oliveira H, Nunes Neto JP, Carvalho VL, da Silva SP, Cardoso JF, de Oliveira RS, da Silva Lemos P, Wood TG, Widen SG, Vasconcelos PFC, Fish D, Vasilakis N, Tesh RB (2017) Genetic characterization, molecular epidemiology, and phylogenetic relationships of insect-specific viruses in the taxon Negevirus. *Virology* 504:152–167. <https://doi.org/10.1016/j.virol.2017.01.022>
- Carapeta S, do Bem B, McGuinness J, Esteves A, Abecasis A, Lopes Â, de Matos AP, Piedade J, de Almeida APG, Parreira R (2015) Negevirus found in multiple species of mosquitoes from southern Portugal: isolation, genetic diversity, and replication in insect cell culture. *Virology* 483:318–328. <https://doi.org/10.1016/j.virol.2015.04.021>
- Kallies R, Kopp A, Zirkel F, Estrada A, Gillespie TR, Drosten C, Junglen S (2014) Genetic characterization of goutanap virus, a novel virus related to negevirus, cileviruses and higrviruses. *Viruses* 6(11):4346–4357. <https://doi.org/10.3390/v6114346>
- Zhao L, Mwaliko C, Atoni E, Wang Y, Zhang Y, Zhan J, Hu X, Xia H, Yuan Z (2019) Characterization of a novel tanay virus isolated from anopheles sinensis mosquitoes in Yunnan, China. *Front Microbiol* 10:1–11. <https://doi.org/10.3389/fmicb.2019.01963>
- Charles J, Tangudu CS, Hurt SL, Tumescheit C, Firth AE, Garcia-Rejon JE, Machain-Williams C, Blitvich BJ (2018) Detection of novel and recognized RNA viruses in mosquitoes from the Yucatan Peninsula of Mexico using metagenomics and characterization of their in vitro host ranges. *J Gen Virol* 99(12):1729–1738. <https://doi.org/10.1099/jgv.0.001165>
- Korhonen EM, Suvanto MT, Uusitalo R, Faolotto G, Smura T, Sane J, Vapalahti O, Huhtamo E (2020) Sindbis virus strains of divergent origin isolated from humans and mosquitoes during a recent outbreak in Finland. *Vector-Borne Zoonotic Dis.* <https://doi.org/10.1089/vbz.2019.2562>
- Becker N, Petric D, Zgomba M, Boase C et al (2010) Mosquitoes and their control. Springer, Berlin, Heidelberg
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30(15):2114–2120
- Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv. 1303.3997v1
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Subgroup GPPD (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* 25(16):2078–2079
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30(4):772–780
- Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ (2014) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32(1):268–274
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* 14(6):587–589
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2017) UFBoot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol* 35(2):518–522

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.