# LETTER TO THE EDITOR



# **Cardiac surgery and COVID-19**

Dear Editor.

I would like to comment on "Cardiac surgery in patients with confirmed COVID-19 infection: early experience." Surgery is usually considered a risky procedure for COVID-19 contact during this pandemic. In general, elective surgery is usually postponed but necessary/emergency surgery may be still indicated. Good prevention is important and the operation for COVID-19 patients does not mean there will be further disease spread. The operation in asymptomatic COVID-19 patients has been reported and there is no further transmission to anyone even though the diagnosis of COVID-19 is made after the operation.<sup>2</sup> The experience of Hussain et al<sup>1</sup> on cardiac surgery on patients with COVID-19 confirms that good planning and management are important for these operations. Furthermore, the surgeon is assured that COVID-19 patients who require cardiac surgery during this outbreak is feasible and can be done safely and effectively. Nevertheless, interesting data that Hussain et al<sup>1</sup> may have omitted is to report on the preparation of the patient before and after surgery, with isolation/surveillance procedures. For the management of a COVID-19 patient, the focus should be not only during the operation phase but also pre-and postsurgery.

### CONFLICT OF INTEREST

The author declares that there is no conflict of interest.

Viroi Wiwanitkit MD (1)

Department of Community Medicine, Dr. D. Y. Patil University, Pune, Maharashtra, India

### Correspondence

Viroj Wiwanitkit, Department of Community Medicine, Dr. D. Y. Patil University, Pune, Maharashtra, India.

Email: wviroj@yahoo.com

#### ORCID

Viroj Wiwanitkit http://orcid.org/0000-0003-1039-3728

## **REFERENCES**

- 1. Hussain A, Khan H, Lopez-Marco A, Roberts N, Oo A. Cardiac surgery in patients with confirmed COVID-19 infection: early experience. J Card Surg. 2020;35(6):1351-1353.
- 2. Joob B, Wiwanitkit V. Carpal fracture and COVID-19 infection: observation from Thailand. Indian J Orthop. 2020;54(3):393.