

## Comment on: Association of Proteinuria with Various Clinical Findings and Morphologic Variables of Oxford Classification in Immunoglobulin A Nephropathy Patients

Mohammad-Reza Ardalan

Chronic Kidney Disease Research Center, Tabriz  
University of Medical Sciences, Tabriz, Iran

**Date of Submission:** Feb 16, 2014

**Date of Acceptance:** Jun 15, 2014

### Correspondence to:

Prof. Mohammad-Reza Ardalan,  
Chronic Kidney Disease Research Center,  
Tabriz University of Medical Sciences,  
Tabriz, Iran.  
E-mail: ardalan34@yahoo.com

**How to cite this article:** Ardalan MR. Comment on: Association of Proteinuria with Various Clinical Findings and Morphologic Variables of Oxford Classification in Immunoglobulin Nephropathy Patients. *Int J Prev Med* 2014;5:1349-50.

### DEAR EDITOR,

I read with great interest, an interesting article by Nasri *et al.*, entitled “association of proteinuria with various clinical findings and morphologic variables of oxford classification in immunoglobulin A nephropathy (IgAN) patients” that was published in the esteemed international journal of preventive medicine. In an observational study on 114 patients with IgAN, they found a positive association between the proportion of crescentic fibrous formation and amount of proteinuria. They also found that Nephrotic syndrome as a definition had also a positive association with the proportion of crescent formation. They suggested the prognostic value of crescent due to its association with proteinuria, and secondly they implied the importance of treatment of proteinuria to prevent progression of IgAN.<sup>[1]</sup> On this study, I would like to point out a few notes. Newly, the Oxford classification is a novel classification for IgAN, and was developed by the renal pathology society and the working group of the international IgAN network, and its clinical validity has been proven in various studies.<sup>[2-4]</sup> This group defined four pathologic features that were more associated with the progression of kidney

disease, including: Mesangial hypercellularity, the presence of endocapillary proliferation, segmental glomerulosclerosis/adhesion, and fourthly the severity of tubular atrophy/interstitial fibrosis.<sup>[3,4]</sup> In fact, this classification offers a histopathologic grading system that is associated with renal disease consequences and by itself it is independent on clinical features. This classification also exhibits an improved capacity to predict the outcome of patients with IgAN.<sup>[2-4]</sup> All four morphologic variables of the Oxford classification have high reproducibility.<sup>[3-5]</sup> However, this classification doesn't include extracapillary proliferation, and base on the results of the Oxford classification, extracapillary proliferation is not associated with renal disease outcome,<sup>[5-9]</sup> meanwhile various investigations, have suggested that crescents have prognostic implication.<sup>[6-10]</sup> The study of Nasri *et al.* further supports the importance of extracapillary crescentic proliferation due to its association with the degree of proteinuria.<sup>[1]</sup> Therefore, we suggest further studies to investigate the prognostic implication of extracapillary proliferation in larger studies, and to examine the interesting findings by Nasri and other investigators in this field. Finally, I should note that the oxford

classification is not driven from a large number of patients ( $n = 265$  patients).<sup>[7]</sup>

## REFERENCES

1. Nasri H, Madihi Y, Merrikhi A, Gheissari A, Baradaran A, Kheiri S, *et al.* Association of proteinuria with various clinical findings and morphologic variables of Oxford classification in immunoglobulin a nephropathy patients. *Int J Prev Med* 2013;4:546-51.
2. Tanaka S, Ninomiya T, Katafuchi R, Masutani K, Tsuchimoto A, Noguchi H, *et al.* Development and validation of a prediction rule using the Oxford classification in IgA nephropathy. *Clin J Am Soc Nephrol* 2013;8:2082-90.
3. Nasri H, Ardalan MR. Significance of hyperuricemia in immunoglobulin a nephropathy. *J Renal Inj Prev* 2013;2:105-6.
4. Nasri H, Ardalan MR. Association between the proportion of globally sclerotic glomeruli and various morphologic variables and clinical data of IgA nephropathy patients. *J Renal Inj Prev* 2012;1:27-30.
5. Spasovski D. Renal markers for assessment of renal tubular and glomerular dysfunction. *J Nephropharmacol.* 2013;2:23-25.
6. Mubarak M, Nasri H. Significance of segmental glomerulosclerosis in IgA nephropathy: What is the evidence? *J Ren Inj Prev* 2013;2:113-5.
7. Working Group of the International IgA Nephropathy Network and the Renal Pathology Society, Cattran DC, Coppo R, Cook HT, Feehally J, Roberts IS, *et al.* The Oxford classification of IgA nephropathy: Rationale, clinicopathological correlations, and classification. *Kidney Int* 2009;76:534-45.
8. Ardalan MR, Sanadgol H, Nasri H, Baradaran A, Tamadon MR, Renal-Kopaei R. Vitamin D therapy in diabetic kidney disease; current knowledge on a public health problem. *J Parathy Dis* 2014;2:15-17.
9. Nasri H, Mubarak M. Significance of vasculopathy in IgA nephropathy patients with regard to Oxford classification and immunostaining findings: A single center experience. *J Ren Inj Prev* 2013;2:41-5.
10. Katafuchi R, Ninomiya T, Nagata M, Mitsuiki K, Hirakata H. Validation study of Oxford classification of IgA nephropathy: The significance of extracapillary proliferation. *Clin J Am Soc Nephrol* 2011;6:2806-13.

**Source of Support:** Nil, **Conflict of Interest:** None declared.