

Research Article

Poor Sleep Quality Worsens Static and Dynamic Balance Control in Individuals With Chronic Low Back Pain: A Cross-Sectional Study

Daniel K. Y. Zheng ^{1,2}, Zhihan Sun,^{1,3} Jeremy R. Chang ¹, Frank F. Huang ¹, Yilin Liu,¹ Siying Yu,² Jinlong Wu ⁴, Zimeng Wang,⁵ Arnold Y. L. Wong ^{1,6} and Xueqiang Wang ⁷

¹Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Hong Kong, China

²Department of Sport Rehabilitation, Shanghai University of Sport, Shanghai, China

³Department of Sport Medicine, Sichuan Province Orthopedic Hospital, Chengdu, China

⁴College of Physical Education, Southwest University, Chongqing, China

⁵Beijing Key Laboratory of Fundamental Research on Biomechanics in Clinical Application, School of Biomedical Engineering, Capital Medical University, Beijing, China

⁶Research Institute for Smart Ageing, The Hong Kong Polytechnic University, Hong Kong, China

⁷Rehabilitation Medicine Center, The Second Affiliated Hospital and Yuying Children's Hospital of Wenzhou Medical University, Wenzhou, China

Correspondence should be addressed to Arnold Y. L. Wong; arnold.wong@polyu.edu.hk and Xueqiang Wang; wangxueqiang@sus.edu.cn

Received 3 November 2024; Accepted 12 February 2025

Academic Editor: Massimiliano Valeriani

Copyright © 2025 Daniel K. Y. Zheng et al. Pain Research and Management published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Objective: To investigate the influence of sleep quality and associated factors on balance control in individuals with chronic low back pain (CLBP).

Methods: 85 participants (mean age 33.2 ± 12.5 years) with CLBP were recruited. Physical and emotional well-beings were evaluated using a battery of questionnaires. Sleep quality over the last month was assessed using the Pittsburgh Sleep Quality Index (PSQI). Participants were dichotomized into the good sleep quality (GSQ) and poor sleep quality (PSQ) groups if their PSQI scores were ≤ 5 and > 5 , respectively. Balance control was measured using the one-leg stance with eyes closed and Y-balance test.

Results: The GSQ group included 37 participants, while the PSQ group comprised 48 participants. After controlling for confounds (including gender, age, disability, anxiety, depression, and fear avoidance beliefs), participants with PSQ displayed significantly poorer performance in the one-leg stance with eyes closed and lower normalized posteromedial, posterolateral, and composite scores of the Y-balance test compared with participants with GSQ. Additionally, sleep quality accounted for 16.9%–24.9% of the variance in balance control, while age explained an additional 5.2%–13.2% of the variance. Additionally, higher levels of physical disability and anxiety were associated with poorer balance control.

Conclusions: Individuals with concurrent CLBP and PSQ exhibit significantly worse balance control than those with CLBP alone. Future studies should investigate whether improving sleep quality, physical disability, and anxiety can enhance balance in individuals with CLBP.

Keywords: anxiety; disability; low back pain; postural balance; sleep disorder

1. Introduction

Low back pain (LBP) is a significant global public health issue, affecting over 80% of individuals at some point in their lives [1, 2]. Approximately 20.1% of LBP cases progress to chronic LBP (CLBP), characterized by pain lasting longer than 12 weeks [3, 4]. Moreover, sleep disturbances have been frequently observed in individuals with CLBP. Previous studies have reported that 50%–89% of individuals with CLBP experience some forms of sleep problems (e.g., difficulties in initiating or maintaining sleep), resulting in a diminished quality of life and substantial costs to society [5–8].

Previous studies have established a prospective and bidirectional association between poorer sleep quality and more severe CLBP [9, 10]. Specifically, individuals with poor sleep quality (PSQ) are more likely to develop CLBP compared to those with good sleep quality (GSQ) [11]. Oliveira et al. found that PSQ at baseline predicted higher LBP-related disability after 6 months in older adults with LBP [12]. Furthermore, compared to asymptomatic controls, individuals with CLBP reported significantly higher Pittsburgh Sleep Quality Index (PSQI) scores, indicating poorer sleep quality [13].

Balance control relies on an intricate integration of the central nervous system with visual, proprioceptive, and vestibular inputs [14, 15]. Previous research has indicated that sleep deprivation disrupted visual, proprioceptive, and vestibular functions in healthy individuals [16–21], which can have a detrimental impact on balance control [18, 22]. PSQ is known to be associated with impaired balance control in both healthy adults and individuals with fibromyalgia [23–25]. PSQ may also lead to certain psychological conditions (such as depression and anxiety) that are commonly experienced by individuals with CLBP [11, 26, 27]. Previous research revealed that anxiety, depression, and fear avoidance beliefs share common pathways that mediate balance control [28, 29]. As such, further investigation is necessary to uncover the impact of these concomitant factors on balance control in individuals with CLBP.

Given this background, the current study aimed to examine the relative influence of sleep quality and various physical and psychological factors on balance control in adults with CLBP. We hypothesized that (1) individuals with CLBP and concomitant PSQ would display inferior balance control compared to those with CLBP and GSQ and (2) other factors (such as depression and anxiety) would also have negative effects on balance control in adults with CLBP. The findings would contribute to a better understanding of the determinants of compromised balance control in individuals with CLBP and provide insights for the comprehensive management of CLBP cases.

2. Materials and Methods

2.1. Sample Size Calculation. Previous studies revealed that the effect sizes for detecting the deficits in static and dynamic balance in healthy adults with GSQ or PSQ ranged from 0.54 to 1.15 [24, 25]. Assuming that individuals with CLBP and

GSQ or PSQ displayed a similar difference in balance control, 35 participants with GSQ or 35 participants with PSQ were needed with an effect size of 0.54, a statistical power of 0.95, and an alpha level of 0.05 as estimated by the G*Power software (Version 3.1.7, Heinrich Heine University Dusseldorf, Germany).

2.2. Participants. The study methods and results were reported following the STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) statement for cross-sectional studies [30]. This cross-sectional study was approved by the Ethics Committee of Shanghai University of Sport (approval number: 2018069). Participants were recruited from the campus, school hospital, and nearby communities through e-posters on a social mobile application, WeChat, using a convenience sampling approach. All participants provided written informed consent and underwent eligibility screening. The inclusion criteria were (1) adults aged between 18–65 years with LBP located between the 12th rib and buttocks, with or without leg pain [31]; (2) a LBP duration for more than 12 weeks [32]; and (3) an average pain intensity of at least 2 points out of 10 on an 11-point Numerical Rating Scale (NRS) in the last seven days, where 0 means no pain and 10 means the worst imaginable pain [33, 34]. Exclusion criteria included (1) the inability to complete the assessments of this study; (2) current pain intensity exceeding eight points on the NRS [35]; (3) the presence of other musculoskeletal conditions (e.g., fractures, multiple joint pain, tumors, spinal stenosis, or arthritis in the lower limbs); (4) a history of spine or lower extremity surgery; (5) pregnancy; or (6) uncorrected visual or vestibular disorders.

2.3. Procedures. Participants were instructed to complete a set of questionnaires assessing their demographics, LBP intensity, LBP-related disability, and psychological well-being. This was followed by self-reported sleep quality assessments and balance tests (Figure 1).

2.3.1. Demographic and Clinical Characteristics. Pain intensity, including current pain, average pain, and maximum pain intensity over the past week, was measured using the 11-point NRS [36]. LBP-related disability was assessed using the Chinese version of the Roland-Morris Disability Questionnaire (RMDQ), with higher scores indicating more disability [37]. Anxiety and depression levels over the last seven days were measured by the Chinese version of the 20-item Zung Self-Rating Anxiety Scale (SAS) and the 20-item Zung Self-Rating Depression Scale (SDS), with higher scores indicating more anxiety and depression, respectively [38]. Fear avoidance beliefs were assessed using Chinese version of the Fear-Avoidance Beliefs Questionnaire (FABQ), which includes subscales for physical activities (FABQ-P) and work activities (FABQ-W) [39]. Higher scores indicate more severe fear-avoidance beliefs toward physical or occupational activities. Additionally, each participant's leg length was determined by measuring the




FIGURE 1: Study flowchart.

distance between the anterior superior iliac spine and the most prominent part of the ipsilateral medial malleolus in a supine position.

2.3.2. Sleep Quality Assessment. The Chinese version of the PSQI was used to evaluate the participants' sleep quality. The PSQI is a reliable and valid measure of sleep quality in the last month [40, 41]. The total score ranges from 0 to 21, with lower scores indicating better sleep quality. A cutoff score of five was used to distinguish between individuals with GSQ and PSQ, with a sensitivity of 89.6% and a specificity of 86.5% [41]. As such, our participants were dichotomized into GSQ (PSQI scores ≤ 5) and PSQ (PSQI scores > 5) groups.

2.3.3. One-Leg Stance With Eyes Closed (OLS-C). The OLS-C was used to assess the static balance ability of the participants. Prior research has shown excellent inter-rater reliability of this test in assessing individuals with LBP (intraclass correlation coefficients (ICC) = 0.88–1.0) [35]. Furthermore, Sung and Leininger (2015) found that the OLS-C was more effective than the eyes-open condition in detecting balance deficits in individuals with LBP [42]. Participants were instructed to stand on a rigid platform with bare feet and their hands on their waists. Upon the “start” cue, the participant closed the eyes and raised one leg simultaneously. The participant was required to maintain balance for up to 120 s [43], while a timer recorded the duration. The timer stopped if the supporting foot moved, the free leg touched the ground or the supporting leg, or if either hand was taken off the waist [44]. Each leg performed three trials, with the order randomized by a coin toss. A 30 s break was given before switching to the other leg to prevent fatigue [45, 46]. Participants were allowed to practice twice on each leg before the actual data collection. The average

duration (in seconds) from both legs was used for the analysis, with a longer duration indicating better static balance performance.

2.3.4. Y-Balance Test (YBT). The lower quarter YBT (FunctionalMovement.com, Danville, VA) was employed to assess the participants' dynamic balance ability. This test has demonstrated excellent inter-rater reliability (ICC = 0.99–1.0) in individuals with CLBP [34]. During this test, participants were instructed to stand on one leg on a fixed platform, place their hands on their waist, and maintain balance, while the foot of contralateral leg pushed an indicator as far as possible in three directions: anterior, posteromedial, and posterolateral. To ensure consistency and reproducibility, the test sequence followed a specific order, starting with the anterior direction and then moving to the posteromedial and posterolateral directions. The trial was considered invalid if the participant (1) failed to maintain balance on one leg or return the reaching foot to the starting position; (2) failed to keep their hands on their waist; (3) used the ground or the indicator for foot support; or (4) kicked the indicator [25, 47]. Three trials were conducted with a 10 s break between each trial, and the average distance was calculated for each direction. The order of legs was randomized using a coin toss, with a 30 s break before switching to another leg to minimize the risk of fatigue [47, 48]. Prior to the formal trials, participants were allowed to practice twice in each direction. The reaching distance in centimeter was measured and then normalized to the length of the reaching leg using the formula: (reaching distance/reaching leg length) $\times 100\%$ [49]. A composite score was calculated by averaging the normalized scores in three directions (anterior, posteromedial, and posterolateral). The average normalized anterior, posteromedial, posterolateral, and composite scores from both legs were used for the analysis, with higher scores indicating better dynamic balance performance.

2.4. Statistical Analysis. For demographic and clinical characteristics, the chi-square tests were utilized to compare between-group differences in categorical variables. The normality of continuous variables was assessed by the Kolmogorov–Smirnov test. Independent *t*-tests were employed to compare differences between groups for parametric variables, while Mann–Whitney *U* tests were used for nonparametric variables. For balance performance, analysis of covariance (ANCOVA) was used to identify the difference between groups after controlling for various confounding demographic and clinical characteristics that showed between-group differences (with $p < 0.1$). Separate forward stepwise regression models were performed to identify factors that were independently related to OLS-C and normalized YBT scores. Variables entered were the abovementioned confounding demographic and clinical characteristics. These variables were entered into the model at a significance level of $p \leq 0.05$, and they were removed from the model if their significance level reached $p \geq 0.1$ [50]. Statistical analyses were performed using the SPSS 29.0 (IBM Corp, Armonk, NY, USA) with the significance level set at $p < 0.05$.

3. Results

3.1. Participants' Characteristics. A total of 85 participants with CLBP, including 37 males and 48 females (33.2 ± 12.5 years), were recruited (Table 1). Among them, 48 participants (56%) reported PSQ, while 37 participants (44%) reported GSQ. The mean PSQI score in the PSQ group (9.2 ± 3) was significantly higher than that in the GSQ group (3.7 ± 1.4) ($p < 0.001$). No significant differences were observed in various demographic variables between the two groups. However, compared to the GSQ group, the PSQ group exhibited higher scores on RMDQ ($p = 0.003$), SAS ($p < 0.001$), SDS ($p < 0.001$), FABQ ($p < 0.001$), FABQ-P ($p = 0.014$), and FABQ-W ($p = 0.002$).

3.2. Static and Dynamic Balance. After controlling for potential confounding demographic variables (gender and age) and clinical variables (RMDQ, SAS, SDS, and FABQ), the PSQ group showed significantly poorer performance on OLS-C than the GSQ group (24.3 ± 18.9 s vs. 50.8 ± 27.8 s, $F = 13.4$, $p < 0.001$) (Figure 2). Likewise, the PSQ group demonstrated significantly lower scores in the posteromedial ($106.8 \pm 14\%$ vs. $119.4 \pm 14.3\%$, $F = 4.9$, $p = 0.029$), posterolateral ($108.2 \pm 10.9\%$ vs. $121 \pm 12.5\%$, $F = 10.0$, $p = 0.002$), and composite measures of the YBT ($94.6 \pm 9.9\%$ vs. $104.8 \pm 10.8\%$, $F = 6.0$, $p = 0.017$) than the GSQ group after controlling for the aforementioned confounding variables (Figure 3). However, no significant between-group difference was noted in the anterior YBT score ($68.7 \pm 8\%$ vs. $74 \pm 10.4\%$, $F = 0.5$, $p = 0.466$).

3.3. Regression Analysis. Sleep quality (PSQ/GSQ) emerged as the independent predictor of OLS-C scores, normalized posteromedial, posterolateral, and composite scores on the YBT, explaining 24.9%, 16.9%, 23.3%, and 20% of the variance, respectively (Table 2). In addition to sleep quality, age accounted for an additional 13.2% of the variance in OLS-C scores, 7.4% in normalized posteromedial scores, 5.2% in normalized posterolateral scores, and 7.5% in normalized composite scores of the YBT. Regarding the normalized anterior YBT scores, RMDQ and SAS scores independently accounted for 11.3% and 5.5% of its variance, respectively.

4. Discussion

This is the first study to investigate the influence of sleep quality on balance control in individuals with CLBP. Compared to those with CLBP alone, individuals with concomitant CLBP and PSQ exhibited inferior static and dynamic balance control, as measured by the OLS-C and YBT, respectively. Additionally, sleep quality accounted for 16.9%–24.9% of the variance in the static and dynamic balance performance, while age explained an additional 5.2%–13.2% of the variances. Furthermore, physical disability and anxiety emerged as potential factors associated with suboptimal dynamic balance control.

Our study revealed that PSQ was significantly associated with elevated levels of physical disability, anxiety, depression, and fear avoidance beliefs in individuals with CLBP, which aligns with prior literature [12, 13]. A systematic review and meta-analysis have shown that insufficient sleep affects both physical and mental health [51]. Notably, individuals with PSQ are more likely to experience impaired agility, mobility, and lower limb function, which can contribute to increased physical disability [52, 53]. Mental health conditions, such as anxiety and depression, have been identified as confounders in the bidirectional association between sleep quality and CLBP [9, 11, 54]. Sleep deprivation can induce or exacerbate anxiety and depression, while these conditions, characterized by hyperarousal, excessive worrying, and negative thought patterns, can also disrupt sleep initiation and maintenance [55]. Furthermore, PSQ has been linked to higher levels of rumination and repetitive negative thinking [56, 57], which are key components of fear avoidance beliefs [58, 59]. Therefore, it is essential to note that individuals with concomitant CLBP and PSQ commonly experience heightened levels of physical disability, anxiety, depression, and fear avoidance beliefs, which can, in turn, affect balance control.

After considering the aforementioned confounding factors, our results showed that PSQ might have a detrimental effect on both OLS-C and YBT performance in individuals with CLBP. This concurs with previous studies on sleep deprivation in healthy individuals, which found that suboptimal static balance (characterized by increased center of pressure displacement during quiet bipedal standing with eyes open on a force plate) could occur after sleep deprivation over two consecutive nights [15, 17, 19]. Our findings on YBT performance were similar to those of Tanwar et al. (2021), who reported that young adults with PSQ exhibited worse normalized composite scores compared to their GSQ counterparts [25]. However, our study found no significant difference in the normalized anterior scores of the YBT between the two groups. The discrepancy may be attributed to the low sensitivity of the YBT in detecting balance deficits in the anterior direction compared to the posteromedial or posterolateral directions, possibly due to visual compensation [60, 61]. Moreover, the association between PSQ and increased fall risk has been documented in patients with fibromyalgia [62], reinforcing the broader relevance of our findings across chronic pain conditions. These results suggest that improving sleep quality could be important for managing balance and preventing falls in individuals with CLBP and other chronic pain conditions.

PSQ is associated with impaired sensorimotor coupling, fatigue, psychological distress (e.g., anxiety and depression), and cognitive deficits (e.g., executive function and attention) in individuals with CLBP, potentially compromise balance control [9, 26, 63–66]. For example, inadequate sleep has been linked to attention impairments, which can reduce available attentional resources and delay reaction times, thereby affecting balance control [64, 67]. Aguiar and Barela (2015) reported that individuals with PSQ faced challenges in uncoupling irrelevant sensory information and were inefficient at selecting the most relevant sensory cues for

TABLE 1: Demographic and clinical characteristics.

	Poor sleep quality (<i>n</i> = 48)	Good sleep quality (<i>n</i> = 37)	$\chi^2/t/z$	<i>p</i>
Gender				
Male	17	20	2.952	0.086
Female	31	17		
Age (years)	35.6 ± 13.5	29.9 ± 10.2	-2.213	0.03
Height (cm)	167.8 ± 9.2	167.6 ± 8.2	-0.092	0.927
Weight (kg)	64.6 ± 12.9	65.6 ± 13.6	0.348	0.728
BMI (kg/m ²)	22.8 ± 2.9	23.2 ± 3.7	-0.31 [†]	0.756
Left leg length (cm)	85.9 ± 5.7	84.5 ± 4.9	-1.204	0.232
Right leg length (cm)	85.7 ± 5.8	84.4 ± 5.1	-1.069	0.288
Pain duration (years)	7 ± 6.6	6.8 ± 6.3	-0.129 [†]	0.898
Average pain	3.8 ± 1.3	3.5 ± 1.2	-1.079 [†]	0.280
Current pain	3.2 ± 1.6	2.6 ± 1.4	-1.596 [†]	0.111
Maximum pain	5.3 ± 1.6	4.9 ± 1.5	-1.201 [†]	0.230
PSQI	9.2 ± 3	3.7 ± 1.4	-7.915 [†]	< 0.001*
RMDQ	7.8 ± 5.7	4.4 ± 3.5	-2.972 [†]	0.003*
SAS	46.9 ± 8.9	36 ± 5.2	-5.683 [†]	< 0.001*
SDS	46.8 ± 10.7	34.5 ± 5.8	-5.384 [†]	< 0.001*
FABQ (total score)	41.9 ± 10.8	32.5 ± 12.4	-3.385 [†]	< 0.001*
FABQ-P	13.5 ± 4.3	10.6 ± 5.5	-2.445 [†]	0.014*
FABQ-W	28.4 ± 8.6	21.9 ± 8.4	-3.087 [†]	0.002*

Note: The values are presented as number or mean ± standard deviation.

Abbreviations: BMI, body mass index; FABQ, Fear Avoidance Beliefs Questionnaire; FABQ-P, Fear Avoidance Beliefs Questionnaire for physical activities; FABQ-W, Fear Avoidance Beliefs Questionnaire for work activities; PSQI, Pittsburgh Sleep Quality Index; RMDQ, Roland Morris disability questionnaire; SAS, Zung Self-Rating Anxiety Scale; SDS, Zung Self-Rating Depression Scale.

[†]Mann-Whitney *U*-tests.

**p* < 0.05.




FIGURE 2: The results of one-leg stance test. Note: Data are expressed as the mean ± standard deviation; GSQ, good sleep quality; PSQ, poor sleep quality. **p* < 0.05.

appropriate motor responses in balance control, suggesting a noisy balance control system [66]. Furthermore, sleep disturbance may induce neurophysiological alterations, such as reduced muscle sympathetic nerve activity and delayed efferent signals from the central nervous system to balance control effectors in both healthy individuals and those with fibrositis syndrome [67–69]. Studies have also demonstrated that sleep deprivation decreases activity in brain regions related to the vestibular system (temporal lobes), proprioceptive system (basal ganglia and cerebellum), and attention and executive function (prefrontal cortex) in healthy

individuals. These brain changes may increase the risk of falls and balance impairments [16–21]. Additionally, Thomas et al. revealed that one night of sleep deprivation resulted in decreased neuronal activity in the cortico-thalamic network, which is involved in attention and higher-order cognitive performance [70]. Given our results, future studies should investigate the central mechanisms underlying balance control impairments in individuals with concurrent CLBP and PSQ using neuroimaging techniques, such as electroencephalography or functional near-infrared spectroscopy [18].




FIGURE 3: The results of Y-balance test scores. *Note:* Data are expressed as the mean \pm standard deviation; “anterior,” “posteromedial,” “posterolateral,” and “composite” are normalized scores of the Y-balance test here. GSQ, good sleep quality; PSQ, poor sleep quality; ns, nonsignificant ($p > 0.05$); * $p < 0.05$.

TABLE 2: Regression models^a summaries for OLS-C and normalized YBT scores including standardized beta coefficients and R^2 .

Variable	Model	Predictor	Standardized beta	R^2	R^2 change
OLS	1	Group	−0.499	0.249	
	2	Group	−0.413		
		Age	−0.373	0.381	0.132
Anterior	3	RMDQ	−0.337	0.113	
	4	RMDQ	−0.263		
		SAS	−0.244	0.168	0.055
Posteromedial	1	Group	−0.411	0.169	
	2	Group	−0.348		
		Age	−0.278	0.243	0.074
Posterolateral	1	Group	−0.482	0.233	
	2	Group	−0.428		
		Age	−0.236	0.285	0.052
Composite	1	Group	−0.448	0.2	
	2	Group	−0.384		
		Age	−0.281	0.275	0.075

Abbreviations: FABQ, Fear Avoidance Beliefs Questionnaire; OLS-C, one-leg stance with eyes closed; RMDQ, Roland Morris Disability Questionnaire; SAS, Self-Rating Anxiety Scale; SDS, Self-Rating Depression Scale; YBT, Y-balance test.

^aVariables entered into the forward stepwise regression models: group (poor/good sleep quality), gender (female/male), age in years, RMDQ, SAS, SDS, and FABQ scores.

Our findings indicate that sleep quality (16.9%–24.9%) is a significant determinant of impaired balance control in individuals with CLBP, followed by age (5.2%–13.2%). These results align with Liu-Ambrose et al., who found that age was the primary determinant of balance control, accounting for 9%–14% of the variance in older women with osteoporosis [50]. Emerging evidence supports the notion that improving sleep quality can reduce pain in various chronic pain conditions, including CLBP and chronic widespread pain [8, 71–74]. Afolalu et al. found that enhanced sleep quality was associated with improved physical function in the general population, although the effect size was small [51]. Therefore, to enhance balance control in adults with CLBP, it is crucial to identify and address those modifiable factors (sleep quality, physical disability, and anxiety) [75, 76]. However, further research, including controlled clinical trials, is necessary to validate these findings.

Several limitations should be considered when interpreting our results. First, our cross-sectional design precludes the establishment of causal relationships between sleep quality and balance control [77]. Second, our assessment of sleep quality relied on a self-reported questionnaire, although the psychometric properties of the PSQI have been established [9, 77, 78]. Future studies should incorporate objective measures such as polysomnography or actigraphy to collect objective data on sleep quality and quantity [77–79]. Third, this study did not include asymptomatic controls for comparison. Nevertheless, prior studies have established that individuals with CLBP generally exhibit poorer balance control than healthy control across adult populations [80–82]. Fourth, most of our participants were young, with an average age of 33.2 ± 12.5 years, and these findings may not be generalized to older adults with CLBP,

who may be more susceptible to falls. Future studies should investigate balance control in older adults with concomitant CLBP and PSQ.

5. Conclusion

This study makes a pioneering contribution, by revealing the association between poorer sleep quality and impaired balance control in individuals with CLBP. PSQ, along with physical disability and anxiety, is a potential factor that contribute to impaired balance control in individuals with CLBP. Addressing sleep quality and managing these factors may be crucial for improving balance control and overall well-being in this population. Further research is warranted to investigate the effects of interventions targeting these modifiable factors on balance control outcomes.

Data Availability Statement

Deidentified individual participant data can be made available upon request to wangxueqiang@sus.edu.cn.

Conflicts of Interest

The authors declare no conflicts of interest.

Funding

This work was supported by the National Natural Science Foundation of China (82372578) and GP Batteries Industrial Safety Trust Fund (R-ZDDR).

Acknowledgments

The authors wish to thank all of the participants supporting this research.

References

- [1] C. Maher, M. Underwood, and R. Buchbinder, "Non-Specific Low Back Pain," *The Lancet* 389, no. 10070 (2017): 736–747, [https://doi.org/10.1016/s0140-6736\(16\)30970-9](https://doi.org/10.1016/s0140-6736(16)30970-9).
- [2] J. Hartvigsen, M. J. Hancock, A. Kongsted, et al., "What Low Back Pain Is and Why We Need to Pay Attention," *The Lancet* 391, no. 10137 (2018): 2356–2367, [https://doi.org/10.1016/s0140-6736\(18\)30480-x](https://doi.org/10.1016/s0140-6736(18)30480-x).
- [3] G. B. Andersson, "Epidemiological Features of Chronic Low-Back Pain," *The Lancet* 354, no. 9178 (1999): 581–585, [https://doi.org/10.1016/s0140-6736\(99\)01312-4](https://doi.org/10.1016/s0140-6736(99)01312-4).
- [4] D. Hoy, C. Bain, G. Williams, et al., "A Systematic Review of the Global Prevalence of Low Back Pain," *Arthritis & Rheumatism* 64, no. 6 (2012): 2028–2037, <https://doi.org/10.1002/art.34347>.
- [5] T. H. T. Wong, K. S. K. Lee, S. M. C. Lo, et al., "Challenges, Concerns, and Experiences of Community-Dwelling Older Women with Chronic Low Back Pain-A Qualitative Study in Hong Kong, China," *Healthcare (Basel)* 11, no. 7 (2023): 945, <https://doi.org/10.3390/healthcare11070945>.
- [6] L. M. McCracken and G. L. Iverson, "Disrupted Sleep Patterns and Daily Functioning in Patients with Chronic Pain," *Pain Research and Management* 7, no. 2 (2002): 75–79, <https://doi.org/10.1155/2002/579425>.
- [7] S. M. Alsaadi, J. H. McAuley, J. M. Hush, and C. G. Maher, "Prevalence of Sleep Disturbance in Patients with Low Back Pain," *European Spine Journal* 20, no. 5 (2011): 737–743, <https://doi.org/10.1007/s00586-010-1661-x>.
- [8] P. K. Morelhão, L. J. Kim, R. Z. Pinto, S. Tufik, and M. L. Andersen, "Should Physical Therapists Assess Sleep Quality in Patients Seeking Care for Low Back Pain?" *Physical Therapy* 99, no. 8 (2019): 961–963, <https://doi.org/10.1093/ptj/pzz058>.
- [9] M. Z. Campanini, A. D. González, S. M. Andrade, et al., "Bidirectional Associations between Chronic Low Back Pain and Sleep Quality: A Cohort Study with Schoolteachers," *Physiology & Behavior* 254 (2022): <https://doi.org/10.1016/j.physbeh.2022.113880>.
- [10] E. M. O'Brien, L. B. Waxenberg, J. W. Atchison, et al., "Intra-individual Variability in Daily Sleep and Pain Ratings Among Chronic Pain Patients: Bidirectional Association and the Role of Negative Mood," *The Clinical Journal of Pain* 27, no. 5 (2011): 425–433, <https://doi.org/10.1097/AJP.0b013e318208c8e4>.
- [11] L. Barazzetti, A. Garcez, P. C. Freitas Sant'Anna, F. Souza de Bairos, J. S. Dias-da-Costa, and M. T. Anselmo Olinto, "Does Sleep Quality Modify the Relationship between Common Mental Disorders and Chronic Low Back Pain in Adult Women?" *Sleep Medicine* 96 (2022): 132–139, <https://doi.org/10.1016/j.sleep.2022.05.006>.
- [12] S. D. Oliveira, R. Z. Pinto, C. Gobbi, et al., "Sleep Quality Predicts Functional Disability in Older Adults with Low Back Pain: A Longitudinal Study," *Journal of Applied Gerontology* 41, no. 11 (2022): 2374–2381, <https://doi.org/10.1177/07334648221113500>.
- [13] M. Marty, S. Rozenberg, B. Duplan, P. Thomas, B. Duquesnoy, and F. Allaert, "Quality of Sleep in Patients with Chronic Low Back Pain: a Case-Control Study," *European Spine Journal* 17, no. 6 (2008): 839–844, <https://doi.org/10.1007/s00586-008-0660-7>.
- [14] D. K. Y. Zheng, G. N. Kawchuk, A. E. Bussi eres, et al., "Trends of Low Back Pain Research in Older and Working-Age Adults from 1993 to 2023: A Bibliometric Analysis," *Journal of Pain Research* 16 (2023): 3325–3341, <https://doi.org/10.2147/jpr.S425672>.
- [15] L. Montesinos, R. Castaldo, F. P. Cappuccio, and L. Pecchia, "Day-to-day Variations in Sleep Quality Affect Standing Balance in Healthy Adults," *Scientific Reports* 8, no. 1 (2018): <https://doi.org/10.1038/s41598-018-36053-4>.
- [16] S. Besnard, B. Tighilet, C. Chabbert, et al., "The Balance of Sleep: Role of the Vestibular Sensory System," *Sleep Medicine Reviews* 42 (2018): 220–228, <https://doi.org/10.1016/j.smrv.2018.09.001>.
- [17] J. C. Wu, J. C. Gillin, M. S. Buchsbaum, et al., "The Effect of Sleep Deprivation on Cerebral Glucose Metabolic Rate in Normal Humans Assessed with Positron Emission Tomography," *Sleep* 14, no. 2 (1991): 155–162.
- [18] J. Ma, Y. J. Yao, R. M. Ma, et al., "Effects of Sleep Deprivation on Human Postural Control, Subjective Fatigue Assessment and Psychomotor Performance," *Journal of International Medical Research* 37, no. 5 (2009): 1311–1320, <https://doi.org/10.1177/147323000903700506>.
- [19] M. Fabbri, M. Martoni, M. J. Esposito, G. Brighetti, and V. Natale, "Postural Control after a Night without Sleep," *Neuropsychologia* 44, no. 12 (2006): 2520–2525, <https://doi.org/10.1016/j.neuropsychologia.2006.03.033>.
- [20] A. Muzur, E. F. Pace-Schott, and J. A. Hobson, "The Prefrontal Cortex in Sleep," *Trends in Cognitive Sciences* 6, no. 11 (2002): 475–481, [https://doi.org/10.1016/s1364-6613\(02\)01992-7](https://doi.org/10.1016/s1364-6613(02)01992-7).

- [21] A. N. Hudson, H. P. A. Van Dongen, and K. A. Honn, "Sleep Deprivation, Vigilant Attention, and Brain Function: a Review," *Neuropsychopharmacology* 45, no. 1 (2020): 21–30, <https://doi.org/10.1038/s41386-019-0432-6>.
- [22] J. B. Caccese, F. V. Santos, F. K. Yamaguchi, T. A. Buckley, and J. J. Jeka, "Persistent Visual and Vestibular Impairments for Postural Control Following Concussion: A Cross-Sectional Study in University Students," *Sports Medicine* 51, no. 10 (2021): 2209–2220, <https://doi.org/10.1007/s40279-021-01472-3>.
- [23] Y. Bolukbas and B. Celik, "Assessment of Sleep Quality in Fibromyalgia Syndrome and its Effect on Postural Balance and Functional Status," *Journal of Back and Musculoskeletal Rehabilitation* 34, no. 2 (2021): 235–242, <https://doi.org/10.3233/bmr-181320>.
- [24] F. Furtado, B. D. Gonçalves, I. L. Abranches, A. F. Abrantes, and A. Forner-Cordero, "Chronic Low Quality Sleep Impairs Postural Control in Healthy Adults," *PLoS One* 11, no. 10 (2016): e0163310, <https://doi.org/10.1371/journal.pone.0163310>.
- [25] T. Tanwar, Z. Veqar, A. K. Ghrouz, D. W. Spence, and S. R. Pandi-Perumal, "Is Poor Sleep Quality Associated with a Deterioration in Postural Control?" *Sleep Sci* 14, no. 3 (2021): 207–213, <https://doi.org/10.5935/1984-0063.20200061>.
- [26] J. A. Robertson, R. J. Purple, P. Cole, Z. Zaiwalla, K. Wulff, and K. T. Pattinson, "Sleep Disturbance in Patients Taking Opioid Medication for Chronic Back Pain," *Anaesthesia* 71, no. 11 (2016): 1296–1307, <https://doi.org/10.1111/anae.13601>.
- [27] R. Marin, T. Cyhan, and W. Miklos, "Sleep Disturbance in Patients with Chronic Low Back Pain," *American Journal of Physical Medicine & Rehabilitation* 85, no. 5 (2006): 430–435, <https://doi.org/10.1097/01.phm.0000214259.06380.79>.
- [28] C. D. Balaban and J. F. Thayer, "Neurological Bases for Balance-Anxiety Links," *Journal of Anxiety Disorders* 15, no. 1-2 (2001): 53–79, [https://doi.org/10.1016/s0887-6185\(00\)00042-6](https://doi.org/10.1016/s0887-6185(00)00042-6).
- [29] M. Belvederi Murri, F. Triolo, A. Coni, et al., "Instrumental Assessment of Balance and Gait in Depression: A Systematic Review," *Psychiatry Research* 284 (2020): <https://doi.org/10.1016/j.psychres.2019.112687>.
- [30] E. Von Elm, D. G. Altman, M. Egger, S. J. Pocock, P. C. Gøtzsche, and J. P. Vandenbroucke, "The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies," *The Lancet* 370, no. 9596 (2007): 1453–1457, [https://doi.org/10.1016/s0140-6736\(07\)61602-x](https://doi.org/10.1016/s0140-6736(07)61602-x).
- [31] N. N. Knezevic, K. D. Candido, J. W. S. Vlaeyen, J. Van Zundert, and S. P. Cohen, "Low Back Pain," *The Lancet* 398, no. 10294 (2021): 78–92, [https://doi.org/10.1016/s0140-6736\(21\)00733-9](https://doi.org/10.1016/s0140-6736(21)00733-9).
- [32] K. R. Cohen, "Management of Chronic Low Back Pain," *JAMA Internal Medicine* 182, no. 2 (2022): 222–223, <https://doi.org/10.1001/jamainternmed.2021.7359>.
- [33] Y. M. Alshehre, K. Alkhathami, K. Brizzolara, M. Weber, and S. Wang-Price, "Effectiveness of Spinal Stabilization Exercises on Dynamic Balance in Adults with Chronic Low Back Pain," *International Journal of Sports Physical Therapy* 18, no. 1 (2023): 173–187, <https://doi.org/10.26603/001c.68075>.
- [34] Y. Alshehre, K. Alkhathami, K. Brizzolara, M. Weber, and S. Wang-Price, "Reliability and Validity of the Y-Balance Test in Young Adults with Chronic Low Back Pain," *International Journal of Sports Physical Therapy* 16, no. 3 (2021): 628–635, <https://doi.org/10.26603/001c.23430>.
- [35] J. Tidstrand and E. Horneij, "Inter-rater Reliability of Three Standardized Functional Tests in Patients with Low Back Pain," *BMC Musculoskeletal Disorders* 10, no. 1 (2009): 58, <https://doi.org/10.1186/1471-2474-10-58>.
- [36] A. Williamson and B. Hoggart, "Pain: a Review of Three Commonly Used Pain Rating Scales," *Journal of Clinical Nursing* 14, no. 7 (2005): 798–804, <https://doi.org/10.1111/j.1365-2702.2005.01121.x>.
- [37] M. Roland and J. Fairbank, "The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire," *Spine* 25, no. 24 (2000): 3115–3124, <https://doi.org/10.1097/00007632-200012150-00006>.
- [38] B. O. Olatunji, B. J. Deacon, J. S. Abramowitz, and D. F. Tolin, "Dimensionality of Somatic Complaints: Factor Structure and Psychometric Properties of the Self-Rating Anxiety Scale," *Journal of Anxiety Disorders* 20, no. 5 (2006): 543–561, <https://doi.org/10.1016/j.janxdis.2005.08.002>.
- [39] G. Waddell, M. Newton, I. Henderson, D. Somerville, and C. J. Main, "A Fear-Avoidance Beliefs Questionnaire (FABQ) and the Role of Fear-Avoidance Beliefs in Chronic Low Back Pain and Disability," *Pain* 52, no. 2 (1993): 157–168, [https://doi.org/10.1016/0304-3959\(93\)90127-b](https://doi.org/10.1016/0304-3959(93)90127-b).
- [40] T. Mollayeva, P. Thurairajah, K. Burton, S. Mollayeva, C. M. Shapiro, and A. Colantonio, "The Pittsburgh sleep Quality Index as a Screening Tool for Sleep Dysfunction in Clinical and Non-clinical Samples: A Systematic Review and Meta-Analysis," *Sleep Medicine Reviews* 25 (2016): 52–73, <https://doi.org/10.1016/j.smrv.2015.01.009>.
- [41] D. J. Buysse, C. F. Reynolds, T. H. Monk, S. R. Berman, and D. J. Kupfer, "The Pittsburgh Sleep Quality Index: a New Instrument for Psychiatric Practice and Research," *Psychiatry Research* 28, no. 2 (1989): 193–213, [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4).
- [42] P. S. Sung and P. M. Leininger, "A Kinematic and Kinetic Analysis of Spinal Region in Subjects with and without Recurrent Low Back Pain during One Leg Standing," *Clinical Biomechanics* 30, no. 7 (2015): 696–702, <https://doi.org/10.1016/j.clinbiomech.2015.05.003>.
- [43] T. Arai, H. Fujita, K. Maruya, Y. Morita, R. Asahi, and H. Ishibashi, "The One-Leg Portion of the Stand-Up Test Predicts Fall Risk in Aged Individuals: A Prospective Cohort Study," *Journal of Orthopaedic Science* 25, no. 4 (2020): 688–692, <https://doi.org/10.1016/j.jos.2019.06.014>.
- [44] S. M. Gorji, H. Mohammadi Nia Samakosh, P. Watt, P. Henrique Marchetti, and R. Oliveira, "Pain Neuroscience Education and Motor Control Exercises versus Core Stability Exercises on Pain, Disability, and Balance in Women with Chronic Low Back Pain," *International Journal of Environmental Research and Public Health* 19, no. 5 (2022): 2694, <https://doi.org/10.3390/ijerph19052694>.
- [45] M. D. Jakobsen, E. Sundstrup, P. Krustrup, and P. Aagaard, "The Effect of Recreational Soccer Training and Running on Postural Balance in Untrained Men," *European Journal of Applied Physiology* 111, no. 3 (2011): 521–530, <https://doi.org/10.1007/s00421-010-1669-2>.
- [46] A. G. Schneiders, S. J. Sullivan, P. Handcock, A. Gray, and P. R. McCrory, "Sports Concussion Assessment: the Effect of Exercise on Dynamic and Static Balance," *Scandinavian Journal of Medicine & Science in Sports* 22, no. 1 (2012): 85–90, <https://doi.org/10.1111/j.1600-0838.2010.01141.x>.
- [47] T. A. Boff, F. Pasinato, Á. J. Ben, J. E. Bosmans, M. van Tulder, and R. L. Carregaro, "Effectiveness of Spinal Manipulation and Myofascial Release Compared with Spinal Manipulation Alone on Health-Related Outcomes in Individuals with Non-specific Low Back Pain: Randomized Controlled Trial,"

- Physiotherapy* 107 (2020): 71–80, <https://doi.org/10.1016/j.physio.2019.11.002>.
- [48] Z. W. Yan, Z. Yang, J. Yang, Y. F. Chen, X. B. Zhang, and C. L. Song, “Tai Chi for Spatiotemporal Gait Features and Dynamic Balancing Capacity in Elderly Female Patients with Non-specific Low Back Pain: A Six-Week Randomized Controlled Trial,” *Journal of Back and Musculoskeletal Rehabilitation* 35, no. 6 (2022): 1311–1319, <https://doi.org/10.3233/bmr-210247>.
- [49] H. Bennett, S. Chalmers, S. Milanese, and J. Fuller, “The Association between Y-Balance Test Scores, Injury, and Physical Performance in Elite Adolescent Australian Footballers,” *Journal of Science and Medicine in Sport* 25, no. 4 (2022): 306–311, <https://doi.org/10.1016/j.jsams.2021.10.014>.
- [50] T. Liu-Ambrose, J. J. Eng, K. M. Khan, A. Mallinson, N. D. Carter, and H. A. McKay, “The Influence of Back Pain on Balance and Functional Mobility in 65- to 75-Year-Old Women with Osteoporosis,” *Osteoporosis International* 13, no. 11 (2002): 868–873, <https://doi.org/10.1007/s001980200119>.
- [51] E. F. Afolalu, F. Ramlee, and N. K. Y. Tang, “Effects of Sleep Changes on Pain-Related Health Outcomes in the General Population: A Systematic Review of Longitudinal Studies with Exploratory Meta-Analysis,” *Sleep Medicine Reviews* 39 (2018): 82–97, <https://doi.org/10.1016/j.smrv.2017.08.001>.
- [52] M. Z. Campanini, A. E. Mesas, J. A. Carnicero-Carreño, F. Rodríguez-Artalejo, and E. Lopez-Garcia, “Duration and Quality of Sleep and Risk of Physical Function Impairment and Disability in Older Adults: Results from the ENRICA and ELSA Cohorts,” *Aging and disease* 10, no. 3 (2019): 557–569, <https://doi.org/10.14336/ad.2018.0611>.
- [53] J. Wang, K. Zheng, J. Wu, R. Wang, X. Zhuang, and X. Wang, “Correlations between Age, Pain Intensity, Disability, and Tactile Acuity in Patients with Chronic Low Back Pain,” *Pain Research and Management* 2022 (2022): 1–9, <https://doi.org/10.1155/2022/2907009>.
- [54] P. K. Morelhão, C. Gobbi, D. G. D. Christofaro, et al., “Bidirectional Association between Sleep Quality and Low Back Pain in Older Adults: A Longitudinal Observational Study,” *Archives of Physical Medicine and Rehabilitation* 103, no. 8 (2022): 1558–1564, <https://doi.org/10.1016/j.apmr.2021.11.009>.
- [55] S. Hyun, H. C. Hahm, G. T. F. Wong, E. Zhang, and C. H. Liu, “Psychological Correlates of Poor Sleep Quality Among U.S. Young Adults during the COVID-19 Pandemic,” *Sleep Medicine* 78 (2021): 51–56, <https://doi.org/10.1016/j.sleep.2020.12.009>.
- [56] F. Clancy, A. Prestwich, L. Caperon, A. Tsipa, and D. B. O’Connor, “The Association between Worry and Rumination with Sleep in Non-clinical Populations: a Systematic Review and Meta-Analysis,” *Health Psychology Review* 14, no. 4 (2020): 427–448, <https://doi.org/10.1080/17437199.2019.1700819>.
- [57] P. Leung, S. H. Li, and B. M. Graham, “The Relationship between Repetitive Negative Thinking, Sleep Disturbance, and Subjective Fatigue in Women with Generalized Anxiety Disorder,” *British Journal of Clinical Psychology* 61, no. 3 (2022): 666–679, <https://doi.org/10.1111/bjc.12356>.
- [58] M. L. Moulds, E. Kandris, S. Starr, and A. C. Wong, “The Relationship between Rumination, Avoidance and Depression in a Non-clinical Sample,” *Behaviour Research and Therapy* 45, no. 2 (2007): 251–261, <https://doi.org/10.1016/j.brat.2006.03.003>.
- [59] H. Kim and M. G. Newman, “Avoidance of Negative Emotional Contrast from Worry and Rumination: An Application of the Contrast Avoidance Model,” *Journal of Behavioral and Cognitive Therapy* 32, no. 1 (2022): 33–43, <https://doi.org/10.1016/j.jbct.2021.12.007>.
- [60] T. L. Hooper, C. R. James, J. M. Brismée, et al., “Dynamic Balance as Measured by the Y-Balance Test Is Reduced in Individuals with Low Back Pain: A Cross-Sectional Comparative Study,” *Physical Therapy in Sport* 22 (2016): 29–34, <https://doi.org/10.1016/j.ptsp.2016.04.006>.
- [61] P. Sun, K. Li, X. Yao, Z. Wu, and Y. Yang, “Association between Functional Disability with Postural Balance Among Patients with Chronic Low Back Pain,” *Frontiers in Neurology* 14 (2023): <https://doi.org/10.3389/fneur.2023.1136137>.
- [62] N. Akkaya, S. Akkaya, N. S. Atalay, M. Acar, N. Catalbas, and F. Sahin, “Assessment of the Relationship between Postural Stability and Sleep Quality in Patients with Fibromyalgia,” *Clinical Rheumatology* 32, no. 3 (2013): 325–331, <https://doi.org/10.1007/s10067-012-2117-y>.
- [63] Z. Zhou, E. S. Hui, G. S. Kranz, et al., “Potential Mechanisms Underlying the Accelerated Cognitive Decline in People with Chronic Low Back Pain: A Scoping Review,” *Ageing Research Reviews* 82 (2022): <https://doi.org/10.1016/j.arr.2022.101767>.
- [64] J. Lim and D. F. Dinges, “Sleep Deprivation and Vigilant Attention,” *Annals of the New York Academy of Sciences* 1129, no. 1 (2008): 305–322, <https://doi.org/10.1196/annals.1417.002>.
- [65] S. P. Becker, M. A. Jarrett, A. M. Luebke, A. A. Garner, G. L. Burns, and M. J. Kofler, “Sleep in a Large, Multi-University Sample of College Students: Sleep Problem Prevalence, Sex Differences, and Mental Health Correlates,” *Sleep Health* 4, no. 2 (2018): 174–181, <https://doi.org/10.1016/j.sleh.2018.01.001>.
- [66] S. A. Aguiar and J. A. Barela, “Adaptation of Sensorimotor Coupling in Postural Control Is Impaired by Sleep Deprivation,” *PLoS One* 10, no. 3 (2015): e0122340, <https://doi.org/10.1371/journal.pone.0122340>.
- [67] R. Robillard, F. Prince, M. Boissonneault, D. Filipini, and J. Carrier, “Effects of Increased Homeostatic Sleep Pressure on Postural Control and Their Modulation by Attentional Resources,” *Clinical Neurophysiology* 122, no. 9 (2011): 1771–1778, <https://doi.org/10.1016/j.clinph.2011.02.010>.
- [68] M. Kato, B. G. Phillips, G. Sigurdsson, K. Narkiewicz, C. A. Pesek, and V. K. Somers, “Effects of Sleep Deprivation on Neural Circulatory Control,” *Hypertension* 35, no. 5 (2000): 1173–1175, <https://doi.org/10.1161/01.hyp.35.5.1173>.
- [69] H. Moldofsky, P. Scarisbrick, R. England, and H. Smythe, “Musculoskeletal Symptoms and Non-REM Sleep Disturbance in Patients with “fibrositis Syndrome” and Healthy Subjects,” *Psychosomatic Medicine* 37, no. 4 (1975): 341–351, <https://doi.org/10.1097/00006842-197507000-00008>.
- [70] M. Thomas, H. Sing, G. Belenky, et al., “Neural Basis of Alertness and Cognitive Performance Impairments during Sleepiness. I. Effects of 24 H of Sleep Deprivation on Waking Human Regional Brain Activity,” *Journal of Sleep Research* 9, no. 4 (2000): 335–352, <https://doi.org/10.1046/j.1365-2869.2000.00225.x>.
- [71] J. R. Chang, Y. K. Cheung, S. Sharma, et al., “Comparative Effectiveness of Non-pharmacological Interventions on Sleep in Individuals with Chronic Musculoskeletal Pain: A Systematic Review with Network Meta-Analysis,” *Sleep Medicine Reviews* 73 (2024): <https://doi.org/10.1016/j.smrv.2023.101867>.
- [72] T. A. Roehrs, “Does Effective Management of Sleep Disorders Improve Pain Symptoms?” *Drugs* 69, no. Supplement 2 (2009): 5–11, <https://doi.org/10.2165/11531260-000000000-00000>.
- [73] K. A. Davies, G. J. Macfarlane, B. I. Nicholl, et al., “Restorative Sleep Predicts the Resolution of Chronic Widespread Pain: Results from the EPIFUND Study,” *Rheumatology* 47, no. 12

- (2008): 1809–1813, <https://doi.org/10.1093/rheumatology/ken389>.
- [74] J. R. Chang, X. Wang, G. Lin, D. Samartzis, S. M. Pinto, and A. Y. L. Wong, “Are Changes in Sleep Quality/Quantity or Baseline Sleep Parameters Related to Changes in Clinical Outcomes in Patients with Nonspecific Chronic Low Back Pain?: A Systematic Review,” *The Clinical Journal of Pain* 38, no. 4 (2022): 292–307, <https://doi.org/10.1097/ajp.0000000000001008>.
 - [75] D. K. Zheng, J. Q. Liu, J. R. Chang, et al., “Are Changes in Pain Intensity Related to Changes in Balance Control in Individuals with Chronic Non-specific Low Back Pain? A Systematic Review and Meta-Analysis,” *Journal of Sport and Health Science* 14 (2025): <https://doi.org/10.1016/j.jshs.2024.100989>.
 - [76] K. Zheng and X. Wang, “Publications on the Association between Cognitive Function and Pain from 2000 to 2018: A Bibliometric Analysis Using CiteSpace,” *Medical Science Monitor* 25 (2019): 8940–8951, <https://doi.org/10.12659/msm.917742>.
 - [77] F. Hita-Contreras, N. Zagalaz-Anula, A. Martínez-Amat, et al., “Sleep Quality and its Association with Postural Stability and Fear of Falling Among Spanish Postmenopausal Women,” *Menopause* 25, no. 1 (2018): 62–69, <https://doi.org/10.1097/gme.0000000000000941>.
 - [78] K. Aoyagi, J. He, D. J. Clauw, and N. K. Sharma, “Sleep Quality in Individuals with Chronic Low Back Pain and Central Sensitization,” *Physiotherapy Research International: The Journal for Researchers and Clinicians in Physical Therapy* 27, no. 4 (2022): e1968, <https://doi.org/10.1002/pri.1968>.
 - [79] W. Yan, K. Zheng, L. Weng, et al., “Bibliometric Evaluation of 2000-2019 Publications on Functional Near-Infrared Spectroscopy,” *NeuroImage* 220 (2020): <https://doi.org/10.1016/j.neuroimage.2020.117121>.
 - [80] J. A. Mingorance, P. Montoya, J. G. Vivas Miranda, and I. Riquelme, “Differences in Postural Balance, Pain Sensitivity and Depression between Individuals with Acute and Chronic Back Pain,” *Journal of Clinical Medicine* 11, no. 10 (2022): 2700, <https://doi.org/10.3390/jcm11102700>.
 - [81] L. Ge, C. Wang, H. Zhou, Q. Yu, and X. Li, “Effects of Low Back Pain on Balance Performance in Elderly People: a Systematic Review and Meta-Analysis,” *Eur Rev Aging Phys Act* 18, no. 1 (2021): 8, <https://doi.org/10.1186/s11556-021-00263-z>.
 - [82] C. Tsigkanos, L. Gaskell, A. Smirniotou, and G. Tsigkanos, “Static and Dynamic Balance Deficiencies in Chronic Low Back Pain,” *Journal of Back and Musculoskeletal Rehabilitation* 29, no. 4 (2016): 887–893, <https://doi.org/10.3233/bmr-160721>.