



Neglected femoral neck fractures in cerebral palsy: a narrative review

Giuseppe Toro^{1,2}
Antimo Moretti¹
Marco Paoletta¹
Annalisa De Cicco¹
Adriano Braile¹
Alfredo Schiavone Panni¹

- Hip fractures are severe conditions with a high morbidity and mortality, especially when the diagnosis is delayed, and if formulated over 30 days after the injury, is termed a 'neglected femoral neck fracture' (NFNF).
- Cerebral palsy (CP) is probably one of the major risk factors for NFNF in Western countries, mainly because of both cognitive and motor impairments. However, considering the high prevalence of fractures in these patients, the incidence of NFNF in this population is probably underestimated, and this condition might result in persistent hip or abdominal pain.
- Several techniques are available for the treatment of NFNF (i.e. muscle pedicle bone graft, fixation with fibular graft, valgisation osteotomy), but most of them could affect motor function.
- Motor function must be preserved for as long as possible, in order to enhance the quality of life of CP patients.
- After discussing published NFNF cases in CP patients and available treatment options, a practical approach is proposed to facilitate the orthopaedic surgeon to both early identify and appropriately manage these challenging fractures.

Keywords: cerebral palsy; hip fracture; motor function; neglected femoral neck fracture; non-union

Cite this article: *EFORT Open Rev* 2020;5:58-64.

DOI: 10.1302/2058-5241.5.190019

Introduction

Hip fracture is a severe condition which is usually due to a fall in old and/or neurologically impaired patients, or due to high-energy trauma, when it is characterized by a

striking clinical picture with hip pain exacerbated by passive movements, inability to fully weight bear and high risk of mortality at short–medium term. Surgery is more successful when it is carried out earlier. However, in some patients this fracture is not diagnosed in a timely fashion, particularly in the case of intracapsular hip fracture. When the diagnosis is made at least 30 days after the injury, hip fracture is more appropriately defined as neglected femoral neck fracture (NFNF).¹

This condition is commonly described in developing countries,^{2–7} because of poverty, ignorance and lack of facilities,⁸ although it can be observed also in developed countries, especially in polytrauma.⁹ Although the frequency of NFNF is unknown, the incidence of this condition is probably expected to decline considering the improvements in healthcare facilities worldwide.¹⁰ As a consequence, the available literature is mostly about case series in non-disabled patients in underdeveloped countries.^{2–7} Disabled patients and especially patients with motor and cognitive impairment are at higher risk of sustaining an NFNF in developed countries.^{11–15}

Communicative impairment is one of the most relevant factors leading to a delay in fracture assessment in this population,^{12,16} because of the difficulties physicians face in obtaining a comprehensive medical history and physical examination.¹⁶ Typically most fractures in cerebral palsy (CP) occur after a low-energy trauma or simple mobilization of the patient,^{14,17} increasing the risk of underestimating patients' symptoms.

NFNF are difficult to treat, considering the loss of bone stock that could modify the femoral neck geometry, and femoral head microarchitecture, negatively affecting the union rate.⁵ Moreover, high risk of avascular necrosis (AVN) of the femoral head^{3,5,10} further influences both treatment options and surgical outcomes.

Therefore, the aim of our narrative review is to discuss published NFNF cases in CP, in order to identify core critical issues in this field and propose an assessment and therapeutic strategy to help the orthopaedic surgeon to appropriately manage affected patients.

CP and hip fractures

Patients with poor mobility and cognitive impairment are at high risk of misdiagnosis of hip fracture.¹³ CP is one of the most common causes of NFNF worldwide, considering the high incidence of osteoporosis due to decreased weight-bearing, endocrine alterations, inadequate calcium and vitamin D intake, and the exposure to medications (i.e. anticonvulsants) that adversely affect bone health.^{11,18,19} In non-ambulatory CP children, the low bone mass density (BMD) Z-score (-2 SD) and post-operative immobilization are risk factors for long bone fractures.²⁰ Moreover, CP is often associated with a high risk of falling due to postural control and gait impairments that further increase fracture risk.¹⁴ Maruyama et al¹⁴ observed a lifetime prevalence of fracture of 10% for disabled children. Furthermore, cognitive impairment affects 23–44% of CP patients.²¹ Considering also the high prevalence of communicative impairment (up to 80%),²¹ a comprehensive medical history is difficult to obtain in this population, leading to a high risk of NFNF.

According to Henderson et al¹² non-significant or unrecognized trauma, communication impairment and minimal displacement are the factors that could lead to a misdiagnosis of a hip fracture in CP. Bajelidze et al¹⁶ observed a positive bone scan in more than half of non-communicative CP patients with persistent pain. Furthermore, as showed by Mariani et al²² and Kenny and Parkinson²³ hip fractures could mimic abdominal pain in the same population. Therefore, it is mandatory to carefully evaluate persistent hip or abdominal pain, both clinically and radiographically^{16,24} in these patients.

As an effect of the extreme variability of the clinical picture of CP, and the lack of available evidence, NFNF are even more difficult to treat in this population. Kenny and Parkinson²³ published a case report of a 16-year-old boy with CP able to walk despite a spastic diplegia, although mentally disabled. The patient sustained bilateral spontaneous hip fractures, initially misdiagnosed because mimicking an abdominal pain. The boy was referred to the authors three months after symptoms began. At that time, he was not able to walk and X-rays showed bilateral subcapital hip fractures. Despite the fact that the patient received no treatment, five years after the diagnosis he was unable to walk, but was pain free.

Mariani et al²² reported a case of bilateral spontaneous hip fractures mimicking abdominal pain. The patient was

24 years old at the time of the fracture. He was a non-communicative CP patient with a Gross Motor Function Classification System (GMFCS) grade II. He was referred to the authors two months after symptoms began and X-rays showed bilateral Garden IV hip fractures. The patient was treated with a simultaneous bilateral total hip arthroplasty (THA) subsequently complicated by surgical wound infection. After debridement and antibiotics therapy he was discharged, and two years after the surgery he returned to pre-fracture functional status.

Toro et al reported a case of an NFNF which occurred in a 15-year-old boy with CP.²⁴ The patient's motor function was GMFCS grade II and, after a fall from a standing height, he started to complain of groin pain. After a clinical and radiographic examination, the patient was discharged with the diagnosis of hip bruise. However, because of the persistence of groin pain and the worsening of function, such that he could no longer sustain full weight bearing and walking without a walker few weeks later, he was referred to our institution three months after the injury. The X-rays showed a Sandhu type 2 NFNF on the affected side. Authors performed an open reduction and fixation with cannulated screws and augmentation with stem cells and cancellous bone graft. After an individual rehabilitation plan, the patient was pain free and able to walk without aids one year later.

Meyers et al,¹ in their series of 32 cases treated using muscle-pedicle bone grafting (MPBG), were the original authors who defined as 'neglected' a femoral neck fracture diagnosed 30 days after the injury.

The management of a neglected hip fracture in CP patients

NFNF in young patients are considered one of the most challenging femoral neck fractures to treat for orthopaedic surgeons.¹⁰ Sandhu et al³ classified the NFNF into three different types based on the shape of fracture surfaces, the proximal fragment length, fracture gap width and femoral head vitality (see Table 1 and Fig. 1 for further details). More recently, Jain et al⁷ summarized the available evidence in a literature review proposing a guide to treatment choice according to the Sandhu classification and the time elapsed since the fracture. The treatment of NFNF is a mechanobiological problem²⁵ and in recent decades several surgical techniques have been proposed for its treatment, including valgus intertrochanteric osteotomy (VIO), MPBG, and reduction and fixation with or without fibular grafting (FG).^{2–7,25}

Pivotal targets for treatment choice are femoral head vitality and proximal fragment size.^{3,7,10} In fact, hip preservation techniques are considered in all cases of viable femoral head,^{3,7,10} with a proximal fragment size of almost 2.5 cm.³ According to Ayoub and Gad,⁵ age (younger

Table 1. Sandhu classification

Sandhu type	Fracture surfaces	Proximal fragment	Fracture gap	Femoral head
1	regular	2.5 cm or more	< 1 cm	no signs of AVN
2	smooth and sclerosed	2.5 cm or more	1.0–2.5 cm	head of the femur still viable
3	smoothed	< 2.5 cm	> 2.5 cm	signs of AVN




Fig. 1 Diagnostic algorithm for persistent hip/abdominal pain in cerebral palsy.

than 30 years) and post-operative neck-shaft angle (lower than 130°) are the main predicting factors for the best outcomes of surgical joint preservation techniques. On the contrary, time elapsed since the trauma might affect micro and macroarchitectural deterioration of the proximal femur, influencing surgical decision-making.⁷

In CP patients with NFNF it is mandatory that the surgical treatment strategy has to be based on a comprehensive evaluation, including patient’s general condition, lower-limb muscle tone, joint contracture and gait impairment,¹⁷ with the ultimate goal of restoring pre-fracture functional status. In CP patients, motor function impairment is commonly classified according to the five stages of the GMFCS.^{26,27} Patients able to walk without aids but limited when walking outdoors should be included in the first two stages, whereas those in stages IV and V require assistive technology for most of their daily activities. Stage III is characterized by an intermediate motor impairment.

Patients classified as GMFCS IV or V show the highest risk of hip dislocation,^{28,29} that has to be considered in the planning of hip surgical procedures. In fact, even though associated with femoral/acetabular relocation

and derotation osteotomy, a joint-preserving technique might be associated with hip instability recurrence in these cases.³⁰ In these patients, assessment of lower-limb muscle tone, using the Tardieu scale, is required to better manage surgical planning and related outcomes.^{31,32}

The surgeon should consider that some of the techniques proposed for the treatment of NFNF could affect the lower-limb biomechanics resulting in worsening of motor function, especially in patients with GMFCS I to III. For example, an excessive valgus alignment after a VIO might negatively affect the abductor lever arm resulting in a persistent limp,³³ whereas the use of fibular graft is associated with gait impairment and ankle instability.³⁴ Therefore, the surgeon should carry out an adequate risk-benefit analysis, preferring less risky procedures (i.e. fixation with cancellous screws and bone graft). Recently, stem-cell transplant was proposed to treat fracture nonunions with encouraging results.^{5,35} Also, our experience in a clinical case treated with cannulated screws, bone graft and stem cells reported good results at one year follow-up.²⁴

Total hip arthroplasty (THA) might be the preferred treatment choice in CP patients with persistent hip pain with good long-term survivorship and a complication

rate comparable with THA in non-disabled patients.^{36,37} According to Houdek et al³⁷ GMFCS stage does not influence THA outcomes. In any case, the surgeon has to consider that performing THA in patients with a neurological disorder has some difficulties to face in terms of pre-operative counselling, altered acetabular and femoral anatomy, dealing with effects of previous surgery to the hip, the need for altered post-operative rehabilitation and a high rate of complications.^{38,39} Some technical precautions may help to improve THA outcomes in these patients.³⁹


Indeed, the use of dual-mobility cups further reduces the THA dislocation risk in this population⁴⁰ and some authors have associated femoral derotational osteotomy with THA in cases of dislocated painful hips.⁴¹ Therefore, hip replacement has been also proposed in ambulatory patients in the more severe GMFCS stages. Mariani et al²² reported a case of NFNF in CP treated with bilateral THA with good outcomes at two-year follow-up. In Sandhu type 3 NFNF, THA should be a treatment option also in CP ambulatory patients. A salvage procedure such as hip arthrodesis might be an alternative choice, especially in CP patients with normal contralateral hip and without structural and/or functional spine impairments,⁴² even if this procedure is associated with high complication rate.⁴³

For non-ambulatory patients or for those with a severe impairment of gross motor function (GMFCS V) the surgeon might also consider proximal femur resection techniques. Generally these procedures are indicated as salvage surgery to treat dislocated painful hip and to allow the patients to sit comfortably and to make perineal care

possible.⁴⁴⁻⁴⁷ Proximal femur resection techniques are generally associated with severe complications, whereas the Castle subtrochanteric resection-interposition arthroplasty (PFRIA) seems associated with more reliable surgical outcomes.^{48,49} In order to reduce the risk of bone-to-bone impingement when proximal migration of the femur occurs, Silverio et al⁵⁰ proposed the interposition of a shoulder prosthesis instead of soft tissues. Kenny and Parkinson,²³ in a case report, scheduled a CP patient with NFNF for proximal femur resection, but the surgery was not performed because of spontaneous pain resolution. According to both literature findings and our expertise, we propose viable options to treat NFNF in CP (Fig. 2).

Conclusions

Although hip fracture has a high incidence in patients with CP, diagnosis is frequently missed. In our opinion, hip fracture should be suspected in a case of persistent and/or unexplained hip or abdominal pain in this population to allow early fracture identification. When diagnosis is delayed for over 30 days, this condition is defined as NFNF, and for its treatment in CP patients the surgeon should consider the GMFCS stage, in order to formulate an appropriate surgical plan to achieve favourable outcomes. Surgical techniques that do not affect motor performance should be preferable to preserve ambulatory function. Total hip arthroplasty is a viable option in ambulatory patients with a Sandhu type 3 NFNF. In non-ambulatory patients, resection of the proximal femur might be indicated.






Fig. 2 Comprehensive algorithm for the treatment of neglected femoral neck fracture (NFNF) in cerebral palsy (CP) according to Sandhu type and GMFCS. (a) Algorithm for the treatment of Sandhu type 1 NFNF. (b) Algorithm for the treatment of Sandhu type 2 NFNF. (c) Algorithm for the treatment of Sandhu type 3 NFNF.

Note. GMFCS, Gross Motor Function Classification System; THA, total hip arthroplasty; PFRIA, Castle subtrochanteric resection-interposition arthroplasty; MPBG, muscle-pedicle bone grafting.

AUTHOR INFORMATION

¹Department of Medical and Surgical Specialties and Dentistry, University of Campania 'Luigi Vanvitelli', Naples, Italy.

²Department of Clinical Sciences and Translational Medicine, University of Rome Tor Vergata, Rome, Italy.

Correspondence should be sent to: Giuseppe Toro, University of Campania 'Luigi Vanvitelli', Department of Medical and Surgical Specialties and Dentistry, Via L. De Crecchio 4, 80138 Naples, Italy.
Email: giusep.toro@gmail.com

ICMJE CONFLICT OF INTEREST STATEMENT

GT reports payment for expert testimony, grants/grants pending, and travel/accommodation/meeting expenses unrelated to activities listed, all outside the submitted work.

The other authors declare no conflict of interest relevant to this work.

FUNDING STATEMENT

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

LICENCE

© 2020 The author(s)

This article is distributed under the terms of the Creative Commons Attribution-Non Commercial 4.0 International (CC BY-NC 4.0) licence (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed.

REFERENCES

- Meyers MH, Harvey JP Jr, Moore TM.** Delayed treatment of subcapital and transcervical fractures of the neck of the femur with internal fixation and a muscle pedicle bone graft. *Orthop Clin North Am.* 1974;5:743–756.
- Huang CH.** Treatment of neglected femoral neck fractures in young adults. *Clin Orthop Relat Res.* 1986;117–126.
- Sandhu HS, Sandhu PS, Kapoor A.** Neglected fractured neck of the femur: a predictive classification and treatment by osteosynthesis. *Clin Orthop Relat Res.* 2005;14–20.
- Magu NK, Singh R, Sharma AK, Ummat V.** Modified Pauwels' intertrochanteric osteotomy in neglected femoral neck fractures in children: a report of 10 cases followed for a minimum of 5 years. *J Orthop Trauma.* 2007;21:237–243.
- Ayoub MA, Gad HM.** Neglected neck femur fractures in adolescents and young adults: factors predicting the surgical outcome. *J Orthop Sci.* 2013;18:93–100.
- Nair N, Patro DK, Babu TA.** Role of muscle pedicle bone graft as an adjunct to open reduction and internal fixation in the management of neglected and ununited femoral neck fracture in young adults: a prospective study of 17 cases. *Eur J Orthop Surg Traumatol.* 2014;24:1185–1191.
- Jain AK, Mukunth R, Srivastava A.** Treatment of neglected femoral neck fracture. *Indian J Orthop.* 2015;49:17–27.
- Nagi ON, Dhillon MS, Goni VG.** Open reduction, internal fixation and fibular autografting for neglected fracture of the femoral neck. *J Bone Joint Surg Br.* 1998;80:798–804.
- Pauyo T, Drager J, Albers A, Harvey EJ.** Management of femoral neck fractures in the young patient: A critical analysis review. *World J Orthop.* 2014;5:204–217.
- Roshan A, Ram S.** The neglected femoral neck fracture in young adults: review of a challenging problem. *Clin Med Res.* 2008;6:33–39.
- Henderson RC, Lark RK, Gurka MJ, et al.** Bone density and metabolism in children and adolescents with moderate to severe cerebral palsy. *Pediatrics.* 2002;110:e5.
- Henderson RC, Berglund LM, May R, et al.** The relationship between fractures and DXA measures of BMD in the distal femur of children and adolescents with cerebral palsy or muscular dystrophy. *J Bone Miner Res.* 2010;25:520–526.
- Hossain M, Akbar SA, Andrew G.** Misdiagnosis of occult hip fracture is more likely in patients with poor mobility and cognitive impairment. *Acta Orthop Belg.* 2010;76:341–346.
- Maruyama K, Nakamura K, Nashimoto M, et al.** Bone fracture in physically disabled children attending schools for handicapped children in Japan. *Environ Health Prev Med.* 2010;15:135–140.
- Iwamoto J, Takeda T, Matsumoto H.** Efficacy of oral bisphosphonates for preventing hip fracture in disabled patients with neurological diseases: a meta-analysis of randomized controlled trials among the Japanese population. *Curr Med Res Opin.* 2011;27:1141–1148.
- Bajelidze G, Belthur MV, Littleton AG, Dabney KW, Miller F.** Diagnostic evaluation using whole-body technetium bone scan in children with cerebral palsy and pain. *J Pediatr Orthop.* 2008;28:112–117.
- Persiani P, Murgia M, Ranaldi FM, et al.** The treatment of femoral fractures in children with cerebral palsy. *Clin Ter.* 2018;169:e18–e22.
- Leet AI, Mesfin A, Pichard C, et al.** Fractures in children with cerebral palsy. *J Pediatr Orthop.* 2006;26:624–627.
- Ozel S, Switzer L, Macintosh A, Fehlings D.** Informing evidence-based clinical practice guidelines for children with cerebral palsy at risk of osteoporosis: an update. *Dev Med Child Neurol.* 2016;58:918–923.
- Ko CH, Tse PWT, Chan AKH.** Risk factors of long bone fracture in non-ambulatory cerebral palsy children. *Hong Kong Med J.* 2006;12:426–431.
- Oding E, Roebroek ME, Stam HJ.** The epidemiology of cerebral palsy: incidence, impairments and risk factors. *Disabil Rehabil.* 2006;28:183–191.
- Mariani P, Buttaro M, Comba F, Zanotti E, Ali P, Piccaluga F.** Bilateral simultaneous femoral neck fracture mimicking abdominal pain in a cerebral palsy patient. *Case Rep Orthop.* 2014;925201.
- Kenny NW, Parkinson RW.** Simultaneous bilateral hip fractures presenting as an acute abdomen. *Injury.* 1993;24:691–692.
- Toro G, Moretti A, Toro G, et al.** Surgical treatment of neglected hip fracture in children with cerebral palsy: case report and review of the literature. *Clin Cases Miner Bone Metab.* 2017;14:317–323.
- Gadegone WM, Ramteke AA, Lokhande V, Salphade Y.** Valgus intertrochanteric osteotomy and fibular strut graft in the management of neglected femoral neck fracture. *Injury.* 2013;44:763–768.
- Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B.** Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol.* 1997;39:214–223.

- 27. Palisano RJ, Rosenbaum P, Bartlett D, Livingston MH.** Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol.* 2008;50:744–750.
- 28. Albiñana J, Gonzalez-Moran G.** Painful spastic hip dislocation: proximal femoral resection. *Iowa Orthop J.* 2002;22:61–65.
- 29. Root L.** Surgical treatment for hip pain in the adult cerebral palsy patient. *Dev Med Child Neurol.* 2009;51:84–91.
- 30. Bayusentono S, Choi Y, Chung CY, Kwon S-S, Lee KM, Park MS.** Recurrence of hip instability after reconstructive surgery in patients with cerebral palsy. *J Bone Joint Surg Am.* 2014;96:1527–1534.
- 31. Tardieu G, Shentoub S, Delarue R.** [Research on a technic for measurement of spasticity]. *Rev Neurol (Paris).* 1954;91:143–144.
- 32. Haugh AB, Pandyan AD, Johnson GR.** A systematic review of the Tardieu Scale for the measurement of spasticity. *Disabil Rehabil.* 2006;28:899–907.
- 33. Varghese VD, Boopalan PR, Titus VTK, Oommen AT, Jepeganmam TS.** Indices affecting outcome of neglected femoral neck fractures after valgus intertrochanteric osteotomy. *J Orthop Trauma.* 2014;28:410–416.
- 34. Ling XF, Peng X.** What is the price to pay for a free fibula flap? A systematic review of donor-site morbidity following free fibula flap surgery. *Plast Reconstr Surg.* 2012;129:657–674.
- 35. Giannoudis PV, Ahmad MA, Mineo GV, Tosounidis TI, Calori GM, Kanakaris NK.** Subtrochanteric fracture non-unions with implant failure managed with the “Diamond” concept. *Injury.* 2013;44:S76–S81.
- 36. Raphael BS, Dines JS, Akerman M, Root L.** Long-term followup of total hip arthroplasty in patients with cerebral palsy. *Clin Orthop Relat Res.* 2010;468:1845–1854.
- 37. Houdek MT, Watts CD, Wyles CC, Trousdale RT, Milbrandt TA, Taunton MJ.** Total hip arthroplasty in patients with cerebral palsy: A cohort study matched to patients with osteoarthritis. *J Bone Joint Surg Am.* 2017;99:488–493.
- 38. Ries MD, Wolff D, Shaul JA.** Hip arthroplasty in mentally impaired patients. *Clin Orthop Relat Res.* 1994;146–154.
- 39. Konan S, Duncan CP.** Total hip arthroplasty in patients with neuromuscular imbalance. *Bone Jt J.* 2018 Jan;100-B:17–21.
- 40. Sanders RJM, Swierstra BA, Goosen JHM.** The use of a dual-mobility concept in total hip arthroplasty patients with spastic disorders: no dislocations in a series of ten cases at midterm follow-up. *Arch Orthop Trauma Surg.* 2013;133:1011–1016.
- 41. Prosser GH, Shears E, O’Hara JN.** Hip resurfacing with femoral osteotomy for painful subluxed or dislocated hips in patients with cerebral palsy. *J Bone Joint Surg Br.* 2012;94:483–487.
- 42. Fucs PM, Yamada HH.** Hip fusion as hip salvage procedure in cerebral palsy. *J Pediatr Orthop.* 2014;34:S32–S35.
- 43. Kolman SE, Ruzbarsky JJ, Spiegel DA, Baldwin KD.** Salvage Options in the Cerebral Palsy Hip: A Systematic Review. *J Pediatr Orthop.* 2016;36:645–650.
- 44. Castle ME, Schneider C.** Proximal femoral resection-interposition arthroplasty. *J Bone Joint Surg Am.* 1978;60:1051–1054.
- 45. McCarthy RE, Simon S, Douglas B, Zawacki R, Reese N.** Proximal femoral resection to allow adults who have severe cerebral palsy to sit. *J Bone Joint Surg Am.* 1988;70:1011–1016.
- 46. Dartnell J, Gough M, Paterson JMH, Norman-Taylor F.** Proximal femoral resection without post-operative traction for the painful dislocated hip in young patients with cerebral palsy: a review of 79 cases. *Bone Joint J.* 2014;96-B:701–706.
- 47. K Graham H, Narayanan UG.** Salvage hip surgery in severe cerebral palsy: some answers, more questions? *Bone Joint J.* 2014;96-B:567–568.
- 48. Toro G, Florio P, Gagliardo N, et al.** Proximal femoral resection-arthroplasty in severe cerebral palsy. *Osteoporos Int* (2015) 26:S318–S318.
- 49. Hwang JH, Varte L, Kim HW, Lee DH, Park H.** Salvage procedures for the painful chronically dislocated hip in cerebral palsy. *Bone Joint J.* 2016;98-B:137–143.
- 50. Silverio AL, Nguyen SV, Schlechter JA, Rosenfeld SR.** Proximal femur prosthetic interposition arthroplasty for painful dislocated hips in children with cerebral palsy. *J Child Orthop.* 2016;1:657–664.