

Complications in Brow Lifts: A Systemic Review of Surgical and Nonsurgical Brow Rejuvenations

Min-Jeong Cho, MD*
 Jourdan A. Carboy, BS*
 Rod J. Rohrich, MD†

Background: Brow lift has been a part of plastic surgeons' armamentarium for over a century. Although there are multiple different techniques available, there has been a dearth of literature comparing which technique is better and has lesser complications. In this study, we performed a systemic literature review of all brow lift techniques to determine and compare complication rates of each technique.

Methods: We performed a computerized search of brow rejuvenation techniques using the MEDLINE database. Data regarding the type of brow lift procedure, number of patients, and complications were collected.

Results: The systematic review was performed in December of 2017. A total of 326 articles were identified initially, and 76 studies remained after final review. The review showed highest revision rate in the hairline brow lift (7.4%), highest numbness rate in the direct brow lift (5.5%), highest asymmetry rate in the temporal/lateral brow lift (1.5%), and highest alopecia rate in the endoscopic brow lift (2.8%). In addition, we found 5 studies in nonsurgical brow rejuvenation (3 botulinum toxin, 1 fat injection, 1 radiofrequency).

Conclusion: Our study shows that each brow technique has a different complication profile, and it is important for plastic surgeons to understand the difference. Furthermore, the study demonstrated the lack of randomized prospective outcome studies and standardization of outcome measures in brow rejuvenation. We believe this information would be the key and next frontier in providing technically safe and long-lasting aesthetic outcomes for the patients. (*Plast Reconstr Surg Glob Open* 2018;6:e1943; doi: 10.1097/GOX.0000000000001943; Published online 15 October 2018.)

INTRODUCTION

Brow lift has been a part of plastic surgeons' armamentarium for over a century. Over time, the techniques have evolved from the most direct approach such as skin excision of upper forehead skin by Passot¹ in 1919 to minimally invasive lift using an endoscopic approach by Isse in 1994². Currently, plastic surgeons have various types of techniques available for their disposal: direct, hairline, temporal, lateral, endoscopic, and transblepharoplasty brow lift³⁻⁸ (Fig. 1).

Although there are multiple different techniques available, there has been a dearth of literature comparing the complication profile of these techniques.⁹ It is critical for

plastic surgeons to understand the complication profile of each technique and anticipate possible postoperative complications for each technique. Therefore, we present a systemic review of all brow rejuvenation techniques.

PATIENTS AND METHODS

A computerized search of the MEDLINE database was performed using OVID. The OVID search was performed using the following search terms and including all sub-headings:

- Brow lift
- Brow rejuvenation
- Periorbital rejuvenation
- Periocular rejuvenation

The following limits were applied for each search term:

- Language: English
- Dates: no limits

From the *Department of Plastic Surgery, University of Texas Southwestern Medical Center, Dallas, Tex.; and †Dallas Plastic Surgery Institute, Dallas, Tex.

Received for publication May 19, 2018; accepted July 27, 2018.

Copyright © 2018 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/GOX.0000000000001943

Disclosure: Rod J. Rohrich, M.D., receives instrument royalties from Eriem Surgical, Inc., and book royalties from Thieme Medical Publishing; he is a Clinical and Research Expert for Allergan, Inc., and MTF Biologics, and the Owner of Medical Seminars of Texas, LLC. No funding was received for this article. Dr. Cho has no financial interest to declare in relation to the content of this article. The Article Processing Charge was paid for by the authors.




Fig. 1. Brow lift evolution.

The resulting articles were reviewed using the following inclusion and exclusion criteria.

Inclusion criteria

- Controlled studies, cohort studies, randomized controlled trials, systemic reviews

Exclusion criteria

- Case reports, review articles, case series, expert opinions
- Publications not in the English language

All studies were reviewed and analyzed by one of the authors (M.J.C.). Data regarding the type of brow lift procedure, number of patients, complications, and outcomes were collected. We categorized articles into 3 groups: endoscopic brow lift, open brow lift, and nonsurgical brow lift. The open brow lift articles were categorized into coronal, hairline, direct, temporal/lateral, and transblepharoplasty brow lifts.

RESULTS

A Systematic Literature Review

The computerized search was performed in December 4 of 2017. A total of 326 articles were identified from the initial search: brow lift, n = 253; brow rejuvenation, n = 6; periorbital rejuvenation, n = 61; and periocular rejuvenation, n = 6. The abstracts of each article were reviewed, and 139 studies of potential relevance remained: brow lift, n = 106; brow rejuvenation, n = 6; periorbital rejuvenation, n = 21; and periocular rejuvenation, n = 6. Then, each article was reviewed, and final 76 studies remained: coronal brow lift, n = 9; endoscopic brow lift, n = 34; direct brow lift, n = 10; temporal/lateral brow, n = 10; hairline brow lift, n = 4; transblepharoplasty brow lift, n = 4; and non-surgical brow lift, n = 5. In total, 94.4% of studied subjects were female, whereas the other 5.6% were male.

Open Brow Lift

For open brow lift, we identified 37 studies from the literature review (2,858 patients). Of the 37 studies, the direct brow lift had the most number of the studies (10 studies), followed by coronal (9), temporal (6), lateral (4), hairline (4), and transblepharoplasty (4).

In comparison, the revision rate was the highest in the hairline brow lift (7.4%), followed by 3.6% in the direct brow lift, 2.4% in the temporal brow lift, 1.8% in the coronal brow lift, and 0.1% in the transblepharoplasty (Tables 1, 2). For the rate of numbness, it was the highest in the direct brow lift (5.5%), followed by the transblepharoplasty (2.1%), and the temporal/lateral brow lift

Table 1. Open Brow Lift Complications

Coronal		
Complication	No. Patients	Percentage (n = 449)
Alopecia	10	2.2
Chemosis	1	0.2
Frontal paralysis	1	0.2
Hematoma	2	0.4
Pruritus	1	0.2
Recurrence	1	0.2
Revision	8	1.8
Visible scar	5	1.1
Direct		
Complication	No. Patients	Percentage (n = 439)
Asymmetry	4	0.9
Eye problem	1	0.2
Nerve injury	1	0.2
Numbness	24	5.5
Overcorrection	3	0.7
Revision upper blepharoplasty due to ptosis	16	3.6
Recurrence	1	0.2
Visible scar	9	2.1
Hairline		
Complication	No. Patients	Percentage (n = 374)
Flap necrosis	3	0.7
Paresthesia	7	1.7
Revision	30	7.4
Widened scar	1	0.2
Temporal/Lateral		
Complication	No. Patients	Percentage (n = 780)
Alopecia	6	0.8
Asymmetry	12	1.5
Bruising	4	0.5
Chronic irritation	1	0.1
Frontal palsy	2	0.3
Hematoma	3	0.4
Infection	16	2.1
Numbness	2	0.3
Palpable suture	2	0.3
Persistent edema	1	0.1
Recurrence	4	0.5
Revision	9	1.2
Transblepharoplasty		
Complication	No. Patients	Percentage (n = 816)
Asymmetry	6	0.8
Dehiscence	3	1.5
Dysesthesia	2	0.5
Eye problem	8	0.1
Hematoma	1	0.3
Lymphedema	1	0.4
Numbness	8	2.1
Revision	1	0.3
Residual ptosis	11	0.3
Suture granuloma	2	0.1
Visible	4	0.5

Table 2. Comparison of Complication Rates

Complication	Complication Rate (%)	Type of Brow Lift
Alopecia	2.8	Highest in endoscopic
	0.8	Lowest in temporal/lateral
Asymmetry	1.5	Highest in temporal/lateral
	0.7	Lowest in transblepharoplasty and endoscopic
Numbness	5.5	Highest in direct
	0.3	Lowest in temporal/lateral
Revision	7.4	Highest in hairline
	0.1	Lowest in transblepharoplasty

Table 3. Endoscopic Brow Lift Complications

Complication	Endoscopic	
	No. Patients	Percentage (n = 7,273)
Alopecia	204	2.8
Asymmetry	54	0.7
Edema	18	0.2
Dehiscence	1	0.0
Eye problem	12	0.2
Glabellar irregularity	2	0.0
Granuloma	1	0.0
Hematoma	7	0.1
Infection	4	0.1
Nerve injury	4	0.1
Numbness	148	2.0
Pain	8	0.1
Palpability	21	0.3
Pruritus	37	0.5
Recurrence	9	0.1
Revision	86	1.2
Screw exposure	1	0.0
Seroma	1	0.0
Skin burn	1	0.0
Visible scar	3	0.0

(0.3%). The rate of asymmetry was the highest in the temporal/lateral brow lift (1.5%), followed by the direct brow lift (0.9%), and the transblepharoplasty (0.7%). Lastly, the rate of alopecia was the highest in the coronal brow lift (2.2%), followed by the temporal/lateral brow lift (1.5%).

Endoscopic Brow Lift

A total of 34 studies were identified (7,273 patients) for the endoscopic brow lift (Table 3). The most common complication was alopecia (2.8%), followed by numbness (2%), revision (1.2%), asymmetry (0.7%), pruritus (0.5%), palpability (0.3%), edema (0.2%), eye-related complication (0.2%), and hematoma/infection/nerve injury/pain/recurrence (0.1%).

Nonsurgical Brow Rejuvenation

The literature review revealed 5 outcome studies in nonsurgical brow rejuvenation: botulinum toxin (3), fat injection (1), and radiofrequency (1; Table 4). The review showed that the most common complication in botulinum toxin was bruising (1.7%), flu-like symptoms/persistent wrinkles/trace ptosis (0.7%), and eyelid ptosis/excessive elevation (0.3%). For the fat injection, infection was the only documented complication (0.4%), and there were no complications documented for radiofrequency brow lift.

Table 4. Nonsurgical Brow Rejuvenation Complications

Complication	Botulinum Toxin	
	No. Patients	Percentage (n = 297)
Bruising	5	1.7
Eyelid ptosis	1	0.3
Excessive elevation	1	0.3
Flu-like symptoms	2	0.7
Persistent wrinkles	2	0.7
Trace ptosis	2	0.7
Complication	Radiofrequency	
	No. Patients	Percentage (n = 125)
No complication	125	100
Complication	Fat	
	No. Patients	Percentage (n = 250)
Infection	1	0.4

DISCUSSION

Reflecting the current trend toward minimally invasive procedures in cosmetic surgery, brow rejuvenation has undergone several transformations past decades.^{1,2,5-8} It has evolved from most invasive approach such as coronal brow lift to endoscopic approach to botulinum toxin injection¹⁰⁻¹² (Fig. 1). Although there are various types of techniques available, very little is known about the relative complication profile of each technique, and how each technique differs from the other approach except few retrospective and cohort studies.^{9,13,14} Therefore, we performed a systemic literature review of all brow lift techniques to determine and compare complication rates of each technique.

Our study shows that the majority of complication rate was minimal (< 5%), and each technique has a different complication profile (Table 1). For the open brow lift techniques, we reviewed 37 studies (2,858 patients), and found the following: (1) numbness (5.5%) was the highest in the direct brow lift; (2) revision rate (7.4%) was the highest in the hairline brow lift; (3) asymmetry rate (1.5%) was the highest in the temporal brow lift; and (4) alopecia rate (2.2%) was the highest in the coronal brow lift. So far, there has been only one retrospective study comparing open brow techniques,⁹ and the sample size for each open brow technique was very small (< 10 patients). In this study, they found that alopecia and sensory-related complications were the highest in the coronal brow lift.⁹ Except for our study and the study by Koch et al.,⁹ the majority of studies compare open versus endoscopic brow lift by combining all open approaches as a one group.^{3,15}

For the endoscopic brow lift, we reviewed 34 studies (7,273 patients) and found that the patients experienced lower complications rates than the open brow lift. The most common complication rate was alopecia (2.8%), followed by numbness (2%), and revision (1.2%). The remaining complication rates were < 1% in asymmetry, pruritus, palpability, edema, eye-related complications, hematoma, infection, nerve injury, pain, and recurrence. This finding is in agreement with our earlier study of open versus endoscopic brow lift.³ In the previous study, we found dysesthesia and alopecia were the common complications in the endoscopic brow lift.




Fig. 2. Trend of brow rejuvenation from 2000 to 2015.

Interestingly, our review revealed that there are only 5 studies (672 patients) on nonsurgical brow rejuvenation: botulinum toxin (3), fat injection (1), and radiofrequency (1). According to American Society of Plastic Surgeons, brow rejuvenation by botulinum toxin has become exponentially popular (797%) past 16 years, whereas the percentage of surgical brow rejuvenation has decreased about 64% during the same period (Fig. 2). Despite the increased popularity of nonsurgical brow rejuvenation, we found a strikingly small number of outcome studies on its use.¹⁶ Nevertheless, the analysis shows that these techniques are associated with minor complications (bruising, flu-like symptoms, persistent wrinkles, trace ptosis, infection).^{10,11,17,18} In addition, the rates of recurrence and revision rates are unknown.

The limitation of our study is the inconsistency in the types of complications reported in the studies. Each study reported different types of complications and had different follow-up periods. Therefore, we were not able to compare a specific type of complication rate in all types of brow lift techniques. In addition, there was no standardization of each type of brow lift. Each study added their own modifications to historical described direct, coronal, temporal/lateral, transblepharoplasty, endoscopic, and botulinum toxin brow lift. Therefore, it is difficult to determine if certain technical modification resulted in lower or higher complications. Lastly, our study is a limited systemic review without a meta-analysis. The results from the identified brow lift studies were not matched using biostatistical techniques.

As demonstrated, our study shows the lack of randomized prospective outcome studies and standardization of outcome measures in brow rejuvenation. As there is a lack

of high-quality evidence-based medicine in aesthetic surgery, we believe future prospective studies with standardized surgical and aesthetic outcome measures of brow rejuvenation are necessary.¹⁹ This information would be the key to provide technically safe and long-lasting aesthetic outcomes for the patients.

CONCLUSIONS

The brow lift approach has transformed from the most invasive technique such as a coronal brow lift to the minimally invasive technique such as botulinum toxin injection. Although there are countless techniques and modifications available at our disposal, there is a limited number of studies on the complication profile of each technique. Our study shows that each brow technique has a different complication profile and it is important for plastic surgeons to understand the difference. Furthermore, the study demonstrated the lack of randomized prospective outcome studies and standardization of outcome measures in brow rejuvenation. We believe this information would be the key and next frontier in providing technically safe and long-lasting aesthetic outcomes for the patients.

Rod J. Rohrich, MD

Dallas Plastic Surgery Institute
9101 North Central Expressway
Suite 600
Dallas, TX 75231
E-mail: rod.rohrich@dpsi.org

REFERENCES

1. Passot R. Chirurgie Esthétique des rides du visage. *Presse Med.* 1919;27:258.
2. Isse NG. Endoscopic facial rejuvenation: endoforehead, the functional lift. Case reports. *Aesthetic Plast Surg.* 1994;18:21–29.
3. Graham DW, Heller J, Kurkjian TJ, et al. Brow lift in facial rejuvenation: a systematic literature review of open versus endoscopic techniques. *Plast Reconstr Surg.* 2011;128:335e–341e.
4. Walrath JD, McCord CD. The open brow lift. *Clin Plast Surg.* 2013;40:117–124.
5. Flowers RS. Periorbital aesthetic surgery for men. Eyelids and related structures. *Clin Plast Surg.* 1991;18:689–729.
6. Regnault P. [Complete face and forehead lifting, with double traction on “crow’s-feet”]. *Plast Reconstr Surg.* 1972;49:123–129.
7. Hunt H. *Plastic Surgery of the Head, Face, and Neck.* Philadelphia, Pa.: Lea&Febiger; 1926.
8. Miller C. *Cosmetic Surgery.* 2nd ed. Chicago, Ill.: Oak Printing and Publishing; 1908.
9. Koch RJ, Troell RJ, Goode RL. Contemporary management of the aging brow and forehead. *Laryngoscope.* 1997;107:710–715.
10. Ahn MS, Catten M, Maas CS. Temporal brow lift using botulinum toxin A. *Plast Reconstr Surg.* 2000;105:1129–1135; discussion 1136.
11. Maas CS, Kim EJ. Temporal brow lift using botulinum toxin A: an update. *Plast Reconstr Surg.* 2003;112:109S–112S; discussion 113S.
12. Steinsapir KD, Rootman D, Wulc A, et al. Cosmetic microdroplet botulinum toxin A forehead lift: a new treatment paradigm. *Ophthalmic Plast Reconstr Surg.* 2015;31:263–268.
13. Roberts TL 3rd, Ellis LB. In pursuit of optimal rejuvenation of the forehead: endoscopic brow lift with simultaneous carbon dioxide laser resurfacing. *Plast Reconstr Surg.* 1998;101:1075–1084.
14. Nassif PS, Kokoska MS, Homan S, et al. Comparison of subperiosteal vs subgaleal elevation techniques used in forehead lifts. *Arch Otolaryngol Head Neck Surg.* 1998;124:1209–1215.
15. Tabatabai N, Spinelli HM. Limited incision nonendoscopic brow lift. *Plast Reconstr Surg.* 2007;119:1563–1570.
16. 2017 Plastic Surgery Statistics Report. *ASPS National Clearinghouse of Plastic Surgery Procedural Statistics.* 2017.
17. Koch RJ. Radiofrequency nonablative tissue tightening. *Facial Plast Surg Clin North Am.* 2004;12:339–346, vi.
18. Kranendonk S, Obagi S. Autologous fat transfer for periorbital rejuvenation: indications, technique, and complications. *Dermatol Surg.* 2007;33:572–578.
19. Rohrich RJ, Cho MJ. Evidence-based medicine in aesthetic surgery: the significance of level to aesthetic surgery. *Plast Reconstr Surg.* 2017;139:1195e–1202e.