

SHORT REPORT

Open Access

The complete *alk* sequences of *Rhodococcus erythropolis* from Lake Baikal

Alexander Likhoshvay, Anna Lomakina* and Mihail Grachev

Abstract

Background: Rhodococci are bacteria able to degrade a wide range of hydrocarbons, including the alkanes present in crude oil, due to *alk* genes in their genomes.

Findings: Genome sequencing of DNA from *Rhodococcus erythropolis* strain 4 (obtained from a deep-water bitumen mound) revealed four *alk* genes, and the predicted amino acid sequences coded by these genes were highly conserved, having sections up to 11 amino acid residues.

Conclusions: Obtained four genes from *Rhodococcus erythropolis* were similar to corresponding genes from other bacteria collected from other environments, including marine sources. This indicated a large-scale horizontal *alk* gene transfer between bacteria from different subgenera.

Keywords: Lake Baikal; Natural oil seep; Alkane hydroxylases; *Rhodococcus erythropolis*

Background

Alkanes may constitute up to 88% of the volume present in natural oil, and due to their high toxicity can serve as a convenient source of energy only for oil-degrading microorganisms (van Beilen et al. 2003).

To overcome this obstacle, bacteria have learned to synthesize specific enzymes for extracting energy from *n*-alkanes. The initial step in *n*-alkane oxidation is catalyzed by a monooxygenase complex composed of an alkane hydroxylase (*alkB*), rubredoxin reductase (*alkT*) and rubredoxin (*alkG*, an electron carrier), which are known to play an important role in oil bioremediation (Kloos et al. 2006; van Beilen et al. 2003; van van Beilen and Funhoff 2007). Biodegradation starts with cleavage of the C-H bond, catalysed by the oxygenase-group enzyme alkane hydroxylase, which inserts an atom of oxygen from O₂ into the hydrocarbon molecule. Prior to this catalysis, the enzyme must be activated by NADH (nicotinamide adenine dinucleotide) which transfers a pair of electrons from FAD (flavin adenine dinucleotide) to rubredoxin. One pair of electrons is transferred to alkane hydroxylase leading to the formation of primary or secondary alcohols [R-CH₃ + O₂ + NAH(P)H + H⁺ → R-CH₂OH + NAD(P)⁺ + H₂O], which

are further converted to dicarboxylic acids (van Beilen et al. 2003).

This enzyme system was originally discovered in *Pseudomonas putida* and further research found that the genes encoding for alkane-degrading enzymes (*alk*-genes, rubredoxin, rubredoxin reductase) are located on a plasmid or chromosomes (van Beilen et al. 2001). Bacteria from different genera, including *Rhodococcus*, possess similar enzyme systems and *alk*-genes (Whyte et al. 1998; Whyte et al. 2002). Members of the genus *Rhodococcus* seem to play significant role in bioremediation of oil spills (Whyte et al. 2002) and are recognized as ideal candidates for the biodegradation of hydrocarbons due to their ability to degrade a wide range of organic compounds (Beard and Page 1998), their hydrophobic cell surface and the production of biosurfactants as well as their ubiquity and robustness in the environment (Larkin et al. 1998; Warhurst and Fewson 1994).

Lake Baikal, the deepest (1637 m) and oldest (25 mln y) lacustrine reservoir on Earth, is located in the middle of Eurasia. During a 2008 exploration using Mir submersibles, natural oil seepages surrounded with "bitumen mounds" were discovered on the lake bottom (Khlystov et al. 2009). These structures are stable, inhabited by living creatures and persist even if the source of oil is depleted. One of these bitumen mounds (No. 8) contained 148 mg/g of aliphatic C₂₂-C₃₄ *n*-alkanes (primarily C₂₅) where several

* Correspondence: lomakina@lin.irk.ru
Limnological Institute of the Siberian Branch of the Russian Academy of Sciences Ulan-Batorskaya 3, Irkutsk 664033, Russia

strains of bacteria were isolated and later identified as *Rhodococcus erythropolis* (Likhoshvay et al. 2013) by means of 16S rRNA analysis.

This study aimed to identify *alk* genes in the genome of one of these isolates (strain 4) by nucleotide and complete genome sequence analysis. Strain 4, identified as *R. erythropolis*, has four *alk*-genes which differed from each other, but were similar to corresponding genes in bacteria from other habitats.

Materials and methods

Bacterial strain

R. erythropolis strain 4 (Acc. No HQ702471), isolated from bitumen mound 8 at the natural oil seep near Cape of Gorevoi Utes (10 km offshore, depth 900 m, Central Baikal) (Likhoshvay et al. 2013).

DNA extraction and sequencing analysis

DNA was extracted by the method of Sambrook et al. (1989) with minor modifications - enzymatic lysis followed by phenol-chloroform extraction.

Complete genome sequencing of DNA was carried out according to the manual/protocol provided with the Illumina GAIIx (India). Number of readings equalled to approximately 10 Mbp. Reassembling of individual nucleotide sequences by Velvet_1.1.02 resulted to 3897 contigs with an average length of 1.8 Kbp and a total length of 6.9 Mbp. The nucleotide sequences of the *alkB* genes were translated into amino acid sequences by the Expsay Translate Tool (<http://web.expasy.org/tools/translate/>) and uploaded to the NCBI data base with the following accession numbers: alkane hydroxylase 1 (KF498365), alkane hydroxylase 2 (KF498366), alkane hydroxylase 3 (KF498367), alkane hydroxylase 4 (KF498368).

Homology between the four sequences was estimated by BLASTX (<http://blast.st-va.ncbi.nlm.nih.gov/Blast.cgi>) where the nucleotide sequences and inferred amino acid sequences were aligned with homologous sequences retrieved from GenBank using the CLUSTAL W software. A phylogenetic tree for the genes was constructed by the neighbor-joining method (Saitou and Nei 1987)

using the MEGA4 program (Tamura et al. 2007). The relative synonymous codon usage (RSCU) was computed for the *alkB* genes and correspondence analysis was performed using CODONW software.

Results

Examination of the complete genome DNA sequence of *R. erythropolis* strain 4 (isolated from the inner part of a bitumen mound) revealed four *alk* genes: *alk1* was 386, *alk2* was 389, *alk3* was 408 and *alk4* was 400 amino acids in length. All four sequences for these alkane hydroxylases had homologous, synonymous and significant substitutions (Figure 1). The highly conserved sequences, typical of amino acid sequences (HE[L/M]xHK, EHxxGHH, LQRH[S/A]DHHA) from alkane-degrading bacteria (van Beilen et al. 2005), were found to have the following sequences - HELGHK, EHNxGHH and LQRHSDHHA. According to data obtained by phylogenetic analysis (Figure 2), the alkane hydroxylases were located on four different branches of the tree with corresponding sequences from other rhodococci.

Complete amino acid sequences obtained from *R. erythropolis* strain 4 were compared to the recently published genome (NCBI) of *R. erythropolis* PR4 (Sekine et al. 2006) (Figure 3) and found to be nearly identical: alkane hydroxylase 1–9 substitutions including 6 synonymous (97.6% identity); alkane hydroxylase 2–1 nonsynonymous substitution (99.7% identity); alkane hydroxylase 3–5 substitutions including 4 synonymous (98.7% identity); alkane hydroxylase 4–2 synonymous, 3 nonsynonymous substitutions and 5 inserts of 2–3 amino acids (94.2% identity).

Comparative analysis of a wide range of homologous eubacterial sequences (NCBI) revealed that the genome of *R. erythropolis* strain 4 contained highly conserved areas. In the sequence of the fourth alkane hydroxylase (BAH36166) – EHNFGHH – polar histidine (H) with basic properties was substituted for a nonpolar hydrophobic phenylalanine (F). The same substitutions were only found in corresponding sequences of *R. erythropolis* SK121 (ZP04385381) and *R. erythropolis* PR4 (YP002768905).




Figure 1 Alignment of the amino acid sequences corresponding to the *Rhodococcus erythropolis* strain 4. One conserved His boxes (Hist-3) and the additional HYG motif are boxed and shaded gray. Amino acid residues, conserved in all alkane hydroxylases, are in black colour. The degree of conservation at each position was created using Clustal X.




Figure 2 Phylogenetic tree based on amino acid sequence alignment for *alkB* genes from *R. erythropolis* strain 4 (in bold). Scale bar, 0.01 substitutions per amino acid site. Numerals indicate the statistical reliability of the branching order as determined by bootstrap analysis of 100 alternative trees. Values exceeding 70% were considered significant.

Discussion

AlkB genes in the genomes of Gram-positive and Gram-negative alkane-degrading bacteria are usually present as several individual copies (van Beilen et al. 2003). In particular, *R. erythropolis* NRRL B-16531 and *R. erythropolis* Q15 possess four *alkB* homologues and suggests these bacteria tend to have several *alkB*-genes encoding for alkane hydroxylase (Whyte et al. 2002). In the genome of the *R. erythropolis* strain 4, 4 nucleotide sequences for (oxygenase group) alkane hydroxylases were identified. A 5th *alkB* gene has also been identified which encoded for rubredoxin reductase, but did not cluster the other 4 and will be discussed in later articles.

The alkane hydroxylase amino acid sequence homologies between *R. erythropolis* strain 4, *R. erythropolis* SK121 and *R. erythropolis* PR4 are remarkable for the following reasons: strain SK121 (Hamamura et al. 2008) was isolated from oil contaminated soil and tends to utilise aromatic hydrocarbons. Strain PR4 was isolated at a depth of 1 km from the Pacific Ocean and is unable to utilise arenes, but does use *n*-alkanes with chain length of C₈-C₂₀ as the sole energy source (Sekine et al. 2006). The *R. erythropolis* strain 4 was isolated from the inner part of bitumen mound, located on the bottom of Lake Baikal and tends to utilise *n*-alkanes with a broader chain length (C₁₂-C₂₉). This adaptation could be explained by



Figure 3 Amino acid sequence alignment of the four new alkane hydroxylases from *R. erythropolis* strain 4 from Lake Baikal with another strain already in the literature (*R. erythropolis* PR4). Conserved regions (*), synonymous substitutions (:), significant substitutions (-) and unspecified substitutions (L).

the composition of the bitumen mound 8, which included *n*-alkanes with chain lengths of C₂₂-C₃₄ (Likhoshvay et al. 2013). However, as a final product of alkanes biodegradation serve fatty acids with chain length of C₁₆-C₁₈. These substances could further be degraded during phospholipid synthesis (Alvarez 2010).

Homologue sequence analysis (NCBI) of the 4 amino acid sequences from *R. erythropolis* strain 4 revealed that the 4 alkane hydroxylases were highly divergent, however each enzyme was similar to the corresponding homologue from *Rhodococcus*. The absence of other bacterial genera in the analyses suggested this was an enzyme system specific to rhodococci, based on the differences in alkane hydroxylases.

All alkane-degrading bacteria have alkane hydroxylases containing the following three sequences: (numbering from *Pseudomonas putida* GPo1): H₁₃₈E[L/M]xHK₁₄₃, E₁₆₇HxxGHH₁₇₃ and L₃₀₉QRH[S/A]DHHA₃₁₇. According to van Beilen et al. (2005), a histidine in the second and third sequences may affect enzyme activity. Furthermore, the histidine residues in these conserved sequences bind two atoms of Fe(II) in the alkane hydroxylase (Whyte et al. 1999; van Beilen et al. 2005). The longest sequence, L₃₀₉QRH[S/A]DHHA₃₁₇, was present in the alkane hydroxylase sequences of most hydrocarbon-oxidising bacteria, including *R. opacus* B4, which was initially isolated from oil contaminated soil and metabolized a wide range of arenes and aliphatics. The genes coding for these enzymes were located in (at least) six replicons: a large linear chromosome of 7,913,450 bp, two linear plasmids - pROB01 (558,192 bp) and pROB02 (244,997 bp), and three circular ones - pKNR (111,160 bp), pKNR01 (4,367 bp) pKNR02 (2,773 bp). Originally isolated at a depth of 1 km in Pacific Ocean, *R. erythropolis* PR4 had a circular chromosome of 6,516,310 bp, a separate linear plasmid - pREL1: 271,577 bp and two circular plasmids - pREC1 - 104,014 bp and pREC2 - 3,637 bp. The first two code for most of the genes responsible for alkane metabolism. Obviously, plasmid *alk* genes could be transferred between bacteria by horizontal gene transfer (Turova et al. 2008). Hence, bacteria of the *Geobacillus* genus could obtain *alk* genes from *Rhodococcus*.

The least ones could be found everywhere and in different climatic zones and they have enormous biodegradation potential to utilise widest range of organic substrates. The structure of *alk* genes apparently has an adaptive character and encodes alkane hydroxylase. This might be necessary for *R. erythropolis* strain 4 to degrade of heavy *n*-alkanes, which are present in bitumen mound 8 at low temperature (3.5°C) and high pressure (90 atm).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AL wrote most of the text and lot of analytical work in the lab and the literature study, MG edited the manuscript and supervised the entire study, AL corrected the manuscript and took care about revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Genotypic (Bangalore, India) and Dr. Roja C. Mugasimangalam personally for assistance in sequencing on the Illumina GAllx; Dr. T.A. Scherbakova for consulting assistance and Dr. Y.P. Galachyants for assembling the contigs. The work was supported by grants from Integration Project SB RAS 82 and Programme of RAS Presidium 23.8.

Received: 13 February 2014 Accepted: 9 October 2014

Published: 21 October 2014

References

- Alvarez HM (2010) *Biology of Rhodococcus*. Springer, London, New York
- Beard TM, Page MI (1998) Enantioselective biotransformations using rhodococci. *Antonie Leeuwenhoek* 74:99–106
- Hamamura N, Fukui M, Ward DM, Inskeep WP (2008) Assessing soil microbial populations responding to crude-oil amendment at different temperatures using phylogenetic, functional gene (*alkB*) and physiological analyses. *Environ Sci Technol* 42(20):7580–7586
- Khlystov OM, Zemskaya TI, Sitnikova TYA, Mekhanikova IV, Kaigorodova IA, Gorshkov AG, Timoshkin OA, Shubenkova OV, Chernitsyna SM, Lomakina AV, Likhoshvai AV, Sagalevich AM, Moskvina VI, Peresyplkin VI, Belyaev NA, Slipenchuk MV, Tulokhonov AK, Grachev MA (2009) Bottom bituminous constructions and biota inhabiting them according to investigation of Lake Baikal with the Mir submersible. *Dokl Earth Sci* 429(8):1333–1336
- Kloos K, Munch JC, Schlöter M (2006) A new method for the detection of alkane-monooxygenase homologous genes (*alkB*) in soils based on PCR-hybridization. *J Microbiol Methods* 66:486–496
- Larkin MJ, De Mot R, Kulakov LA, Nagy I (1998) Applied aspects of *Rhodococcus* genetics. *Antonie Leeuwenhoek* 74:133–153
- Likhoshvay A, Khanaeva T, Gorshkov A, Zemskaya T, Grachev M (2013) Do oil-degrading Rhodococci contribute to the genesis of deep water bitumen mounds in Lake Baikal? *Geomicrobiol* 30(3):209–213
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning. A Laboratory Manual* Cold. Spring Harbor Laboratory Press, New York
- Sekine M, Tanikawa S, Omata S, Saito M, Fujisawa T, Tsukatani N, Tajima T, Sekigawa T, Kosugi H, Matsuo Y, Nishiko R, Imamura K, Ito M, Narita H, Tago S, Fujita N, Harayama S (2006) Sequence analysis of three plasmids harboured in *Rhodococcus erythropolis* strain PR4. *Environ Microbiol* 8(2):334–346
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599
- Turova TP, Nazina TN, Mikhailova EM, Rodionova TA, Ekimov AN, Mashukova AV, Poltarauk AB (2008) *AlkB* Homologues in thermophilic bacteria of the genus *Geobacillus*. *Mol Biol* 42(2):247–257
- van Beilen JB, Funhoff EG (2007) Alkane hydroxylase homologues involved in microbial alkane degradation. *Appl Microbiol Biotechnol* 1:13–21
- van Beilen JB, Panke S, Lucchini S, Franchini AG, Röthlisberger M, Witholt B (2001) Analysis of *Pseudomonas putida* alkane degradation gene clusters and flanking insertion sequences: Evolution and regulation of the *alk*-genes. *Microbiol* 147:1621–1630
- van Beilen JB, Li Z, Duetz WA, Smits THM, Witholt B (2003) Diversity of alkane hydroxylase systems in the environment. *Oil Gas Sci Technol* 58(4):427–440
- van Beilen JB, Smits THM, Roos FF, Brunner T, Balada SB, Röthlisberger M, Witholt B (2005) Identification of an amino acid position that determines the substrate range of integral membrane alkane hydroxylases. *J Bacteriol* 187(1):85–91
- Warhurst AW, Fewson CA (1994) Biotransformations catalyzed by the genus *Rhodococcus*. *Crit Rev Biotechnol* 14:29–73
- Whyte LG, Hawari J, Zhou E, Bourbonnière L, Inniss WE, Greer CW (1998) Biodegradation of variable-chain-length alkanes at low temperatures by a psychrotrophic *Rhodococcus* sp. *Appl Environ Microbiol* 64:2578–2584
- Whyte LG, Slagman SJ, Pietrantonio F, Bourbonnière L, Koval SF, Lawrence JR, Inniss WE, Greer CW (1999) Physiological adaptations involved in alkane

assimilation at a low temperature by *Rhodococcus* sp. strain Q15. *Appl Environ Microbiol* 65(7):2961–2968

Whyte LG, Smits THM, Labbe' D, Witholt B, Greer CW, van Beilen JB (2002) Gene cloning and characterization of multiple alkane hydroxylase systems in *Rhodococcus* strains Q15 and NRRL B-16531. *Appl Environ Microbiol* 68:5933–5942

doi:10.1186/2193-1801-3-621

Cite this article as: Likhoshvay *et al.*: The complete *alk* sequences of *Rhodococcus erythropolis* from Lake Baikal. *SpringerPlus* 2014 **3**:621.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
