

# Antitumor pharmacotherapy of colorectal cancer in kidney transplant recipients

Yuanyuan Fu, Chengheng Liao, Kai Cui, Xiao Liu and Wentong Fang

*Ther Adv Med Oncol*

2019, Vol. 11: 1–20

DOI: 10.1177/  
1758835919876196

© The Author(s), 2019.  
Article reuse guidelines:  
sagepub.com/journals-  
permissions

**Abstract:** Renal transplantation has become the sole most preferred therapy modality for end-stage renal disease patients. The growing tendency for renal transplants, and prolonged survival of renal recipients, have resulted in a certain number of post-transplant colorectal cancer patients. Antitumor pharmacotherapy in these patients is a dilemma. Substantial impediments such as carcinogenesis of immunosuppressive drugs (ISDs), drug interaction between ISDs and anticancer drugs, and toxicity of anticancer drugs exist. However, experience of antitumor pharmacotherapy in these patients is limited, and the potential risks and benefits have not been reviewed systematically. This review evaluates the potential impediments, summarizes current experience, and provides potential antitumor strategies, including adjuvant, palliative, and subsequent regimens. Moreover, special pharmaceutical care, such as ISDs therapeutic drug monitoring, metabolic enzymes genotype, and drug interaction, are also highlighted.

**Keywords:** anticancer drug, colorectal cancer, immunosuppressive agent, renal transplantation

Received: 11 September 2018; revised manuscript accepted: 19 August 2019.

## Introduction

Renal transplant, commonly performed for end-stage renal disease (ESRD), is an alternative to dialysis. After renal transplant, immunosuppressive drugs (ISDs) are used continuously to prevent allograft rejection. Improvements in ISDs have led to significantly enhanced graft survival. As a result, kidney transplant recipients (KTRs) have greater accumulation and longer exposure to ISDs than ever before. Concomitantly, the incidence of severe side effects, such as malignancies, has increased, which has become a major obstacle for long term survival in KTRs.<sup>1</sup>

Colorectal cancer (CRC) is the third most common cancer worldwide, and the second highest cause of cancer mortality. It has been reported that the risk of CRC is higher in KTRs than in the nontransplant population.<sup>2</sup> Antitumor therapy, especially antitumor pharmacotherapy, can significantly improve the prognosis of CRC. However, substantial impediments such as carcinogenesis of ISDs, drug interaction between ISDs and anticancer drugs, and toxicity of anticancer drugs exist. In

this review, we systematically evaluate the potential risks of antitumor drugs in order to provided optimal antitumor pharmacotherapy in KTRs with CRC.

## Epidemiology of CRC after renal transplant

Malignancy is one of the complications most often encountered in the later period post transplantation. Well-recognized types of malignancies are skin, kidney, lung, bladder, liver, lymphoma, Kaposi's sarcoma, and carcinoma of the vulva perineum.<sup>3</sup> It is generally recognized that renal transplant has also been related to increased risk of CRC,<sup>4,5</sup> although the risk increasing of rectal cancer is much more controversial.<sup>6</sup> Previously reported incidence of CRC in KTRs ranges dramatically from 0.02% to 3.62%,<sup>3, 7–14, 2, 6, 15–38</sup> (Table 1). This variation might be due to differences in geographical location, duration and degree of immunosuppression, age of renal transplant, complication, etc. Anyway, the growing tendency for renal transplant and prolonged survival of KTRs have resulted in a certain number of CRC patients after renal transplant.

Correspondence to:  
**Wentong Fang**  
Department of Pharmacy,  
First Affiliated Hospital  
of Nanjing Medical  
University, No 300  
Guangzhou Road, Nanjing,  
Jiangsu Province, 210029,  
China  
[fangwentong@jnsph.org.cn](mailto:fangwentong@jnsph.org.cn)

**Yuanyuan Fu**  
Department of Pharmacy,  
First Affiliated Hospital  
of Nanjing Medical  
University, China

**Chengheng Liao**  
Lineberger Comprehensive  
Cancer Center, University  
of North Carolina School  
of Medicine, Chapel Hill,  
NC, USA

**Kai Cui**  
Department of Pharmacy,  
Liaocheng Infectious  
Disease Hospital,  
Liaocheng, Shandong,  
China

**Xiao Liu**  
Department of Pharmacy,  
Qinghai provincial Peoples  
Hospital, Xining, China



**Table 1.** The incidence of colorectal cancer in renal transplant recipients.

Source	No of renal transplant recipients	No of malignancy	Incidence of malignancy (%)	No of CRCs	Incidence of CRCs (%)
Adami <sup>7</sup>	5004	639	12.77	37	0.74
Birkeland <sup>8</sup>	2272	97	4.27	0	0.00
Birkeland <sup>9</sup>	1821	209	11.48	13	0.71
Blohme <sup>10</sup>	934	32	3.43	2	0.21
Cheung <sup>11</sup>	4895	299	6.11	29	0.59
Collett <sup>12</sup>	25,104	4422	17.61	181	0.72
Demir <sup>3</sup>	100	14	14.00	2	2.00
Disney <sup>13</sup>	6641	2449	36.88	38	0.57
Ebisui <sup>14</sup>	309	20	6.47	3	0.97
Ho <sup>15</sup>	1387	—	—	9	0.65
Hoover <sup>16</sup>	6297	63	1.00	1	0.02
Hsiao <sup>17</sup>	642	54	8.42	3	0.47
Ishikawa <sup>18</sup>	1312	35	2.67	3	0.23
Ju <sup>19</sup>	2630	177	6.73	22	0.84
Kasiske <sup>20</sup>	1500	87	5.80	11	0.73
Kasiske <sup>21</sup>	35,765	—	—	182	0.51
Kehinde <sup>22</sup>	492	32	6.50	3	0.61
Kim <sup>23</sup>	2630	—	—	17	0.65
Krynitz <sup>24</sup>	7952	2774	34.88	81	1.02
Kwon <sup>25</sup>	248	—	—	4	1.61
Kyllönen <sup>26</sup>	2090	94	4.50	8	0.38
Kyllönen <sup>27</sup>	2890	230	7.96	13	0.45
Langer <sup>28</sup>	2159	116	5.37	4	0.19
Lj <sup>29</sup>	4716	320	6.76	15	0.32
Papaconstantinou <sup>2</sup>	—	—	—	93	—
Parikshak <sup>30</sup>	638	—	—	1	0.16
Penn <sup>31</sup>	10,667	—	—	386	3.62
Popov <sup>32</sup>	185	15	8.11	1	0.54
Rostami <sup>33</sup>	—	—	—	8	—
Saidi <sup>34</sup>	556	31	5.58	3	0.54
Stewart <sup>6</sup>	62,088	—	—	68	0.11
Vajdic <sup>35</sup>	10,180	1236	12.14	95	0.93
Vilardell <sup>36</sup>	2222	66	2.97	2	0.09
Villeneuve <sup>37</sup>	11,033	778	7.05	51	0.46
Zilinska <sup>38</sup>	1421	85	5.98	11	0.77

CRC, colorectal cancer.

CRC in KTRs has a worse 5-year survival rate than in the general population.<sup>2,23</sup> Papaconstantinou and colleagues researched patients in Israel Penn International Transplant Tumor Registry and found that transplantation patients had a younger mean age at colorectal cancer diagnosis (58 *versus* 70 years;  $p < 0.001$ ), and a worse 5-year survival (overall, 44% *versus* 62%,  $p < 0.001$ ; Dukes A and B, 74% *versus* 90%,  $p < 0.001$ ; Dukes C, 20% *versus* 66%,  $p < 0.001$ ; and Dukes D, 0% *versus* 9%,  $p = 0.08$ ).<sup>2</sup> Kim and colleagues investigated the CRC in KTRs in Korea from 1994 to 2007, and found that recurrence (35.2% *versus* 15.2%,  $p = 0.048$ ) and 5-year survival (40.2% *versus* 67.8%;  $p = 0.044$ ) were worse in the transplant group than in the non-transplant group.<sup>23</sup> This difference was more significant in advanced cancers. The 2-year survival of the transplant group was significantly worse than the nontransplant group with advanced cancer (stages III–IV; 45.7% *versus* 71.6%;  $p = 0.023$ ).

Besides immunosuppression, the decreased survival is also due to inadequate anticancer treatment.<sup>23</sup> Surgical resection and radiotherapy remain the preferred treatment for most early-stage and locally invasive CRC in KTRs. Pharmacotherapy is necessary to prevent tumor relapse and development of metastases and achieve adequate palliation. However, pharmacotherapy of CRC in KTRs is a dilemma, and has significant impediments.

## Immunosuppressive therapy in KTRs with CRC

### *ISD selection in KTRs with CRC*

Induction period of immunosuppressive therapy starts before, or at the time of, kidney transplantation. Hereafter, initial maintenance therapy initiates and a combination of ISDs is recommended. If there is no acute rejection, lowest planned doses of maintenance ISDs are used as long-term maintenance therapy starting 2–4 months after transplantation. The mean onset time of CRC is 10.4 years after transplantation,<sup>39</sup> which is the long-term maintenance period of immunosuppressive therapy. In this period, the combination of ISDs with different mechanisms of action and at reduced doses has recommended by most society- and government-sponsored guidelines. Conventional maintenance regimens includes a calcineurin inhibitor (CNI) and an antiproliferative agent, with or without corticosteroids. This

strategy minimizes morbidity and mortality associated with each class of agent while maximizing overall effectiveness.

Carcinogenesis is an important consideration for choosing of ISD in the maintenance period. Calcineurin controls the nuclear factor of activated T cells (NFAT), an essential transcriptional factor for the activation of T lymphocytes,<sup>40</sup> and is, therefore, a target of cyclosporine A (CsA) and tacrolimus (TAC). The overexpression of calcineurin in human colorectal adenocarcinomas, and the activation of NFAT in human colon cancer cell lines, have been reported.<sup>41</sup> CsA and TAC were shown to reduced cell growth in colony cell lines (HT29).<sup>41</sup> Paradoxically, some research also showed that CsA and TAC increased transforming growth factor-beta, an effector clearly associated with tumor growth.<sup>42,43</sup> The controversy surrounding the anti-CRC activity of CNI is still unresolved, but TAC is associated with decreased acute rejection rates and is generally better tolerated than CsA.<sup>44</sup> Thus, we suggest the administration of TAC rather than CsA.

Azathioprine (AZA), sirolimus (SIR),<sup>45</sup> and mycophenolate mofetil (MMF)<sup>46</sup> are commonly used antiproliferative agents. The use of AZA has been associated with neoplastic development post-transplantation, particularly an increased risk of cutaneous squamous cell carcinomas. The mechanism of action is postulated to be the inhibition of repair splicing and induction of codon misreads via intercalation of DNA.<sup>47</sup> AZA is not preferred because of its inferior ability to prevent acute rejection and its poorer side effect profile than MMF. SIR, an inhibitor of the mammalian target of rapamycin (mTOR), can suppress the growth of tumors.<sup>45</sup> A meta-analysis of 21 randomized trials showed that SIR was associated with a 40% reduction in the risk of malignancy (adjusted HR 0.60, 95% CI 0.39–0.93,  $p = 0.02$ ), but 43% increased risk of death (adjusted HR 1.43, 1.21 to 1.71;  $p < 0.001$ ) compared with other ISD.<sup>46</sup> Elevation of inosine monophosphate dehydrogenase was found in some solid tumors. MMF blocked purine biosynthesis *via* inhibition of inosine monophosphate dehydrogenase.<sup>18,48</sup> Some population studies have suggested that the risk of developing a malignancy is not higher with MMF, and may actually be associated with a decreased risk.<sup>18,48,49</sup> It is still not clear whether SIR- or MMF-containing regimens improve patient outcomes. SIR showed no particular superiority to MMF, but was associated with

increased risk of graft loss when combined with CNI, even when combined with a reduced dose of CNI. Hence, we suggest administration of MMF rather than SIR.

**Kidney Disease: Improving Global Outcomes (KDIGO) guidelines** suggested that, among patients at low immunologic risk and who receive induction therapy, prednisone should be discontinued during the first week after transplantation. This is based on the desire to minimize long-term glucocorticoid exposure, thereby decreasing the risk of adverse effects with this agent. However, many transplant centers continue low-dose glucocorticoid therapy in all patients, regardless of the risk of acute rejection.

#### *Future perspectives and clinical relevance of ISD*

The controversy surrounding the anti-CRC activity of CNI is still unresolved, despite its ability to act both as a tumor suppressor and as an oncogenic activity inducer in CRC. CNI dose may be an important factor in its immunosuppressive and anticancer effects. In a phase I/II trial on 44 patients with advanced non-small cell lung carcinoma, low-dose CsA (1–2 mg/kg per day) was compared with high-dose CsA (3–6 mg/kg per day) in association with etoposide and cisplatin. In this small series, the authors reported a significant increase in survival of patients treated with low-dose CsA, with a 2-year survival of 25% compared with 4% with high-dose CsA. Kaplan–Meier survival curves were significantly different for these two groups by log-rank test ( $p = 0.047$ ).<sup>50</sup> In this regard, more studies are needed to evaluate the effects of different doses of CNIs on CRC and immunology.

NFAT is a family of common cellular transcription factors that is initially identified in T cells. The NFAT family includes five members: NFAT1, NFAT2, NFAT3, NFAT4, and NFAT5. Previous research showed that NFATc2 is overexpressed in CRC tissues compared with normal tissue margins ( $p \leq 0.05$ ), and might be used as a therapeutic target.<sup>51</sup> The study of Vaeth and colleagues also showed that selective NFAT targeting (NFAT1, NFAT2, or a combination of both) in T cells might ameliorates graft-versus-host disease while maintaining antitumor activity.<sup>40</sup> Hence, further investigations should identify specific targets for calcineurin or NFAT to anti-tumor. More importantly, selective CNIs or natural compounds

against targets could be useful clinically for treating CRC in KTRs.

#### **Pharmacotherapy of CRC**

For CRC, the available agents include oxaliplatin, irinotecan, parenteral and oral fluoropyrimidines, anti-angiogenesis drugs, anti-epidermal growth factor receptor (EGFR) monoclonal antibody (MoAb), and drugs targeting the programmed cell death-1 (PD-1)/programmed death-ligand 1 (PD-L1).<sup>52</sup>

For patients with indications for adjuvant chemotherapy, a combination chemotherapy of 5-fluorouracil (5-FU)/leucovorin (LV)/oxaliplatin (FOLFOX) or capecitabine/oxaliplatin (CapeOx) is the preferred option, while fluoropyrimidine-based regimen (5-FU/LV or capecitabine alone) is recommended for patients who cannot tolerate combined chemotherapy.<sup>53</sup> 5-FU/LV/irinotecan (FOLFIRI) and 5-FU/LV/oxaliplatin/irinotecan (FOLFOXIRI), as well as the regimens mentioned above, are recommended as palliative regimens. Bevacizumab is recommended to add to chemotherapy in palliative therapy. Cetuximab or panitumumab is superior to bevacizumab for patients with a wild-type KRAS gene.<sup>54,55</sup> Some new drugs, such as aflibercept, regorafenib, trifluridine-tipiracil, S-1, tegafur-uracil, nivolumab, and pembrolizumab, have been suggested in subsequent therapy.<sup>56</sup> For KTRs with CRC, anticancer regimens should be optimized as far as possible to avoid interfering with immunosuppressive therapy.

#### **Possible interaction between ISD and anticancer drugs**

Whenever patients take more than one medication, they are at risk of a drug interaction. For post-transplantation CRC patients, potential interaction between ISDs and anticancer drugs must be checked before it occurs. Drug interactions occur in two different ways: pharmacokinetic interaction and pharmacodynamic interaction.

#### *Pharmacokinetic interaction*

Pharmacokinetic interaction may occur if one drug affects another drug's absorption, distribution, metabolism, or excretion. There is no research on pharmacokinetic interactions between ISDs and antitumor drugs, although some

interaction in absorption, distribution, metabolism, and excretion could be predicted.

*Interaction in drug absorption.* ISDs are absorbed in the gastrointestinal tract, and bioavailability varies from 14% to 94% (Table 2). Many factors can affect the absorption of ISDs; for example, colitis can affect the secretion and excretion of bile, which disturbs the enterohepatic circulation and bioavailability of MMF. Anticancer drugs have gastrointestinal toxicity, leading to diarrhea, vomiting, and constipation, which disturbs gastrointestinal function and subsequently ISDs absorption.

*Diarrhea.* Diarrhea is most commonly described with chemotherapeutic drugs such as fluoropyrimidines and irinotecan. Irinotecan is associated with both early- and late-onset diarrhea. Early-onset diarrhea occurs within several hours of drug infusion and has an incidence of 45–50%. It is cholinergically mediated and usually well controlled by subcutaneous or intravenous atropine. Late-onset diarrhea induced by irinotecan is not cholinergically mediated, and is unpredictable. The median time to onset is approximately 6 days, with a 350 mg/m<sup>2</sup> every 3-week schedule, and 11 days with a weekly schedule (125 mg/m<sup>2</sup>).<sup>57</sup> The pathophysiology of late diarrhea appears to be multifactorial, with some association with the polymorphism of UGT1A1.<sup>58</sup> In some studies, homozygotes for the UGT1A1\*28 (and, to a lesser degree, heterozygotes) have had significantly higher rates of diarrhea with irinotecan. The UGT1A1 genotype should be detected before the initiation of irinotecan-containing regimens such as FOLFIRI and FOLFOXIRI. It is advised that the starting dose of irinotecan should be lowered in patients known to be homozygous for UGT1A1\*28.

Diarrhea can occur in all schedules with fluoropyrimidines. A meta-analysis showed that more frequent grade 3/4 diarrhea could be found in capecitabine-based groups (507/2658) than in 5-FU-based groups (333/2624).<sup>59</sup> The pathophysiology of diarrhea appears to be multifactorial, with some association with the polymorphisms of dihydropyrimidine dehydrogenase (DPD) and thymidylate synthetase (TYMS).<sup>60,61</sup> Polymorphisms of DPD and TYMs lead to deficient enzyme activity and inadequate degradation of fluoropyrimidines, and, thus, subsequently increase the risk of severe toxicity such as diarrhea. Testing of DPD and TYMS polymorphism for fluoropyrimidine-based

chemotherapy is a controversial area nowadays,<sup>60</sup> and has not been widely adopted. We advise detecting DPD and TYMS genotype in KTRs to identify high-risk patients, and making an early decision to reduce dose or select an alternative treatment regimen.

In contrast to fluoropyrimidines and irinotecan, anti-EGFR MoAbs related diarrhea is generally not pervasive or severe.<sup>62,63</sup> But adding anti-EGFR MoAbs to standard chemotherapy increased the rates of diarrhea. In a meta-analysis of 18 randomized controlled clinical trials, the incidence of grade 3–4 diarrhea was 18% (95% CI 14–23%) in the anti-EGFR MoAbs arm and 11% (95% CI 8–14%) in the control arm.<sup>64</sup> Using a fixed-effects model, the overall result showed that anti-EGFR regimens are associated with a significantly higher risk of severe diarrhea (RR of 1.66, 95% CI 1.52–1.80).<sup>64</sup> Other studies also showed similar results.<sup>46,65</sup>

Considering the cornerstone status of fluoropyrimidines in the treatment of CRC, fluoropyrimidine monotherapy or combination chemotherapy still applies to KTRs with CRC, although they have a high risk of diarrhea. DPD and TYMS genotype test might be helpful in these patients. Regimens containing fluoropyrimidines and irinotecan (FOLFIRI and FOLFOXIRI) are not preferred, but can be used in oxaliplatin-refractory patients. UGT1A1 genotype should be detected before the initiation of irinotecan-based regimens. Triple combinations of fluoropyrimidines, irinotecan and anti-EGFR MoAbs (cetuximab/panitumumab + FOLFIRI) are less recommended due to their synergistic effect of diarrhea. For patients needing intensive treatment, the triple combination of fluoropyrimidines, oxaliplatin, and anti-EGFR MoAbs (cetuximab/panitumumab + FOLFOX or cetuximab/panitumumab + CapeOx) might be a choice, but also increase the risk of other adverse drug reactions (ADRs) besides diarrhea.

Anticancer regimens optimization can minimize the risk of diarrhea, but cannot avoid diarrhea completely. It has been confirmed that the absorption of MMF<sup>66,67</sup> and CsA<sup>68</sup> is little affected by diarrhea, and needs no further dose adjustments. However, severe diarrhea (more than three loose stools daily) substantially increased with exposure to TAC, which is also reflected as an elevation of the trough level. Therefore, dose adjustments of TAC in patients with severe diarrhea must be



**Table 2.** The absorption and distribution of immunosuppressive agents and anticancer drugs.

Drugs	Bioavailability	Distribution	Plasma protein binding
Methylprednisolone (tablet)	82%	Widely distributed in the organization	40–90% (mainly albumin)
Prednison (tablet)	Almost 100%	Liver, plasma, cerebrospinal fluid, hydrothorax, ascites	Corticosteroid-binding globulin
Cyclosporine A (soft capsules)	20–30%	Fat, liver, adrenal gland and pancreas	About 90% (mainly lipoprotein)
Tacrolimus (capsules)	20–25%	Lungs, spleen, heart, kidneys and pancreas	About 99% (mainly albumin and alpha 1- acidic glycoprotein)
Sirolimus (tablet)	about 27%	Widely distributed in the blood	About 99% (mainly albumin, alpha 1- acidic glycoprotein and lipoprotein)
Sirolimus (oral solution)	about 14%	Widely distributed in the blood	About 99% (mainly albumin, alpha 1- acidic glycoprotein and lipoprotein)
Mycophenolate Mofetil (tablet, capsules)	94% (enterohepatic circulation)	Not reported	About 97% (mainly albumin)
5-FU(injection)	—	Intestinal mucosa, bone marrow, liver, cerebrospinal fluid, and brain tissue.	Not reported
Capecitabine (tablet)	Not reported	Not reported	Less than 60% (about 35% albumin)
Oxaliplatin (injection)	—	85% is rapidly distributed into tissues	More than 75% (about 95% on the 5th day after injection) (mainly albumin and gamma-globulins)
Irinotecan (injection)	—	Not reported	Irinotecan: 30–68%; SN-38: 95%;
Bevacizumab (injection)	—	Not reported	Not reported
Ramucirumab (injection)	—	Not reported	Not reported
Aflibercept (injection)	—	Not reported	Not reported
Regorafenib (tablet)	Not reported	Not reported	About 99.5%;
Cetuximab (injection)	—	Not reported	Not reported
Panitumumab (injection)	—	Not reported	Not reported
Nivolumab (injection)	—	Not reported	Not reported
Pembrolizumab (injection)	—	Not reported	Not reported

monitored carefully, especially when doses of MMF are also reduced.<sup>68</sup> Various explanations for the effect of severe diarrhea upon exposure to TAC have been proposed, but the most important one is enhanced absorption as a consequence of a damaged intestinal barrier and reduced intestinal metabolism.<sup>69</sup> It is generally considered that mild diarrhea might be an effect of exposure to

TAC, but probably to a lesser extent.<sup>69</sup> However, in the study of van Boekel and colleagues,<sup>69</sup> mild diarrhea did not affect the exposure and trough level of TAC, and thus showed no evidence for the presence of hidden TAC overexposure in patients with mild diarrhea while on treatment with TAC and MMF.

*Emesis and constipation.* Trifluridine/tipiracil has moderate to high emetic risk, which means more than 30% frequency of emesis. Oxaliplatin and irinotecan have moderate emetic risk, which means a 30–90% frequency of emesis. The emetic risk of other anticancer drugs used in CRC is low or minimal. Preventive antiemetic therapies were used widely before chemotherapy and can prevent most of the emesis. Additional antiemetic therapy will be given when patients have breakthrough nausea. Dosing interval should be adjusted in order to minimize the risk of vomiting with ISDs. The T<sub>max</sub> values of TAC, CsA, SIR, and MMF are 1–3 h, 1.5–2.0 h, 1.0 h, and 0.9–1.8 h, respectively. Hence, ISDs should be taken orally 1–3 h before the beginning of an anticancer regimen.

The effects of constipation on ISDs exposure have not been reported. Anticancer drugs rarely cause constipation directly. Antiemetic therapies, such as 5-hydroxytryptamine receptor antagonists, often induce constipation that slows intestinal transit time. The initial management of constipation includes patient education, dietary changes, bulk-forming laxatives, or the use of nonbulk-forming laxatives or enemas. Considering multiple concurrent drugs, it is preferred for KTRs with CRC to increase fluid and fiber intake to prevent constipation. Laxatives are used only for existing constipation, because laxatives affect CNI absorption.<sup>70</sup>

Attention should be paid to the interaction between ISD and proton pump inhibitors (PPIs) as it is widely implicated in chemotherapy-induced gastrointestinal dysfunction. It has been confirmed that PPIs reduce absorption of MMF by elevating gastric pH value and decreasing dissolution of MMF.<sup>71,72</sup> Rupperecht and colleagues demonstrated that the maximum concentration of a single dose of MMF was reduced by 57% in healthy volunteers, and exposure was reduced by 27% when given 1 h after a single dose of pantoprazole 40 mg.<sup>73</sup> A number of studies designed to evaluate the impact of MPA pharmacokinetics in transplant recipients on various PPI therapies have shown similar effects on MMF exposure, but failed to demonstrate a significant impact on rejection rates, graft function, or graft survival.<sup>74</sup> However, physicians should pay more attention to monitoring MMF levels in the presence of PPIs.

The presence of clinically relevant pharmacokinetic drug interaction between PPIs and TAC

remains a matter of controversy.<sup>75</sup> *In vivo* studies using human liver microsomes have shown that omeprazole inhibits CYP3A4-mediated metabolism of TAC competitively.<sup>75</sup> In contrast, Pascual and colleagues estimated the potential interaction between omeprazole and TAC in renal transplant recipients and concluded an absence of important drug interaction.<sup>76</sup> In these conditions, rabeprazole or H2RA were proposed as a safer treatment option than omeprazole in KTRs receiving TAC.

*Interaction in drug distribution.* When a drug is displaced from its plasma binding protein, increased unbound drug concentrations theoretically cause an increase in drug effect, with potentially toxic results.<sup>77</sup> However, some scholars have presented theoretical arguments from a few cases where protein-binding changes were clinically significant. Benet and colleagues considered that changes in plasma protein binding have little clinical relevance.<sup>77</sup> We also summarize the distribution and plasma protein binding of ISD (Table 2) in order to minimize the potential risk of interaction in drug distribution. Methylprednisolone, TAC, SIR, and MMF bind mainly to albumin, and prednisone binds mainly to corticosteroid transporters, while CsA binds mainly to lipoprotein. Plasma protein binding rates of ISDs vary from 40% to 99%. TAC, MMF, and oxaliplatin bind mainly to albumin, but the specific binding site and binding pattern to albumin are still unknown. On the other hand, it is still unclear what proportion of plasma protein is bound to each drug. So it is uncertain whether oxaliplatin competes for protein binding with TAC and MMF. Therapeutic drug monitoring of ISD cannot predict protein binding competition because it represents the total plasma concentration not the free concentration. But we also recommend therapeutic drug monitoring of ISDs because no other indicator can be used in this condition.

*Interaction in drug metabolism.* The metabolism of ISD and anticancer drugs are summarized in Table 3. In humans, three carboxylesterases (CES) have been identified: human liver CES (CES1), human intestinal CES (CES2), and human brain CES (CES3).<sup>78</sup> MMF is an inactive ester prodrug, and undergoes hydrolysis to form an active drug, mycophenolic acid. Hydrolysis occurs in the intestine, plasma, and liver. Liver hydrolysis by CES1 has been demonstrated to be the most efficient pathway.<sup>79</sup> Capecitabine, a prodrug of 5-fluorouracil, is first metabolized to 5'-deoxy-5-fluorocytidine (5'-DFCR), mainly by CES2.<sup>80</sup>

Hence, the combination of MMF and capecitabine rarely causes competition of CES.

Irinotecan is a prodrug whose hydrolysis results in the formation of the active metabolite, 7-ethyl-10-hydroxy camptothecin (SN-38).<sup>81</sup> Both CES1 and CES2 appear to contribute to the hydrolysis of irinotecan to SN-38.<sup>82</sup> Drug activation in the intestine and kidney are likely major contributors to SN-38 production *in vivo*. Studies have also shown that CES2 is commonly expressed in tumor tissue and is correlated with activation of irinotecan.<sup>83</sup> As MMF is hydrolyzed mainly by liver CES1, competition with irinotecan for CES seems unlikely.

CYP3A4 activates regorafenib and prednisone, but inactivates CsA, TAC, and SIR. Competition of CYP3A4 between regorafenib and ISD has not been reported, but can be anticipated. Competition of CYP3A4 might increase the plasma concentration of regorafenib and decreased the plasma concentrations of the active metabolites M-2 and M-5, and may lead to decreased efficacy and toxicity. The plasma concentration and immunosuppressive effect of CsA, TAC, and SIR might be increased, while that of prednisone might be decreased when competition of CYP3A4 occurs. As CNIs such as CsA or TAC are necessary for maintenance immunosuppressive therapy, the use of regorafenib should be avoided as far as possible.

The active metabolites of irinotecan (SN-38) and MMF (mycophenolic acid) are inactivated by uridine diphospho-glucuronosyltransferase 1A (UGT1A). The combination of irinotecan and MMF has the potential for competition with UGT1A, although this has not yet been confirmed. The potential competition might lead to the accumulation of the active metabolites SN-38, which significantly enhance the possibility of fatal late diarrhea.<sup>58</sup>

Regorafenib has proved to be a UGT1A inhibitor.<sup>84</sup> Inhibition of UGT1A by regorafenib or competition of UGT1A by irinotecan might result in the accumulation of mycophenolic acid, and thus reinforce the immunosuppressive effect. Considering the wide use of MMF in immunosuppressive therapy, the use of irinotecan and regorafenib is restricted for these patients.

*Interaction in drug excretion.* The excretion of ISD and anticancer drugs are summarized in

Table 3. Capecitabine, oxaliplatin, prednisone, and MMF are excreted in the urine without the participation of a carrier. So the probability of excretive interaction is small.

#### *Pharmacodynamic interaction*

Pharmacodynamic interaction occurs when two drugs given together act at the same or similar receptor site and lead to a greater (additive or synergistic) effect or a decreased (antagonist) effect. ISDs, such as CsA, TAC, and SIR, block the critical events that lead to the activation of alloreactive T cells and the subsequent amplification of the signals involved in T cell proliferation. MMF inhibits the *de novo* synthesis of purines, which are necessary for the proliferation and function of T and B lymphocytes. PD-1 is a cell surface receptor that belongs to the immunoglobulin superfamily and is expressed on T cells and pro-B cells.<sup>85</sup> PD-1 binds two ligands, PD-L1, and PD-L2, and guards against autoimmunity through promoting apoptosis (programmed cell death) of antigen-specific T-cells in lymph nodes and reducing apoptosis in regulatory T cells. PD-1/PD-L1 inhibitors attack tumors by activating the immune system,<sup>86</sup> which potentially antagonize the immunosuppressive effect of ISDs. Though there are no published data on the potential risk, we do not suggest the application of PD-1/ PD-L1 inhibitors in KTRs.

Though acting at different receptor sites, ISDs and cytotoxic chemotherapy simultaneously enhance the systemic level of immunosuppression. Molecular targeted drugs have a very low risk of infection, and rarely increase the infection risk of chemotherapy. Nephrotoxicity, which is the potential ADR of CsA and TAC,<sup>87</sup> is also the potential ADR of cytotoxic chemotherapy and molecular targeting drugs. Hence, it is necessary to choose regimens with less risk of infection and nephrotoxicity.

#### *Risk of infection*

Infection is one of the major causes of death following renal transplant. The risk of infection in KTRs is determined by the synergy between two factors: the epidemiologic exposure of the individual and the 'net state of immunosuppression', which is a conceptual measure of all of the factors that contribute to the individual's susceptibility (or resistance) to infection.<sup>88,89</sup> Infections are most likely to occur between the 1st and 3rd month



**Table 3.** The metabolism and excretion of immunosuppressive agents and anticancer drugs.

Drugs	Metabolism	Clearance	Enzyme/transporter	
			Substrate	Inhibition
Prednison	CYP3A4, CYP3A5 (possibly)	urine	—	—
Cyclosporine A	CYP3A4	feces	P-gp	P-gp
Tacrolimus	CYP3A4, CYP3A5	feces	P-gp	P-gp (supportive data are limited)
Sirolimus	CYP3A4	feces	P-gp	—
Mycophenolate Mofetil	CES, UGT	urine	—	—
5-FU	DPYP, DPYS, UPB1;	Feces (~80%); Urine (~20%)	—	—
Capecitabine	CES, DPYP, DPYS, UPB1;	Urine (main)	—	—
Oxaliplatin	Nonenzymatic conversion	Urine (main)	—	—
Irinotecan	BCHE, UGT1A	Feces and urine	—	—
Trifluridine-tipiracil	No-CYP conversion	Trifluridine: urine(main); Tipiracil: feces (main)	—	—
Bevacizumab	No-CYP conversion	—	—	—
Ramucirumab	No-CYP conversion	—	—	—
Aflibercept	No-CYP conversion	—	—	—
Regorafenib	CYP3A4, UGT1A9	Feces (~70%); Urine (~20%)	—	—
Cetuximab	No-CYP conversion	—	—	—
Panitumumab	No-CYP conversion	—	—	—
Nivolumab	No-CYP conversion	—	—	—
Pembrolizumab	No-CYP conversion	—	—	—

5-FU, 5-fluorouracil; CES, carboxylesterases; CYP, cytochrome P450; GMPS, Guanosine monophosphate synthetase; HGPRT, Hypoxanthine-guanine-phosphoribosyl-transferase; IMPD, Inosine monophosphate dehydrogenase; P-gp, P-glycoprotein; TPMT, thiopurine S-methyltransferase; UGT, UDP-glucuronosyltransferase; XO, Xanthine oxidase.

following transplant, since immune suppression is at its maximum due to induction therapy.<sup>90</sup> The risk of infection is relatively low due to stable and reduced levels of immunosuppression after 3 months of transplant. Cytotoxic chemotherapy adds to the risk of infection by suppressing the production of neutrophils, and by cytotoxic effects on the cells that line the alimentary tract.

The chemotherapy regimen is one of the primary determinants of the risk of neutropenia, and some regimens are more myelotoxic than others. Febrile neutropenia (FN) occurred in 17.6% (190/1078) of patients receiving FOLFIRI treatment, 13.4% (530/3957) of FOLFOX treatment, and 11.6%

(165/1419) of 5-FU treatment.<sup>91</sup> CapeOx (0-1%) has been reported to be less myeloablative than FOLFOX and FOLFIRI.<sup>92</sup> As for monotherapy, the FN risk of capecitabine (0-1%) is lower than that of 5-FU CI (10.0-13.4%).<sup>93</sup> Molecular targeting drugs have very low risk of FN, and rarely increase the FN risk of chemotherapy. For metastatic CRC (mCRC) patients treated with FOLFOX and FOLFIRI regimens, the percentages of FN were 13.4% and 17.6%, respectively, while, combined with bevacizumab, the percentages were 11.8% and 13.3%, respectively.<sup>91</sup>

Cytotoxic chemotherapy can adversely affect the developmental integrity of the gastrointestinal

mucosa, which increases the risk of invasive infection due to colonizing bacteria or fungi that translocate across intestinal mucosal surfaces. Mucositis is the principal manifestation of mucosal toxicity related to chemotherapy.<sup>94</sup> Sonis and colleagues retrospectively analyzed mucositis in previous studies, and found the incidence of grade 3–4 oral mucositis in 5-FU/LV (14%, 95% CI 12–15%) is higher than that of FOLFOX (5%, 95% CI 4–7%), FOLFIRI (5%, 95% CI 3–8%), and FOLFOXIRI (6%, 95% CI 2–11%) regimens, while the incidence of gastrointestinal mucositis of FOLFOXIRI (38%, 95% CI 30–47%) is higher than that of FOLFIRI (25%, 95% CI 20–30%), 5-FU/LV (11%, 95% CI 10–12%), and FOLFOX (8%, 95% CI 6–10%) regimens.<sup>94</sup> As for oral *versus* intravenous fluoropyrimidines, there is no difference in the OR (0.64, 95% CI 0.25–1.62) for grade 3–4 mucositis between capecitabine based regimens, and 5-FU based regimens in adjuvant chemotherapy, while the pooled OR favored capecitabine-based regimens in palliative chemotherapy (OR 0.17, 95% CI 0.12–0.24).<sup>95</sup> Mucositis induced by molecular targeting drugs and immunotherapy drugs have not been reported in clinical trials.

Overall, no regimen for CRC is forbidden for KTRs when considering infection. Considering the immunosuppressive status of KTRs, it is wise to prevent or minimize the risk of infection. CapeOx and FOLFOX regimens have relatively low risk of FN and mucositis in the combined chemotherapy, while capecitabine has a relatively low risk in monotherapy. Adding molecular targeting drugs to chemotherapy does not significantly increase the risk of FN and mucositis. Special attention should be paid to infection during anticancer therapy.

#### *Risk of nephrotoxicity*

Nephrotoxicity is a potential ADR of anticancer pharmacotherapy and may result in a variety of functional abnormalities, including glomerular or tubular dysfunction, hypertension and disturbance of the renal endocrine function. In KTRs with CRC, the successful treatment of diseases might be limited by drug-related renal injury. The nephrotoxicity of these drugs are summarized in Table 4.

Nephrotoxicity is rarely encountered with corticosteroids, SIR, and MMF, but is common with CsA and TAC.<sup>87</sup> Acute kidney injury (AKI) of

CsA and TAC manifests mainly as oliguric AKI and primary nonfunction,<sup>87</sup> while chronic nephrotoxicity is manifested by renal insufficiency due to glomerular and vascular disease, abnormalities in tubular function, and an increase in blood pressure.<sup>96</sup> Trials comparing the risk of nephrotoxicity of CsA and TAC have shown contradictory results. Early researches found a similar incidence of early acute renal failure (ARF) and late hypertension, while late renal insufficiency was more prevalent with TAC.<sup>54,97</sup> In contrast, some trials found that TAC was associated with less nephrotoxicity in liver transplants and similar nephrotoxicity in other transplants.<sup>42,98</sup> Hence, it is generally considered that nephrotoxicity of TAC is generally similar to that of CsA.<sup>47</sup>

CNIs, such as CsA or TAC, is necessary for the maintenance immunosuppressive therapy, so it is wise to minimize the nephrotoxicity of anticancer regimens. In patients with mild renal impairment (eGFR 50–70 ml/min), capecitabine treatment can induce diarrhea-related dehydration, and dehydration may cause acute renal failure.<sup>99</sup> Capecitabine-related renal failure rarely occurs in patients with normal renal function.

Vascular endothelial growth factor (VEGF) blockade is associated with proteinuria, which is rarely in the nephrotic range, and even more rarely associated with the nephrotic syndrome.<sup>100</sup> In patients treated with bevacizumab, the overall incidence of mild proteinuria ranges from 21% up to 63%, and incidence of grade 3 or 4 proteinuria is about 2%.<sup>101</sup> Aflibercept plus chemotherapy led to 62.2% of all-grade proteinuria and 7.9% of grade 3 or 4 proteinuria, while aflibercept alone led to 40.7% of all-grade proteinuria and 1.2 % of grade 3 or 4 proteinuria in a phase III trial.<sup>101</sup> Among patients treated with ramucirumab, the risk of proteinuria may be lower. In a meta-analysis of six placebo-controlled randomized trials, the incidence of all-grade proteinuria for ramucirumab *versus* placebo was 9.4% *versus* 3.1%, while the risk of severe (grade 3 or 4) proteinuria was 1.1% *versus* 0.04%.<sup>102</sup> Previous studies have shown contradictory results on the proteinuria risk of regorafenib. The United States package insert lists the rate of all grade proteinuria for single agent regorafenib to be 51–84% compared with 37–61% in the placebo treatment arm. In a phase III trial involving 500 patients receiving regorafenib, the incidence of all-grade proteinuria was 7.0% and grade 3 or 4 proteinuria was 1.0%.<sup>103</sup>

**Table 4.** Nephrotoxicity of anti-cancer agents and ISD.

Drug	Manifestation of nephrotoxicity
Cyclosporin A	AKI mainly manifests as oliguric AKI and primary nonfunction; Chronic nephrotoxicity is manifested by renal insufficiency. Metabolic abnormalities: hyperkalemia, hyperuricemia, metabolic acidosis, hypophosphatemia, hypomagnesemia, and hypercalciuria.
Tacrolimus	Similar with Cyclosporin A
Sirolimus	Rare
Mycophenolate mofetil	Rare
Corticosteroid	Rare
5-FU	Not reported
Capecitabine	Dehydration, acute renal failure (not common)
Trifluridine-tipiracil	Not reported
Irinotecan	Dehydration, acute renal failure (rare)
Oxaliplatin	Renal tubular vacuolization, acute tubular necrosis, renal tubular acidosis (rare)
Bevacizumab	Proteinuria (All grade: 21–63%; Grade 3–4: 2%)
Aflibercept	Proteinuria (Aflibercept+Chemotherapy: All grade: 62.2%; Grade 3–4: 7.9%) Proteinuria (Aflibercept alone: All grade: 40.7%; Grade 3–4: 1.2%)
Ramucirumab	Proteinuria (All grade: 9.4%; Grade 3–4: 1.1%)
Regorafenib	Proteinuria (All grade: 7.0%; Grade 3–4: 1.0%), hypophosphatemia, hypocalcemia, hyponatremia, and hypokalemia
Cetuximab	Hypomagnesemia (All grade: 37%; Grade 3–4: 5.6%)
Panitumumab	Hypomagnesemia (All grade: 27.6%; Grade 3–4: 7.0%), hypocalcemia, hypokalemia, hyponatremia
Nivolumab	AKI (rare)
Pembrolizumab	AKI (rare)

5-FU, 5-fluorouracil; AKI, acute kidney injury; ISD, immunosuppressive drugs.

Bevacizumab is recommended as a first-line antiangiogenic drug in National Comprehensive Cancer Network (NCCN) guidelines, but ramucirumab seems to have the lowest risk of nephrotoxicity. It is hard to decide the best antiangiogenic drugs for CRC in KTRs. Anti-VEGF associated proteinuria is usually an asymptomatic event detected only through laboratory analysis. Hence, all antiangiogenic drugs are applicable for CRC in KTRs, and bevacizumab might be the first-line choice.

Cetuximab and panitumumab are both associated with the progressive development of

hypomagnesemia due to renal magnesium wasting.<sup>95,96</sup> Nephrotoxicity of cetuximab has also been manifested as hypocalcemia, hypokalemia, and hyponatremia.<sup>104–106</sup> In a meta-analysis of 19 clinical reports, cetuximab-based treatment led to 37% of all-grade hypomagnesemia and 5.6% of grade 3 or 4 hypomagnesemia.<sup>104</sup> Panitumumab led to 27.6% of all-grade hypomagnesemia and 7.0% of grade 3 or 4 hypomagnesemia in patients with chemotherapy-refractory mCRC.<sup>95</sup> Hypomagnesemia resolves after treatment is discontinued. Hence, cetuximab and panitumumab are equally applicable for CRC in KTRs.

Clinically significant renal toxicity induced by chemotherapy regimens is rare in CRC patients. Adding molecular targeted drugs to chemotherapy increase the risk of nephrotoxicity, but is feasible. We suggest measuring urine volume, urine protein excretion, and serum creatinine after every cycle of anticancer therapy or every 1 month. Glomerular filtration rate (GFR) should be estimated whenever serum creatinine is measured. We also suggest kidney allograft ultrasound examination as part of the assessment of kidney allograft dysfunction.

### Other considerations in antitumor pharmacotherapy

#### *Fluoropyrimidine-related cardiotoxicity*

Fluoropyrimidine-related cardiotoxicity is an infrequent but potentially lethal ADR. The most common clinical manifestation is chest pain, which can be either nonspecific or anginal and is usually associated with electrocardiographic (ECG) changes.<sup>107</sup> Symptoms occur with or without elevated serum biomarkers of cardiac injury. Chest pain tends to occur most often in the first cycle of administration, but can be delayed. The incidence of cardiotoxicity in capecitabine (3–9%)<sup>108,109</sup> is within the range of that reported with infusional 5-FU (1–19%).<sup>107,110,111</sup> The incidence in CapeOx regimens (12% in one report) may be higher than with capecitabine alone.<sup>108</sup> Cardiotoxicity appears to be completely reversible after termination of FU treatment.<sup>107</sup> The preferred strategy in most cases is to switch to a nonfluoropyrimidine-containing chemotherapy regimen or a different treatment modality.

#### *Hypertension of antiangiogenesis drugs*

All commercially available angiogenesis inhibitors have been implicated in the development of hypertension. Previous studies have shown similar incidence of all-grade hypertension (24% versus 21.3%) and severe hypertension (8.0% versus 9.0%) between bevacizumab and ramucirumab.<sup>102,104</sup> The incidence of severe hypertension may be higher with aflibercept (aflibercept + chemotherapy 19.1% versus chemotherapy 1.5%) than with other VEGF-targeted therapies, but no direct comparisons of aflibercept with other VEGF inhibitors exist.<sup>112</sup> All patients receiving angiogenesis inhibitor should have their blood pressure actively monitored

during treatment, with more frequent measurement in the first few weeks of therapy. Manage blood pressure during therapy, with an ideal goal, where possible, of keeping blood pressure less than 130/80mmHg for most patients, and lower in those with specific pre-existing cardiovascular risk factors, such as diabetes or chronic kidney disease.

#### *Skin toxicity*

Given the fact that EGFR is expressed in skin and adnexal structures, EGFR inhibitors are associated with skin toxicity, mainly acneiform eruption and xerosis, but also paronychia, abnormal scalp, facial hair, and eyelash growth, maculopapular rash, mucositis, and postinflammatory hyperpigmentation.<sup>113,114</sup> PRIDE syndrome comprises the most frequent reactions associated with anti-EGFR reactions (papules, pustules, paronychia, hair growth disorders, pruritus, and skin and mucosal xerosis).<sup>114</sup> In a recent meta-analysis involving 38 studies, any grade skin toxicities were identical in cetuximab and panitumumab (83.6% versus 84.4%;  $p = 0.99$ ). However, cetuximab was associated with fewer G3-4 skin toxicities (15.7 versus 23.2%; OR = 0.62, 95% CI 0.53–0.62;  $p < 0.001$ ).<sup>115</sup> Conversely, cetuximab was associated with slightly more frequent G3-4 acne-like rash (16 versus 13%; OR = 1.24, 95% CI 1.04–1.48;  $p = 0.04$ ) and paronychia (23 versus 18%; OR 1.36, 95% CI 1.1–1.7) but fewer cases of all-grade skin fissures (8.5 versus 12.8%; OR = 0.64, 95% CI 0.44–0.93;  $p = 0.02$ ) and all-grade pruritus (17.4 versus 32%; OR = 0.45, 95% CI 0.35–0.58;  $p < 0.001$ ) than panitumumab.<sup>115</sup> Incidence of skin toxicity of EGFR inhibitor in renal transplant recipients has not been reported. Because of the common use of corticosteroids, the incidence and severity of EGFR-inhibitor-induced acneiform eruption in kidney recipients might be lower in kidney recipients than in non-transplant patients.

The initial pustules are sterile, but secondary infection of the affected skin may occur. Eilers and colleagues reported that 38% (84 of 221) of patients treated with EGFR inhibitor developed infections with bacteria, dermatophytes, or viruses.<sup>116</sup> *Staphylococcus aureus* was the most common pathogen, found in approximately 60% of infected lesions. Secondary infection may result in sudden worsening of cutaneous rashes and an alteration in clinical status.<sup>114</sup> Hence,

preventive measures are needed, such as adequate hydration of dry areas, decrease in sun exposure, avoiding prolonged skin contact with water, irritants, and solvents. Topical agents can reduce reaction severity. Moderate reactions may be treated with oral antibiotics, including tetracycline, doxycycline, and minocycline besides topical agents. Tetracycline is not recommended for kidney recipients due to its nephrotoxicity. Doxycycline and minocycline can be used in these patients, but the dose of minocycline should be reduced in patients with renal injury.

Paronychia and nail lesions occur in 25% of patients treated with panitumumab and 16% of those treated with cetuximab.<sup>113</sup> Avoiding friction and pressure on the nailfold, and application of topical steroids, are commonly used to reduce reaction severity. Paronychia is not considered an infectious process, but secondary infection with *S. aureus* or coagulase-negative, Gram-positive bacteria (nosocomial colonization) can occur.<sup>117</sup> In this condition, topical antimicrobial agents (mupirocin or nystatin ointment) and oral doxycycline might be used.

#### *Neurotoxicity of oxaliplatin*

The major dose-limiting toxicity with oxaliplatin is neurotoxicity. There are two distinct syndromes: acute neurotoxicity and cumulative sensory neuropathy. Acute neurotoxicity develops within 24–72 h after each dose and recurs with each dose.<sup>118</sup> Symptoms consist of striking paresthesias and dysesthesias of the hands, feet, and perioral region, jaw tightness, and unusual pharyngo-laryngo-dysesthesias.<sup>119,120</sup> Patients need to be warned not to drink cold fluids in the days around their oxaliplatin infusions. Lengthening the infusion duration from 2 h to 6 h can prevent recurrence of the pseudolaryngo spasm.

Late-onset neuropathy with oxaliplatin is cumulative, dose-dependent, and typically consists of a sensory, symmetric distal axonal neuropathy without motor involvement. Incidence and severity are related predominantly to cumulative dose. For example, the incidence of grade 3 neuropathy with cumulative doses of 850 mg/m<sup>2</sup> is 10–15%, and rises thereafter. As the clinical situation permits, if significant neuropathy develops during treatment, we suggest discontinuing oxaliplatin and switching to a non-oxaliplatin chemotherapy to permit as much recovery as possible before reintroducing oxaliplatin.

#### **Experience of pharmacotherapy on post-transplantation CRC**

Pharmacotherapy is of particular important for the treatment of CRC. The median overall survival (OS) is about 6 months in untreated mCRC. The OS has been extended up to 12–20 months by treatment with combined chemotherapy such as FOLFOX, FOLFIRI, and CapeOx. The OS has been further improved to 24–41 months after adding target agents.<sup>34</sup> A few cases have explored the pharmacotherapy of CRC in KTRs (Table 5).<sup>1, 121–124</sup>

Modified chemotherapy, such as single agent therapy, was attempted to improve the survival and compliance of CRC in KTRs. In a study by Liu and colleagues, three cases of advanced rectal cancer were treated with capecitabine adjuvant chemotherapy after surgery.<sup>1</sup> Gu and colleagues reported a case of a 57-year-old female with advanced rectal cancer who received three cycles of oxaliplatin chemotherapy.<sup>122</sup> Trivedi and colleagues also reported the application of continuous 5-FU infusion in a colon cancer patient. Chemotherapy was well tolerated in all these patients.<sup>124</sup>

The application of combination therapy for post-transplantation CRC has been explored. In our previous study, an advanced colon cancer was treated with three cycles of FOLFOX chemotherapy. Plasma concentration of ISDs (CsA and SIR) and renal function were not affected by chemotherapy. No serious ADR was observed.<sup>121</sup>

Targeted therapy also has been reported. Mısıri and colleagues reported a case of post-transplantation mCRC treated with bevacizumab + FOLFIRI. The dose of bevacizumab was 5 mg/kg/day for 14 days. Proteinuria (2.5 g/day) existed before the start of the treatment, and increased to 4 g/day at the end of the fifth course.<sup>123</sup> It is wise to exclude patients with proteinuria before the start of antiangiogenic therapy.

#### **Potential management strategies**

##### *Immunosuppressive management in KTRs with CRC*

In KTRs with CRC, immunosuppressive therapy and antitumor therapies are both important, and constrict each other. Disturbance of immunosuppressive level, either under-immunosuppression or over-immunosuppression, can lead to other serious complications, such as graft rejection,



**Table 5.** Parameters of patients with CRC after renal transplantation.

Source	Age (years)	Gender	Location	ISD	Anticancer	ADR	Outcome
Fang <sup>121</sup>	39	Male	Colon	CSA+SIR	FOLFOX*3 cycles	—	PD after 3 cycles
Gu <sup>122</sup>	57	Female	Rectum	TAC+MMF+P	oxaliplatin	—	—
Liu <sup>1</sup>	68	Male	Rectum	—	Capecitabine*3 cycles	—	Died after 31 months
Liu <sup>1</sup>	44	Male	Rectum	—	Capecitabine*1 cycles	—	Alive after 21 months
Liu <sup>1</sup>	54	Male	Rectum	—	Capecitabine*1 cycles	—	Alive after 8 months
Musri <sup>123</sup>	64	Male	CRC	EVE+TAC	FOLFIRI+bevacizumab*5 cycles	proteinuria 4 g/day, GFR 24 ml/min	—
Trivedi <sup>124</sup>	44	Female	Colon	Pred+Aza+CsA	continuous 5-FU infusion	Not reported	Died after 7 months

5-FU, 5-fluorouracil; ADR, adverse drug reaction; Aza, azathioprine; CRC, colorectal cancer; CSA, cyclosporin A; EVE, Everolimus; ISD, immunosuppressive drugs; MMF, mycophenolate mofetil; P, prednisone; PD, programmed cell death; TAC, Tacrolimus.

infection, and renal dysfunction. Those are also contraindications for antitumor therapy. Preservation of graft function is of primary importance, and informs the basis of antitumor therapy. Hence our suggestions of ISDs for KTRs with CRC are similar to KDIGO clinical practice guidelines, which recommend TAC as the first-line CNI, and MMF as the first-line antiproliferative agent. CsA is recommended as the alternative CNI, and SIR as the alternative antiproliferative agent.

Although lacking trial-based evidence, judicious reduction in immunosuppression with regular monitoring of disease progression and graft function may be warranted, particularly among those with high-grade and advanced malignancy. We suggest KTRs with CRC reach the trough concentration recommended by KDIGO guideline for KTRs after 1 year of renal transplant, which is that the trough concentration of TAC should be more than 5–10 ng/ml. We also recommend intensive therapeutic drug monitoring of ISD such as testing of trough concentration before and after anticancer therapy.

#### *Antitumor management in KTRs with CRC*

For patients with indication of adjuvant chemotherapy, FOLFOX or CapeOx is preferred for patients who can tolerate combined chemotherapy, and

5-FU/LV or capecitabine is recommended for those patients who cannot tolerate combined chemotherapy. These regimens are also recommended as first-line palliative chemotherapy, and can be combined with molecular targeted drugs (anti-angiogenesis drugs and anti-EGFR MoAbs). FOLFIRI only can be used in oxaliplatin-refractory patients. In subsequent therapy, aflibercept, trifluridine-tipiracil, and tegafur-uracil are suggested for these patients.

#### *Pharmaceutical care in KTRs with CRC*

Regardless of the antitumor regimen, trough concentration of ISDs, renal function, blood routine, and cardiac function should be intensively monitored. DPD and TYMS genotype should be detected in order to make an early decision regarding fluoropyrimidines dose reduction. Attention should be paid to neurotoxicity if an oxaliplatin-based regimen (CapeOx or FOLFOX) is chosen. Blood pressure and urine protein excretion should be intensively monitored if an antiangiogenic drug is added. Attention should be paid to hypomagnesemia and skin toxicity if anti-EGFR MoAb is added.

Irinotecan has the risk of severe diarrhea, which substantially increased the exposure of TAC. Irinotecan also has the potential of competing for

UGT1A, which might lead to the accumulation of MMF and irinotecan. Hence irinotecan-based regimens (FOLFIRI and FOLFOXIRI) are not preferred, but can be used in oxaliplatin-refractory patients. UGT1A1 genotype should be determined before the initiation of irinotecan-based regimens, and the starting dose of irinotecan should be lowered in patients known to be homozygous for UGT1A1\*28. Cetuximab or panitumumab combined with FOLFIRI are not recommended for the synergistic effect of diarrhea.

Nivolumab or pembrolizumab should be avoided due to their immune enhancement effect. Regorafenib is not recommended because of its inhibiting effect on UGT1A and potential competition for CYP3A4. Attention should be paid to the interaction between ISDs and PPIs. All PPIs reduce absorption of MMF, and omeprazole might inhibit CYP3A4-mediated metabolism of TAC.

### Conclusion

Antitumor pharmacotherapy of CRC in KTRs is a dilemma. Impediments such as drug interactions and potential ADR really exist. But these should not be contraindications for KTRs to adopt anticancer therapy. Toxicity can be minimized by selecting regimens and managing ADR, and a few cases have also proved the feasibility of pharmacotherapy in these patients. We also provide potential antitumor regimens for the adjuvant, palliative, and subsequent therapy of CRC in KTRs. Future clinical trials or case reports are needed to confirm the relative benefit and risk of antitumor pharmacotherapy in KTRs.

### Acknowledgments

Authors Yuanyuan Fu and Chengheng Liao contributed equally.

### Funding

The author(s) received no financial support for the research, authorship, and publication of this article.

### Conflict of interest statement

The authors declare that there is no conflict of interest.

### Reference

1. Liu HY, Liang XB, Li YP, *et al.* Treatment of advanced rectal cancer after renal transplantation. *World J Gastroenterol* 2011; 17: 2058–2060.
2. Papaconstantinou HT, Sklow B, Hanaway MJ, *et al.* Characteristics and survival patterns of solid organ transplant patients developing de novo colon and rectal cancer. *Dis Colon Rectum* 2004; 47: 1898–1903.
3. Demir T, Ozel L, Gokce AM, *et al.* Cancer screening of renal transplant patients undergoing long-term immunosuppressive therapy. *Transplant Proc* 2015; 47: 1413–1417.
4. Park JM, Choi MG, Kim SW, *et al.* Increased incidence of colorectal malignancies in renal transplant recipients: a case control study. *Am J Transplant* 2010; 10: 2043–2050.
5. Webster AC, Craig JC, Simpson JM, *et al.* Identifying high risk groups and quantifying absolute risk of cancer after kidney transplantation: a cohort study of 15,183 recipients. *Am J Transplant* 2007; 7: 2140–2151.
6. Stewart T, Henderson R, Grayson H, *et al.* Reduced incidence of rectal cancer, compared to gastric and colonic cancer, in a population of 73,076 men and women chronically immunosuppressed. *Clin Cancer Res* 1997; 3: 51–55.
7. Adami J, Gabel H, Lindelof B, *et al.* Cancer risk following organ transplantation: a nationwide cohort study in Sweden. *Br J Cancer* 2003; 89: 1221–1227.
8. Birkeland SA. Malignant tumors in renal transplant patients. The Scandia transplant material. *Cancer* 1983; 51: 1571–1575.
9. Birkeland SA, Lokkegaard H and Storm HH. Cancer risk in patients on dialysis and after renal transplantation. *Lancet* 2000; 355: 1886–1887.
10. Blohme I and Brynner H. Malignant disease in renal transplant patients. *Transplantation* 1985; 39: 23–25.
11. Cheung CY, Lam MF, Chu KH, *et al.* Malignancies after kidney transplantation: Hong Kong renal registry. *Am J Transplant* 2012; 12: 3039–3046.
12. Collett D, Mumford L, Banner NR, *et al.* Comparison of the incidence of malignancy in recipients of different types of organ: a UK Registry audit. *Am J Transplant* 2010; 10: 1889–1896.
13. Disney AP. Demography and survival of patients receiving treatment for chronic renal failure in Australia and New Zealand: report on dialysis and renal transplantation treatment from the Australia and New Zealand Dialysis and Transplant Registry. *Am J Kidney Dis* 1995; 25: 165–175.

14. Ebisui C, Okazaki M, Kanai T, *et al.* Clinicopathological study of colorectal cancers after renal transplantation. *Transplant Proc* 2000; 32: 1984–1985.
15. Hoover R and Fraumeni JF Jr. Risk of cancer in renal-transplant recipients. *Lancet* 1973; 2: 55–57.
16. Hsiao FY and Hsu WW. Epidemiology of post-transplant malignancy in Asian renal transplant recipients: a population-based study. *Int Urol Nephrol* 2014; 46: 833–838.
17. Ishikawa N, Tanabe K, Tokumoto T, *et al.* Clinical study of malignancies after renal transplantation and impact of routine screening for early detection: a single-center experience. *Transplant Proc* 2000; 32: 1907–1910.
18. Kasiske BL and Danpanich E. Malignancies in renal transplant recipients. *Transplant Proc* 2000; 32: 1499–1500.
19. Ju MK, Joo DJ, Kim SJ, *et al.* Chronologically different incidences of post-transplant malignancies in renal transplant recipients: single center experience. *Transpl Int* 2009; 22: 644–653.
20. Ho HT, Russ G, Stephens J, *et al.* Colorectal cancer in patients with chronic renal failure: the effect of dialysis or renal transplantation. *Colorectal Dis* 2002; 4: 193–196.
21. Kasiske BL, Snyder JJ, Gilbertson DT, *et al.* Cancer after kidney transplantation in the United States. *Am J Transplant* 2004; 4: 905–913.
22. Kehinde EO, Petermann A, Morgan JD, *et al.* Triple therapy and incidence of de novo cancer in renal transplant recipients. *Br J Surg* 1994; 81: 985–986.
23. Kim JY, Ju MK, Kim MS, *et al.* Clinical characteristics and treatment outcomes of colorectal cancer in renal transplant recipients in Korea. *Yonsei Med J* 2011; 52: 454–462.
24. Krynitz B, Edgren G, Lindelof B, *et al.* Risk of skin cancer and other malignancies in kidney, liver, heart and lung transplant recipients 1970 to 2008—a Swedish population-based study. *Int J Cancer* 2013; 132: 1429–1438.
25. Kwon JH, Koh SJ, Kim JY, *et al.* Prevalence of advanced colorectal neoplasm after kidney transplantation: surveillance based on the results of screening colonoscopy. *Dig Dis Sci* 2015; 60: 1761–1769.
26. Kyllonen L, Pukkala E and Eklund B. Cancer incidence in a kidney-transplanted population. *Transpl Int* 1994; 7(Suppl 1): S350–S352.
27. Kyllonen L, Salmela K and Pukkala E. Cancer incidence in a kidney-transplanted population. *Transpl Int* 2000; 13(Suppl 1): S394–398.
28. Langer RM, Jaray J, Toth A, *et al.* De novo tumors after kidney transplantation: the Budapest experience. *Transpl Proc* 2003; 35: 1396–1398.
29. Li WH, Chen YJ, Tseng WC, *et al.* Malignancies after renal transplantation in Taiwan: a nationwide population-based study. *Nephrol Dial Transplant* 2012; 27: 833–839.
30. Parikshak M, Pawlak SE, Eggenberger JC, *et al.* The role of endoscopic colon surveillance in the transplant population. *Dis Colon Rectum* 2002; 45: 1655–1660.
31. Penn I. Tumors after renal and cardiac transplantation. *Hematol Oncol Clin North Am* 1993; 7: 431–445.
32. Popov Z, Ivanovski O, Kolevski P, *et al.* De novo malignancies after renal transplantation—a single-center experience in the Balkans. *Transpl Proc* 2007; 39: 2589–2591.
33. Rostami Z, Einollahi B, Lessan-Pezeshki M, *et al.* Old male living renal transplant recipients more likely to be at risk for colorectal cancer. *Transpl Proc* 2011; 43: 588–589.
34. Saidi RF, Dudrick PS and Goldman MH. Colorectal cancer after renal transplantation. *Transplant Proc* 2003; 35: 1410–1412.
35. Vajdic CM, McDonald SP, McCredie MR, *et al.* Cancer incidence before and after kidney transplantation. *JAMA* 2006; 296: 2823–2831.
36. Vilardell J, Oppenheimer F, Castela AM, *et al.* Cancer after transplantation in Catalonia. *Transplant Proc* 1992; 24: 124.
37. Villeneuve PJ, Schaubel DE, Fenton SS, *et al.* Cancer incidence among Canadian kidney transplant recipients. *Am J Transplant* 2007; 7: 941–948.
38. Zilinska Z, Sersenova M, Chrastina M, *et al.* Occurrence of malignancies after kidney transplantation in adults: slovak multicenter experience. *Neoplasma* 2017; 64: 311–317.
39. Pendon-Ruiz de Mier V, Navarro Cabello MD, Martinez Vaquera S, *et al.* Incidence and long-term prognosis of cancer after kidney transplantation. *Transplant Proc* 2015; 47: 2618–2621.
40. Vaeth M, Bauerlein CA, Pusch T, *et al.* Selective NFAT targeting in T cells ameliorates GvHD while maintaining antitumor activity. *Proc Natl Acad Sci USA* 2015; 112: 1125–1130.

41. Werneck MB, Hottz E, Bozza PT, *et al.* Cyclosporin A inhibits colon cancer cell growth independently of the calcineurin pathway. *Cell Cycle* 2012; 11: 3997–4008.
42. Hojo M, Morimoto T, Maluccio M, *et al.* Cyclosporine induces cancer progression by a cell-autonomous mechanism. *Nature* 1999; 397: 530–534.
43. Maluccio M, Sharma V, Lagman M, *et al.* Tacrolimus enhances transforming growth factor-beta1 expression and promotes tumor progression. *Transplantation* 2003; 76: 597–602.
44. Tsiptotis E, Gupta NR, Raman G, *et al.* Bioavailability, efficacy and safety of generic immunosuppressive drugs for kidney transplantation: a systematic review and meta-analysis. *Am J Nephrol* 2016; 44: 206–218.
45. Guba M, Graeb C, Jauch KW, *et al.* Pro- and anti-cancer effects of immunosuppressive agents used in organ transplantation. *Transplantation* 2004; 77: 1777–1782.
46. Knoll GA, Kokolo MB, Mallick R, *et al.* Effect of sirolimus on malignancy and survival after kidney transplantation: systematic review and meta-analysis of individual patient data. *BMJ* 2014; 349: g6679.
47. Swann PF, Waters TR, Moulton DC, *et al.* Role of postreplicative DNA mismatch repair in the cytotoxic action of thioguanine. *Science* 1996; 273: 1109–1111.
48. Huang M-Y, Lee H-H, Tsai H-L, *et al.* Comparison of efficacy and safety of preoperative Chemoradiotherapy in locally advanced upper and middle/lower rectal cancer. *Radiat Oncol* 2018; 13.
49. O'Neill JO, Edwards LB and Taylor DO. Mycophenolate mofetil and risk of developing malignancy after orthotopic heart transplantation: analysis of the transplant registry of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant* 2006; 25: 1186–1191.
50. Ross HJ, Cho J, Osann K, *et al.* Phase I/II trial of low dose cyclosporin A with EP for advanced non-small cell lung cancer. *Lung Cancer* 1997; 18: 189–198.
51. Zafari V, Hashemzadeh S, Hosseinpour Feizi M, *et al.* mRNA expression of nuclear factor of activated T-cells, cytoplasmic 2 (NFATc2) and peroxisome proliferator-activated receptor gamma (PPARG) transcription factors in colorectal carcinoma. *Bosn J Basic Med Sci* 2017; 17: 255–261.
52. Mody K and Bekaii-Saab T. Clinical trials and progress in metastatic colon cancer. *Surg Oncol Clin N Am* 2018; 27: 349–365.
53. Casadaban L, Rauscher G, Aklilu M, *et al.* Adjuvant chemotherapy is associated with improved survival in patients with stage II colon cancer. *Cancer* 2016; 122: 3277–3287.
54. Heinemann V, von Weikersthal LF, Decker T, *et al.* FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2014; 15: 1065–1075.
55. Van Cutsem E, Cervantes A, Adam R, *et al.* ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol* 2016; 27: 1386–1422.
56. Jawed I, Wilkerson J, Prasad V, *et al.* Colorectal cancer survival gains and novel treatment regimens: a systematic review and analysis. *JAMA Oncol* 2015; 1: 787–795.
57. Hecht JR. Gastrointestinal toxicity of irinotecan. *Oncology* 1998; 12: 72–78.
58. Mego M, Chovanec J, Vochyanova-Andrezalova I, *et al.* Prevention of irinotecan induced diarrhea by probiotics: a randomized double blind, placebo controlled pilot study. *Complement Ther Med* 2015; 23: 356–362.
59. Zhang L, Xing X, Meng F, *et al.* Oral fluoropyrimidine versus intravenous 5-fluorouracil for the treatment of advanced gastric and colorectal cancer: meta-analysis. *J Gastroenterol Hepatol* 2018; 33: 209–225.
60. Lorient MA, Ciccolini J, Thomas F, *et al.* [Dihydropyrimidine dehydrogenase (DPD) deficiency screening and securing of fluoropyrimidine-based chemotherapies: update and recommendations of the French GPCO-Umicancer and RNP Gx networks]. *Bull Cancer* 2018; 105: 397–407.
61. Naghibalhossaini F, Shefaghat M, Mansouri A, *et al.* The impact of thymidylate synthase and methylenetetrahydrofolate reductase genotypes on sensitivity to 5-fluorouracil treatment in colorectal cancer cells. *Acta Med Iran* 2017; 55: 751–758.
62. Fornasier G, Francescon S and Baldo P. An update of efficacy and safety of cetuximab in metastatic colorectal cancer: a narrative review. *Adv Ther* 2018; 35: 1497–1509.
63. Battaglin F, Dadduzio V, Bergamo F, *et al.* Anti-EGFR monoclonal antibody panitumumab for the treatment of patients with metastatic colorectal cancer: an overview of current practice



- and future perspectives. *Expert Opin Biol Ther* 2017; 17: 1297–1308.
64. Miroddi M, Sterrantino C, Simonelli I, *et al.* Risk of grade 3–4 diarrhea and mucositis in colorectal cancer patients receiving anti-EGFR monoclonal antibodies regimens: a meta-analysis of 18 randomized controlled clinical trials. *Crit Rev Oncol Hematol* 2015; 96: 355–371.
  65. Zhang D, Ye J, Xu T, *et al.* Treatment related severe and fatal adverse events with cetuximab in colorectal cancer patients: a meta-analysis. *J Chemother* 2013; 25: 170–175.
  66. Fernandez A, Villafruela J, Amezcua Y, *et al.* Mycophenolate mofetil absorption quotient: interest to clinical practice. *Transplant Proc* 2009; 41: 2317–2319.
  67. Heller T, van Gelder T, Budde K, *et al.* Plasma concentrations of mycophenolic acid acyl glucuronide are not associated with diarrhea in renal transplant recipients. *Am J Transplant* 2007; 7: 1822–1831.
  68. Maes BD, Lemahieu W, Kuypers D, *et al.* Differential effect of diarrhea on FK506 versus cyclosporine A trough levels and resultant prevention of allograft rejection in renal transplant recipients. *Am J Transplant* 2002; 2: 989–992.
  69. van Boekel GA, Aarnoutse RE, van der Heijden JJ, *et al.* Effect of mild diarrhea on tacrolimus exposure. *Transplantation* 2012; 94: 763–767.
  70. Lien YH. Top 10 things primary care physicians should know about maintenance immunosuppression for transplant recipients. *Am J Med* 2016; 129: 568–572.
  71. Shamliyan TA, Middleton M and Borst C. Patient-centered outcomes with concomitant use of proton pump inhibitors and other drugs. *Clin Ther* 2017; 39: 404–427.e436.
  72. Patel KS, Stephany BR, Barnes JF, *et al.* Renal transplant acute rejection with lower mycophenolate mofetil dosing and proton pump inhibitors or histamine-2 receptor antagonists. *Pharmacotherapy* 2017; 37: 1507–1515.
  73. Rupprecht K, Schmidt C, Raspe A, *et al.* Bioavailability of mycophenolate mofetil and enteric-coated mycophenolate sodium is differentially affected by pantoprazole in healthy volunteers. *J Clin Pharmacol* 2009; 49: 1196–1201.
  74. Knorr JP, Sjeime M, Braitman LE, *et al.* Concomitant proton pump inhibitors with mycophenolate mofetil and the risk of rejection in kidney transplant recipients. *Transplantation* 2014; 97: 518–524.
  75. Katsakiori PF, Papapetrou EP, Goumenos DS, *et al.* Investigation of clinical interaction between omeprazole and tacrolimus in CYP3A5 non-expressors, renal transplant recipients. *Ther Clin Risk Manag* 2010; 6: 265–269.
  76. Takahashi K, Yano I, Fukuhara Y, *et al.* Distinct effects of omeprazole and rabeprazole on the tacrolimus blood concentration in a kidney transplant recipient. *Drug Metab Pharmacokinet* 2007; 22: 441–444.
  77. Benet LZ and Hoener BA. Changes in plasma protein binding have little clinical relevance. *Clin Pharmacol Ther* 2002; 71: 115–121.
  78. Zou LW, Jin Q, Wang DD, *et al.* Carboxylesterase inhibitors: an update. *Curr Med Chem* 2018; 25: 1627–1649.
  79. Fujiyama N, Miura M, Kato S, *et al.* Involvement of carboxylesterase 1 and 2 in the hydrolysis of mycophenolate mofetil. *Drug Metab Dispos* 2010; 38: 2210–2217.
  80. Quinney SK, Sanghani SP, Davis WI, *et al.* Hydrolysis of capecitabine to 5'-deoxy-5-fluorocytidine by human carboxylesterases and inhibition by loperamide. *J Pharmacol Exp Ther* 2005; 313: 1011–1016.
  81. Laizure SC, Herring V, Hu Z, *et al.* The role of human carboxylesterases in drug metabolism: have we overlooked their importance? *Pharmacotherapy* 2013; 33: 210–222.
  82. Hatfield MJ, Tsurkan L, Garrett M, *et al.* Organ-specific carboxylesterase profiling identifies the small intestine and kidney as major contributors of activation of the anticancer prodrug CPT-11. *Biochem Pharmacol* 2011; 81: 24–31.
  83. Xu G, Zhang W, Ma MK, *et al.* Human carboxylesterase 2 is commonly expressed in tumor tissue and is correlated with activation of irinotecan. *Clin Cancer Res* 2002; 8: 2605–2611.
  84. Tlemsani C, Huillard O, Arrondeau J, *et al.* Effect of glucuronidation on transport and tissue accumulation of tyrosine kinase inhibitors: consequences for the clinical management of sorafenib and regorafenib. *Expert Opin Drug Metab Toxicol* 2015; 11: 785–794.
  85. Pedoem A, Azoulay-Alfaguter I, Strazza M, *et al.* Programmed death-1 pathway in cancer and autoimmunity. *Clin Immunol* 2014; 153: 145–152.
  86. Homet Moreno B and Ribas A. Anti-programmed cell death protein-1/ligand-1 therapy in different cancers. *Br J Cancer* 2015; 112: 1421–1427.



87. Burdmann EA, Andoh TF, Yu L, *et al.* Cyclosporine nephrotoxicity. *Semin Nephrol* 2003; 23: 465–476.
88. Fishman JA. Infection in solid-organ transplant recipients. *N Engl J Med* 2007; 357: 2601–2614.
89. Blumberg EA, Danziger-Isakov L, Kumar D, *et al.* Foreword: guidelines 3. *Am J Transplant* 2013; 13(Suppl. 4): 1–2.
90. Fischer SA and Avery RK; AST Infectious Disease Community of Practice. Screening of donor and recipient prior to solid organ transplantation. *Am J Transplant* 2004; 4(Suppl. 10): 10–20.
91. Weycker D, Li X, Edelsberg J, *et al.* Risk and consequences of chemotherapy-induced febrile neutropenia in patients with metastatic solid tumors. *J Oncol Pract* 2015; 11: 47–54.
92. Guo Y, Xiong BH, Zhang T, *et al.* XELOX vs. FOLFOX in metastatic colorectal cancer: an updated meta-analysis. *Cancer Invest* 2016; 34: 94–104.
93. Petrelli F, Cabiddu M and Barni S. 5-Fluorouracil or capecitabine in the treatment of advanced colorectal cancer: a pooled-analysis of randomized trials. *Med Oncol* 2012; 29: 1020–1029.
94. Sonis ST, Elting LS, Keefe D, *et al.* Perspectives on cancer therapy-induced mucosal injury: pathogenesis, measurement, epidemiology, and consequences for patients. *Cancer* 2004; 100: 1995–2025.
95. Chionh F, Lau D, Yeung Y, *et al.* Oral versus intravenous fluoropyrimidines for colorectal cancer. *Cochrane Database Syst Rev* 2017; 7: CD008398.
96. Vermorken JB, Stohlmacher-Williams J, Davidenko I, *et al.* Cisplatin and fluorouracil with or without panitumumab in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck (SPECTRUM): an open-label phase 3 randomised trial. *Lancet Oncol* 2013; 14: 697–710.
97. Platz KP, Mueller AR, Blumhardt G, *et al.* Nephrotoxicity following orthotopic liver transplantation. A comparison between cyclosporine and FK506. *Transplantation* 1994; 58: 170–178.
98. Lucey MR, Abdelmalek MF, Gagliardi R, *et al.* A comparison of tacrolimus and cyclosporine in liver transplantation: effects on renal function and cardiovascular risk status. *Am J Transplant* 2005; 5: 1111–1119.
99. Baird R, Biondo A, Chhaya V, *et al.* Toxicity associated with capecitabine plus oxaliplatin in colorectal cancer before and after an institutional policy of capecitabine dose reduction. *Br J Cancer* 2011; 104: 43–50.
100. Fukasawa H, Furuya R, Yasuda H, *et al.* Anti-cancer agent-induced nephrotoxicity. *Anticancer Agents Med Chem* 2014; 14: 921–927.
101. Grothey A, Van Cutsem E, Sobrero A, *et al.* Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013; 381: 303–312.
102. Arnold D, Fuchs CS, Tabernero J, *et al.* Meta-analysis of individual patient safety data from six randomized, placebo-controlled trials with the antiangiogenic VEGFR2-binding monoclonal antibody ramucirumab. *Ann Oncol* 2017; 28: 2932–2942.
103. Grothey A, Sobrero AF, Shields AF, *et al.* Duration of adjuvant chemotherapy for stage III colon cancer. *N Engl J Med* 2018; 378: 1177–1188.
104. An MM, Zou Z, Shen H, *et al.* Incidence and risk of significantly raised blood pressure in cancer patients treated with bevacizumab: an updated meta-analysis. *Eur J Clin Pharmacol* 2010; 66: 813–821.
105. Jhaveri KD, Sakhiya V, Wanchoo R, *et al.* Renal effects of novel anticancer targeted therapies: a review of the food and drug administration adverse event reporting system. *Kidney Int* 2016; 90: 706–707.
106. Berardi R, Santoni M, Rinaldi S, *et al.* Risk of hyponatraemia in cancer patients treated with targeted therapies: a systematic review and meta-analysis of clinical trials. *PLoS One* 2016; 11: e0152079.
107. Saif MW, Shah MM and Shah AR. Fluoropyrimidine-associated cardiotoxicity: revisited. *Expert Opin Drug Saf* 2009; 8: 191–202.
108. Kwakman JJ, Simkens LH, Mol L, *et al.* Incidence of capecitabine-related cardiotoxicity in different treatment schedules of metastatic colorectal cancer: a retrospective analysis of the CAIRO studies of the Dutch Colorectal Cancer Group. *Eur J Cancer* 2017; 76: 93–99.
109. Kosmas C, Kallistratos MS, Kopterides P, *et al.* Cardiotoxicity of fluoropyrimidines in different schedules of administration: a prospective study. *J Cancer Res Clin Oncol* 2008; 134: 75–82.
110. Jensen SA, Hasbak P, Mortensen J, *et al.* Fluorouracil induces myocardial ischemia with increases of plasma brain natriuretic peptide

- and lactic acid but without dysfunction of left ventricle. *J Clin Oncol* 2010; 28: 5280–5286.
111. Lestuzzi C, Vaccher E, Talamini R, *et al.* Effort myocardial ischemia during chemotherapy with 5-fluorouracil: an underestimated risk. *Ann Oncol* 2014; 25: 1059–1064.
112. Van Cutsem E, Tabernero J, Lakomy R, *et al.* Addition of aflibercept to fluorouracil, leucovorin, and irinotecan improves survival in a phase III randomized trial in patients with metastatic colorectal cancer previously treated with an oxaliplatin-based regimen. *J Clin Oncol* 2012; 30: 3499–3506.
113. Owczarek W, Slowinska M, Lesiak A, *et al.* The incidence and management of cutaneous adverse events of the epidermal growth factor receptor inhibitors. *Postepy Dermatol Alergol* 2017; 34: 418–428.
114. Stanculeanu DL, Zob D, Toma OC, *et al.* Cutaneous toxicities of molecular targeted therapies. *Maedica* 2017; 12: 48–54.
115. Petrelli F, Ardito R, Ghidini A, *et al.* Different toxicity of cetuximab and panitumumab in metastatic colorectal cancer treatment: a systematic review and meta-analysis. *Oncology* 2018; 94: 191–199.
116. Eilers RE Jr, Gandhi M, Patel JD, *et al.* Dermatologic infections in cancer patients treated with epidermal growth factor receptor inhibitor therapy. *J Natl Cancer Inst* 2010; 102: 47–53.
117. Lupu I, Voiculescu N, Bacalbasa N, *et al.* Cutaneous complications of molecular targeted therapy used in oncology. *J Med Life* 2016; 9: 19–25.
118. Pachman DR, Qin R, Seisler DK, *et al.* Clinical course of oxaliplatin-induced neuropathy: results from the randomized phase III trial N08CB (Alliance). *J Clin Oncol* 2015; 33: 3416–3422.
119. Velasco R, Bruna J, Briani C, *et al.* Early predictors of oxaliplatin-induced cumulative neuropathy in colorectal cancer patients. *J Neurol Neurosurg Psychiatry* 2014; 85: 392–398.
120. Alejandro LM, Behrendt CE, Chen K, *et al.* Predicting acute and persistent neuropathy associated with oxaliplatin. *Am J Clin Oncol* 2013; 36: 331–337.
121. Fang W. Chemotherapy in patient with colon cancer after renal transplantation: a case report with literature review. *Medicine* 2018; 97: e9678.
122. Gu JY, Zhang W and Gu RG. The clinical analysis of solid tumors after renal transplantation. *J Shandong Med Coll* 2017; 39: 6.
123. Musri FY, Mutlu H, Eryilmaz MK, *et al.* Experience of bevacizumab in a patient with colorectal cancer after renal transplantation. *J Cancer Res Ther* 2015; 11: 1018–1020.
124. Trivedi MH, Agrawal S, Muscato MS, *et al.* High grade, synchronous colon cancers after renal transplantation: were immunosuppressive drugs to blame? *Am J Gastroenterol* 1999; 94: 3359–3361.