CORRECTION

Correction: Elevated triglycerides and reduced high-density lipoprotein cholesterol are independently associated with the onset of advanced chronic kidney disease: a cohort study of 911,360 individuals from the United Kingdom

Misghina Weldegiorgis[1,2,3*] and Mark Woodward[1,4,5]

Correction: BMC Nephrol 23, 312 (2022) https://doi.org/10.1186/s12882-022-02932-2

Following publication of the original article [1], the authors informed us that the confidence interval is missing in **Table 2, Column 5, row Q2**. It should be **1.01 [0.95-1.09]**.

The original article has been corrected.

Reference

1. Weldegiorgis M, Woodward M. Elevated triglycerides and reduced high-density lipoprotein cholesterol are independently associated with the onset of advanced chronic kidney disease: a cohort study of 911,360 individuals from the United Kingdom. BMC Nephrol. 2022;23:312. https://doi.org/10.1186/s12882-022-02932-2.

Author details
[1]The George Institute for Global Health, School of Public Health, Imperial College London, London, UK. [2]St George and Sutherland Clinical School, Faculty of Medicine, University of New South Wales Sydney, Sydney, Australia. [3]Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK. [4]The George Institute for Global Health, Faculty of Medicine, University of New South Wales Sydney, Sydney, Australia. [5]Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA.

Published online: 21 September 2022

*Correspondence: m.weldegiorgis@imperial.ac.uk
[3] Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK
Full list of author information is available at the end of the article