

Letter to the Editor

Carcinoembryonic antigen and cytokeratin 20 in peritoneal cells of cancer patients: are we aware of what we are detecting by mRNA examination?

M Kowalewska^{*,1}, M Chechlinska² and R Nowak¹

¹Department of Molecular Biology, The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology, Roentgena 5, Warsaw 02-781, Poland; ²Department of Immunology, The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology, Roentgena 5, Warsaw 02-781, Poland

British Journal of Cancer (2008) 98, 512–513. doi:10.1038/sj.bjc.6604189 www.bjcancer.com

Published online 15 January 2008

© 2008 Cancer Research UK

Sir,

In a recently published paper in your journal, Katsuragi *et al* (2007) show that peritoneal recurrence in patients with early-stage gastric cancer relates to the levels of carcinoembryonic antigen (CEA) and/or cytokeratin 20 (CK 20) transcripts in the peritoneal lavage specimens from these patients. The authors are convinced that a positive RT-PCR result identifies the presence of 'micrometastatic cells'. In our opinion, there are no grounds for this presumption. We and others have raised the issue of the nonspecific expression of so-called tumour markers by activated lymphoid cells (Kowalewska *et al*, 2006). CEA and CK 20 transcripts are markers of limited value for the detection of cancer cells for several reasons. Both markers have been shown to be expressed by the haematopoietic cells of healthy volunteers and of patients with chronic inflammatory diseases (Jung *et al*, 1998, 1999; Champelovier *et al*, 1999; Vlems *et al*, 2002). The qRT-PCR-based techniques of CEA and CK 20 detection present the same limitations as conventional RT-PCR, as researchers were unable to set the cut-off values to distinguish between cancer and haematopoietic cell expression (Bustin *et al*, 2004; Schuster *et al*, 2004). Especially in the experimental setting studied by Katsuragi *et al* (2007), one should consider that CEA expression may be induced in peripheral blood mononuclear cells by cytokines (Jung *et al*, 1998; Goeminne *et al*, 1999), as increased concentrations of an array of cytokines characterise peritoneal fluids of cancer patients (Chechlinska *et al*, 2007). Furthermore, peritoneal lavage samples of cancer-free patients were found to contain CEA transcripts (Broll *et al*, 2001). In addition, normal granulocytes that are likely to be recruited to peritoneal cavity (Hau, 1990) have been shown to express CK 20 (Jung *et al*, 1999; Kruger *et al*, 2001). In fact, the increased numbers of granulocytes as well as of activated lymphocytes were demonstrated in the peritoneal fluids of gastric and colon cancer patients as compared with those of cancer-free controls (Olszewski *et al*, 2007).

Unfortunately, none of these limitations, although widely discussed in numerous papers, have been considered or even discussed by Katsuragi *et al* (2007). In effect, the observed correlation between the positive CEA and CK 20 qRT-PCR signals and cancer recurrence seems reasonably well documented, whereas the statement that the authors detect 'free cancer cells' in their study, expressed already in the title and then subsequently used through the methods section to discussion, is totally unwarranted.

Considering the above, the measurements of the markers described by Katsuragi *et al* (2007) may reflect the presence of inflammatory cells rather than micrometastatic cells: the more so because as many as 35% of cytology-negative/PCR-positive patients developed no peritoneal metastases. Nevertheless, prognosis for PCR-positive patients was shown to be significantly worse than for PCR-negative patients, and this was in accordance with the recent data of Crumley *et al* (2006) and Deans *et al* (2007), who linked inflammation symptoms with adverse prognosis in patients with gastric cancer. However, there are simpler and cheaper methods than qRT-PCR to assess inflammatory parameters.

The study of Katsuragi *et al* (2007) is also methodologically flawed. The *GAPDH* gene was used as an internal control and a reference gene. The *GAPDH* gene is known for the presence of its numerous pseudogenes, which makes it inadequate as a reference gene in non-DNased total RNA samples, such as those prepared by Katsuragi *et al* (2007). In addition, the reader learns nothing about the standardisation of RNA quantities subjected to reverse transcription, necessary to make a reliable comparison of samples containing different cell counts.

Finally, Katsuragi *et al* (2007) use the inexplicable and inappropriate term 'tumour nucleotides', and for reasons that are far from obvious the disseminated cancer cells are called 'isolated' cells, although no isolation or enrichment procedure was performed. The authors have not applied the idea of micrometastases detection in cancer cell-enriched populations, one of many detection methods focused to enhance specificity and the only one which produced an assay that received FDA clearance (Smerage and Hayes, 2006).

*Correspondence: Dr M Kowalewska; E-mail: magdak@coi.waw.pl
Published online 15 January 2008

REFERENCES

- Broll R, Weschta M, Windhoevel U, Berndt S, Schwandner O, Roblick U, Schiedeck TH, Schimmelpenning H, Bruch HP, Duchrow M (2001) Prognostic significance of free gastrointestinal tumor cells in peritoneal lavage detected by immunocytochemistry and polymerase chain reaction. *Langenbecks Arch Surg* **386**: 285–292
- Bustin SA, Siddiqi S, Ahmed S, Hands R, Dorudi S (2004) Quantification of cytokeratin 20, carcinoembryonic antigen and guanylyl cyclase C mRNA levels in lymph nodes may not predict treatment failure in colorectal cancer patients. *Int J Cancer* **108**: 412–417
- Champelovier P, Mongelard F, Seigneurin D (1999) CK20 gene expression: technical limits for the detection of circulating tumor cells. *Anticancer Res* **19**: 2073–2078
- Chechlińska M, Kaminska J, Markowska J, Kramar A, Steffen J (2007) Peritoneal fluid cytokines and the differential diagnosis of benign and malignant ovarian tumors and residual/recurrent disease examination. *Int J Biol Markers* **22**: 172–180
- Crumley AB, McMillan DC, McKernan M, McDonald AC, Stuart RC (2006) Evaluation of an inflammation-based prognostic score in patients with inoperable gastro-oesophageal cancer. *Br J Cancer* **94**: 637–641
- Deans DA, Wigmore SJ, de Beaux AC, Paterson-Brown S, Garden OJ, Fearon KC (2007) Clinical prognostic scoring system to aid decision-making in gastro-oesophageal cancer. *Br J Surg* **94**: 1501–1508
- Goeminne JC, Guillaume T, Salmon M, Machiels JP, D'Hondt V, Symann M (1999) Unreliability of carcinoembryonic antigen (CEA) reverse transcriptase-polymerase chain reaction (RT-PCR) in detecting contaminating breast cancer cells in peripheral blood stem cells due to induction of CEA by growth factors. *Bone Marrow Transplant* **24**: 769–775
- Hau T (1990) Bacteria, toxins, and the peritoneum. *World J Surg* **14**: 167–175
- Jung R, Krüger W, Hosch S, Holweg M, Kröger N, Gutensohn K, Wagener C, Neumaier M, Zander AR (1998) Specificity of reverse transcriptase polymerase chain reaction assays designed for the detection of circulating cancer cells is influenced by cytokines *in vivo* and *in vitro*. *Br J Cancer* **78**: 1194–1198
- Jung R, Petersen K, Kruger W, Wolf M, Wagener C, Zander A, Neumaier M (1999) Detection of micrometastasis by cytokeratin 20 RT-PCR is limited due to stable background transcription in granulocytes. *Br J Cancer* **81**: 870–873
- Katsuragi K, Yashiro M, Sawada T, Osaka H, Ohira M, Hirakawa K (2007) Prognostic impact of PCR-based identification of isolated tumour cells in the peritoneal lavage fluid of gastric cancer patients who underwent a curative R0 resection. *Br J Cancer* **97**: 550–556
- Kowalewska M, Chechlińska M, Markowicz S, Kober P, Nowak R (2006) The relevance of RT-PCR detection of disseminated tumour cells is hampered by the expression of markers regarded as tumour-specific in activated lymphocytes. *Eur J Cancer* **42**: 2671–2674
- Kruger WH, Jung R, Detlefsen B, Mumme S, Badbaran A, Brandner J, Renges H, Kroger N, Zander AR (2001) Interference of cytokeratin-20 and mammaglobin-reverse-transcriptase polymerase chain assays designed for the detection of disseminated cancer cells. *Med Oncol* **18**: 33–38
- Olszewski WL, Kubicka U, Tarnowski W, Bielecki K, Ziolkowska A, Wesolowska A (2007) Activation of human peritoneal immune cells in early stages of gastric and colon cancer. *Surgery* **141**: 212–221
- Schuster R, Max N, Mann B, Heufelder K, Thilo F, Grone J, Rokos F, Buhr HJ, Thiel E, Keilholz U (2004) Quantitative real-time RT-PCR for detection of disseminated tumor cells in peripheral blood of patients with colorectal cancer using different mRNA markers. *Int J Cancer* **108**: 219–227
- Smerage JB, Hayes DF (2006) The measurement and therapeutic implications of circulating tumour cells in breast cancer. *Br J Cancer* **94**: 8–12
- Vlems FA, Diepstra JH, Cornelissen IM, Ruers TJ, Ligtenberg MJ, Punt CJ, van Krieken JH, Wobbes T, van Muijen GN (2002) Limitations of cytokeratin 20 RT-PCR to detect disseminated tumour cells in blood and bone marrow of patients with colorectal cancer: expression in controls and downregulation in tumour tissue. *Mol Pathol* **55**: 156–163