



# Myasthenia gravis in pregnancy: Systematic review and case series

Harrison Banner<sup>1</sup> , Kirsten M Niles<sup>1</sup>, Michelle Ryu<sup>2</sup>,  
 Mathew Sermer<sup>1</sup>, Vera Bril<sup>3</sup> and Kellie E Murphy<sup>1</sup>

## Abstract

**Background:** Myasthenia gravis is an autoimmune disease which can impact pregnancy.

**Methods:** Six databases were systematically searched for studies with at least five subjects reporting pregnancy outcomes for women with myasthenia gravis in pregnancy. Assessment of bias was performed for all included studies. Forty-eight cases from our own centre were also included in the analysis.

**Results:** In total, 32 publications met inclusion criteria for systematic review, for a total of 33 unique data sets including 48 cases from our institution. Outcome data was available for 824 pregnancies. Spontaneous vaginal delivery occurred in 56.3% of pregnancies. Overall risk of myasthenia gravis exacerbation was 33.8% with a 6.4% risk of myasthenic crisis in pregnancy and 8.2% postpartum. The incidence risk of transient neonatal myasthenia gravis was 13.0%.

**Conclusions:** The current systematic review provides the best estimates of risk currently available to aid in counselling women with myasthenia gravis in pregnancy.

## Keywords

Myasthenia gravis, pregnancy, obstetric outcomes, obstetric complications, systematic review

Date Received: 7 July 2021; accepted: 19 July 2021

## Introduction

Myasthenia gravis (MG) is an antibody-mediated disorder affecting the neuromuscular junction.<sup>1</sup> The majority of affected individuals will have antibodies against acetylcholine receptor (AChR) or muscle specific kinase. Patients experience fluctuating weakness in the skeletal muscles, primarily affecting the ocular, bulbar, limb and respiratory muscles. Though uncommon, with a reported prevalence in the general population of 0.3–7.7 per 100,000,<sup>2–3</sup> MG incidence spikes for women during the second and third decades of life and therefore can impact pregnancy. Recent guidelines make recommendations about the optimal management of MG in pregnancy<sup>4, 5</sup> though high-quality data is lacking.

MG is a disease with a variable course, with episodic exacerbations requiring more intensive treatment. During an exacerbation, individuals experience worsening muscle weakness or other related symptoms and may require increased dosages or additional types of medication for symptom control. Myasthenic crisis on the other hand, is a life-threatening complication involving weakness of the respiratory and/or bulbar muscles leading to respiratory compromise requiring ventilatory support.<sup>6</sup>

MG treatment primarily involves symptomatic management with anticholinesterase agents, with steroids and other immunosuppressive agents used in more severe disease. For refractory cases or myasthenic crises, intravenous immunoglobulin (IVIG) and/or plasmapheresis can be employed. Thymectomy has been demonstrated to improve symptoms for individuals with a thymoma or those with thymic hyperplasia in young-onset AChR antibody cases.<sup>7</sup>

MG has a variable course in pregnancy. Though it may happen at any time, exacerbation seems most likely to occur in the first trimester and during the postpartum period.<sup>8</sup> Previous studies report overall pregnancy exacerbation rates on the order of 30%,<sup>9, 10</sup> though higher<sup>11, 12</sup> and lower<sup>13</sup> rates have been reported. The literature is similarly discordant with regard to pregnancy outcome, with some studies reporting increased rates of preterm prelabour rupture of membranes (PPROM), preterm birth (PTB), small for gestational age (SGA) babies and caesarean delivery,<sup>14, 15</sup>

and others reporting rates similar to that of the general population.<sup>16, 17</sup> Transient neonatal myasthenia gravis (TNMG) is a complication whereby the autoantibodies pass through the placenta causing a self-limited myasthenic syndrome in the neonate. Risk of TNMG is generally quoted to be 10–15%, with individual series demonstrating rates ranging between 3–4%<sup>15, 18</sup> and 33%.<sup>19</sup>

The present study sought to systematically review all reported case series (5 or more cases) of MG during pregnancy and to include a large case series from our institution in an effort to summarize the present literature and better understand the risks associated with MG in pregnancy.

## Materials and methods

The review was registered in the PROSPERO database (2019; registration number CRD42019143072) prior to study initiation. The methods were performed in accordance with the preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines (<http://www.prisma-statement.org>).

<sup>1</sup>Division of Maternal Fetal Medicine, Department of Obstetrics and Gynaecology, Mount Sinai Hospital, University of Toronto, Canada

<sup>2</sup>Sidney Liswood Library, Mount Sinai Hospital, University of Toronto, Canada

<sup>3</sup>Division of Neurology, University Health Network, University of Toronto, Canada

### Corresponding author:

Harrison Banner, Department of Obstetrics and Gynaecology, Western University, London Health Sciences Centre, 800 Commissioners Rd E – Rm B2 412, London, Ontario, N6A 5W9, Canada.  
 Email: [harrison.banner@lhsc.on.ca](mailto:harrison.banner@lhsc.on.ca)

## Search strategy

The following electronic bibliographic databases were searched using a comprehensive search strategy developed by an information specialist (M.R.): (1) Ovid MEDLINE, (2) Ovid EMBASE, (3) Ovid EBM Reviews – Cochrane Central Register of Controlled Trials, (4) PubMed (Non-Medline records), (5) Web of Science and (6) LILACS (Latin American & Caribbean Health Science Literature). Additionally, all registered controlled trials were screened in ClinicalTrials.gov, the WHO International Clinical Trials Registry Platform and the International Standard Randomised Controlled Trial Number Registry. The search strategy was structured according to the 2015 Peer Reviewed Electronic Search Strategies Guidelines. Subject headings and free text terms related to ‘myasthenia gravis’, ‘pregnancy’ and ‘obstetric delivery’ were included. No restrictions were applied to publication year, language or age. All databases were searched for relevant references published from their inception up to 31 July 2020. All references and duplicate records were managed and screened using EndNote citation management software and Covidence systematic review software (Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia; www.covidence.org).

Inclusion criteria included randomized controlled trials, prospective or retrospective cohort studies and case series of at least five subjects with MG in pregnancy. Exclusion criteria included reviews without individual patient-level data, publication of data reported in previous publications and abstract presentations.

## Data extraction and assessment of risk of bias

Two authors (HB and KMN) independently screened all titles and abstracts. Relevant studies were then pulled for full text review. These two authors independently reviewed, extracted data and assessed risk of bias, using “Quality Assessment Tool for Case Series” (NIH.gov), for each full text article. Disagreements were settled via consensus or if necessary, with discussion involving a third author (KEM).

Data was extracted from included studies using a standardized data extraction form. Extracted data included disease information, including MG treatment, proportion of pregnancies with exacerbation, stable disease and asymptomatic/symptomatic improvement in pregnancy, and the rate of myasthenic crisis during pregnancy and up to 6 weeks postpartum. Data was also extracted regarding pregnancy outcomes, including mode of delivery as well as the rate of pregnancy complications, including: PTB less than 34 weeks and less than 37 weeks gestational age, PPROM, SGA less than 10th centile (based on birthweight for given gestational age) and the proportion of neonates experiencing TNMG. Studies were screened to exclude data sets which included previously published patient data.

## Institutional case series

After local Research Ethics Board approval (MSH REB #19-0183-C, 26 July 2019), a retrospective chart review of all pregnant women with myasthenia gravis cared for at Mount Sinai Hospital, Toronto, was conducted from the Medical Disorders of Pregnancy clinic database from 1 January 2000 to 31 August 2019.

Cases were included if there was a documented diagnosis of myasthenia gravis made prior to or during the pregnancy based on clinical or electrophysiological criteria. Cases of congenital myasthenic syndromes were excluded as this was felt to represent a separate disease entity. Only cases which were followed for the duration of pregnancy at our institution with available birth outcomes were included.

For the purposes of this study, each pregnancy was treated as a separate case for women having more than one pregnancy. Although multiple pregnancies in the same individual will have correlated outcomes, the

previously published literature on this topic included subsequent pregnancies in the same woman and it was not possible to assign pregnancies to individuals in these studies. Therefore, this convention was followed in our case series to maintain consistency. This was felt to be reasonable as comparative statistics were not planned.

For each case, data regarding demographics, diagnosis, MG treatment, disease course during pregnancy and up to 6 weeks postpartum, mode of delivery and maternal-fetal-neonatal outcomes were collected. Disease course was defined as “asymptomatic/symptomatic improvement” if the woman had an improvement in their symptoms or remained asymptomatic throughout the pregnancy, “unchanged” if symptoms were stable throughout pregnancy and “exacerbation” if the woman experienced a documented increase in the frequency or severity of symptoms or required an increase in medication dose or an additional treatment modality in pregnancy. Neonates were considered to be SGA if their birthweight was less than the 10th percentile for gestational age based on the growth curves created by Kramer et al.<sup>20</sup>

## Statistical methods

Descriptive statistics were performed using SPSS (IBM SPSS Statistics, Version 23.0, Armonk, NY: IBM Corp).

## Results

Our literature search yielded 980 citations. After removing 334 duplicates, 646 titles and abstracts were screened for inclusion. Of these, 227 were retrieved for full-text review and 32 ultimately were included in our review (Figure 1). Our case series data was included in the final review for a total of 33 unique data sets involving MG in pregnancy.

In our institutional case series, 39 women with MG had 48 pregnancies. The demographic information, MG treatment and pregnancy outcomes for women at our centre are summarized in Table 1. Individual patient level data is available as a Supplemental Table.

The data collected from the 32 studies<sup>8, 11, 16, 21–46</sup> included in the systematic review as well as our own case series are summarized in Table 2. The included studies represent a heterogeneous group from 22 different countries with case data collected from 1947 up until 2020. Twenty-eight of the included studies were case series, four used a retrospective cohort design and one case-control study was included. Sample sizes ranged from 5 to 163 subjects with MG in pregnancy. There was significant variability in the type of data collected between studies. The significant heterogeneity in study design and reporting precluded meaningful comparative analysis of the study data.


Risk of bias was assessed and this is summarized in Figure 2. Due to the nature of methodology used for these studies, the overall risk of bias is quite high.

## MG treatment in pregnancy

The proportion of women with a thymectomy prior to pregnancy ranged from 16.7%<sup>36</sup> to 100%.<sup>19, 37</sup> Across studies, the majority of women were treated with anti-cholinesterase medications during pregnancy, though a significant number also required prednisone.<sup>34, 39, 46</sup> More recent studies were more likely to include women treated with Azathioprine,<sup>39</sup> IVIG<sup>11</sup> or plasmapheresis.<sup>41</sup> Management changes over time reflect practice changes as new medications and evidence for treatment has become available.

## MG course in pregnancy

Across all studies reporting this outcome, 181/535 (33.8%) women experienced an exacerbation of their MG during pregnancy. While not all studies differentiated between stable disease and improvement, in those that did,



**Figure 1.** Preferred reporting items for systematic reviews and meta-analyses (PRISMA) flow diagram of systematic literature search for studies related to myasthenia gravis in pregnancy.

242/505 (47.9%) had stable disease and 100/505 (19.8%) had an improvement in their symptoms from baseline.

In studies reporting this outcome, 23/357 (6.4%) women experienced an episode of myasthenic crisis in pregnancy, and 31/380 (8.2%) experienced a myasthenic crisis during the postpartum period.

## Mode of delivery

The overall rate of caesarean delivery done for MG was 25/824 (3.0%), with an additional 249/824 (30.2%) women having a caesarean delivery for an obstetric indication. The rate of operative vaginal delivery was 86/824 (10.4%), with 464/824 (56.3%) of women having a spontaneous vaginal birth.

## Pregnancy complications

PTB less than 34 and 37 weeks' gestation occurred in 14/327 (4.3%) and 69/579 (11.9%) pregnancies, respectively. The overall risk of PPRM was 27/404 (6.7%).

In studies providing information on birthweight for gestational age, 55/390 (14.1%) neonates were smaller than the 10th percentile. The risk of TNMG was 89/686 (13.0%).

## Discussion

Myasthenia gravis is an uncommon neuromuscular condition which can have a significant impact on pregnancy. Previous publications have described a wide range of risk of MG exacerbation during pregnancy,

ranging in individual series from 0%<sup>37</sup> to 60%<sup>40, 47</sup> depending on the study. In our systematic review, 33.8% of women with MG overall experienced an exacerbation of their symptoms during pregnancy, a finding consistent with the generally quoted risk of 30–45%.<sup>9, 10</sup> Previous authors have stated that exacerbation is most likely to occur in the first trimester and in the postpartum period,<sup>8</sup> though the current review did not examine the timing of exacerbation within pregnancy since the timing of exacerbation was only reported in a small minority of studies.

Myasthenic crisis, an important and potentially life-threatening complication of MG, occurred in 6.4% of women during pregnancy and in 8.2% of women during the postpartum period.

The method by which women with MG give birth shows significant regional and temporal variation in the previously published literature, with the proportion of women having a vaginal birth ranging from 0%<sup>36</sup> to 100%<sup>35</sup> in some small series. Overall, 56.3% of births across all included studies occurred by SVD and an additional 10.4% were operative vaginal deliveries. In total, 33.3% of deliveries were via caesarean section, with 30.2% done for an obstetric indication. Only 3.0% of women had a caesarean delivery with MG as the indication.

For women with symptomatic MG at the time of birth, there is some concern that the voluntary striated muscles used during active pushing may be weakened by MG exacerbation and further compromised by excessive maternal effort, precipitating myasthenic crisis.<sup>48</sup> For women with exacerbation symptoms at the time of delivery, an assisted second stage of labour may decrease the need for maternal effort and minimize this risk. This has not been universally adopted as standard practice, and indeed current guidelines note that spontaneous vaginal delivery is the objective and should be encouraged.<sup>4</sup>

**Table 1.** Demographic information, disease course and pregnancy outcomes of women with myasthenia gravis in case series.

<b>Patient characteristics</b>		
Maternal age at delivery	(mean)	32.7
Parity	(mean)	1.75
	Nulliparous	22/48 (45.8%)
	Multiparous	26/48 (54.2%)
Disease type:	Ocular	8/48 (16.7%)
	Generalized	40/48 (83.3%)
MG diagnosed in pregnancy		2/48 (4.1%)
Thymectomy prior to pregnancy		34/48 (70.8%)
Treatment in pregnancy:	None	9/48 (18.8%)
	Pyridostigmine	36/48 (75.0%)
	Steroids	22/48 (45.8%)
	Azathioprine	11/48 (22.9%)
	IVIg	7/48 (14.6%)
	PLEX	6/48 (12.5%)
<b>Patient outcomes</b>		
Disease course in pregnancy:	Exacerbation	12/48 (25.0%)
	Unchanged	33/48 (68.8%)
	Asymptomatic/ symptomatic improvement	3/48 (6.3%)
Myasthenic crisis in pregnancy		4/48 (8.3%)
Myasthenic crisis in postpartum		0/48 (0%)
Planned mode of delivery:	SVD	37/48 (77.1%)
	Operative vaginal delivery	4/48 (8.3%)
	Caesarean section	7/48 (14.6%)
Actual mode of delivery:	SVD	22/48 (45.8%)
	Operative vaginal delivery	10/48 (20.8%)
	Caesarean section	16/48 (33.3%)
Delivery prior to 37 completed weeks		9/48 (18.8%)
Delivery prior to 34 completed weeks		5/48 (10.4%)
Small for gestational age		6/44 (12.5%)
PPROM		4/48 (8.3%)
TNMG		1/48 (2.0%)

IVIg: intravenous immunoglobulin; PLEX: plasmapheresis; SVD: spontaneous vaginal delivery; PPRM: preterm prelabour rupture of membranes; TNMG: transient neonatal myasthenia gravis.

Women with MG seem to be at increased risk of PPRM compared to the 3%<sup>49</sup> risk in the general population, with this complication occurring in 6.7% of all pregnancies in women with MG. This finding has been previously reported,<sup>8, 39</sup> though a pathophysiologic explanation remains elusive.

Whether or not MG increases the risk of preterm labour and birth has been the subject of previous controversy in the literature, with some populations, our own included, appearing to demonstrate an elevated risk.<sup>33</sup> Our systematic review did not find an elevated risk of PTB in women with MG, with overall risks of PTB of 11.9% prior to 37 weeks and 4.3% prior to 34 weeks, which is similar to the population risk.<sup>50</sup> Tanacan et al.<sup>8</sup> have previously suggested that an elevated risk of PTB may exist in the population

who experience exacerbation in pregnancy<sup>8</sup> though it was not possible to assess this in the current study.

Previous studies have demonstrated conflicting findings regarding the association between MG and fetal growth, with high rates of SGA babies reported in some series<sup>12, 16, 17</sup> but not others.<sup>11, 17, 41</sup> Our review found that 14.1% of babies born to mothers with MG had an infant with a birthweight less than the 10th percentile for gestational age overall.

Finally, the occurrence of TNMG in individual studies has ranged from 0%<sup>12, 21, 36, 37, 46</sup> to 33.3%.<sup>19, 34, 41</sup> In our large sample, the overall rate was 89/686 (13.0%). Previous studies have demonstrated a lack of correlation between anti-AChR antibody titre,<sup>42</sup> duration of illness and treatment modalities<sup>34</sup> and the risk of TNMG. The current understanding of risk factors for this complication remains limited. Women should be counselled about this risk and parents and care providers should be aware of the signs of this condition and the fact that there is no known association between this outcome and any predictive risk factor.

The current systematic review represents an accumulation of data, which to our knowledge has not previously been synthesized systematically. Previous estimates of risk and recommendations for management in the literature have been based on individual single-centre case series, the results of which differ considerably.

We used a search strategy which encompassed multiple databases and did not eliminate studies based on publication language which allowed us to capture a diverse group of data sets. By excluding studies with sample sizes less than 5, some of the bias inherent in the publication of extreme case studies was minimized. This study extracted data on a large range of factors related to the treatment and outcomes of MG in pregnancy, and due to the large total sample represents the best estimate of the true risk presently available.

The heterogeneous nature of the included studies means that data is represented from many different geographical regions. The fact that we did not use time or geographical exclusions increases the generalizability of our data, but does mean that our findings include regional and temporal variations in practice.

The main limitation of a study of this nature is the inclusion of low-quality evidence, as the majority of included studies were small and did not use a comparative design. Case series are particularly vulnerable to convenience sampling and selection bias, and may not appropriately represent the wider population of women with MG embarking on pregnancy. The lack of a comparator group in the majority of included studies compromises internal validity and means that results are presented in a descriptive manner rather than employing meaningful comparative statistical analysis. Finally, most studies reported on pregnancies rather than by individual patients. It is possible that women with less severe disease and fewer complications may have contributed more pregnancies to the overall number. Risk estimates may have been under- or over-estimated as a result of the nature and quality of the studies included in the review, however the large cumulative number of included women allows the risk estimates to be as accurate as possible.

For women with MG entering pregnancy, the current study allows the current best risk estimates for use in counselling regarding disease behaviour as well as the risk of myasthenic and pregnancy complications.

## Conclusion

Myasthenia gravis has important implications and considerations for pregnancy which should be discussed with women. A counselling summary based on the findings of our study is provided in Figure 3. Women with MG are at increased risk of requiring assisted vaginal delivery or caesarean delivery compared to the general population, though more than half of women with MG will be able to have a spontaneous vaginal birth. The overall risk of exacerbation in pregnancy is about 33%, with a small but

**Table 2.** Description of individual studies retrieved in the systematic review of myasthenia gravis in pregnancy.

Study characteristics		MG treatment in pregnancy									
Author Year	Country	Study Era	Design	Total population number	Previous Thymectomy	No treatment	Anti-cholinesterase	Prednisone	Azathioprine	IVIg	PLEX
Hoff 2003	Norway	1967–2000	Retrospective cohort	127	45/127	–	–	–	–	–	–
Braga 2016	Portugal	2005–2013	Case series	30	18/25	–	24/30	13/30	2/30	12/30	0/30
Assis 1983	Brazil	1983	Case series	39	–	–	–	–	–	–	–
Wen 2009	Taiwan	2001–2003	Retrospective cohort	163	11/163	–	–	–	–	–	–
Tellez-Zenteno 2004	Mexico	1996–2003	Case series	18	17/18	–	15/18	2/18	1/18	–	–
Tanacan 2019	Turkey	2010–2017	Retrospective cohort	27	7/27	–	–	–	–	–	–
Slama 2008	Tunisia	2008	Case series	5	–	–	–	–	–	–	–
Shi 2018	China	2004–2012	Case series	8	5/8	1/8	4/8	5/8	–	1/8	–
Ramirez 2006	France	1994–2003	Case series	36	12/18	5/34	27/34	13/34	6/34	0/34	2/34
Qi 2012	China	1983–2010	Case series	7	4/7	–	–	–	–	–	–
Picone 2003	France	1994–2002	Case series	13	8/13	0/13	9/13	1/13	1/13	1/13	1/13
Garcia 1989	Brazil	1989	Case series	6	1/6	0/6	6/6	1/6	0/6	0/6	0/6
Tsurane 2019	Japan	2016–2017	Case series	6	5/6	0/6	2/6	5/6	2/6	0/6	0/6
Thoulon 1978	France	1950–1978	Case series	24	13/24	–	–	–	–	–	–
Roth 2006	Switzerland	1986–2005	Retrospective cohort	15	7/15	–	–	–	–	–	–
Podciechowski 2005	Poland	1992–2004	Case series	13	6/13	–	12/13	–	–	–	–
Drazancic 1978	Yugoslavia	1963–1978	Case series	5	2/5	1/5	4/5	1/5	0/5	0/5	0/5
Kostera-Pruszyk 2005	Poland	2005	Case series	26	22/26	0/26	16/26	13/26	6/26	–	–
Mitchell 1992	Canada	1979–1989	Case series	11	8/11	–	–	–	–	–	–
Perucca 2006	Chile	1983–2004	Case series	19	7/19	0/19	19/19	1/19	0/19	0/19	0/19
Miller 1968	United States	1942–1962	Case series	7	–	1/7	6/7	–	–	–	–
Chabert 2004	France	1994–2002	Case series	10	–	0/10	10/10	–	1/10	1/10	1/10
Castro 1995	Brazil	1995	Case series	9	8/9	0/9	9/9	0/9	0/9	0/9	0/9
Gordon 1968	USA	1953–1968	Case series	11	–	–	–	–	–	–	–
Gametz 2017	Spain	2013–2014	Case series	5	5/5	0/5	0/5	0/5	0/5	5/5	0/5
Fraser 1953	England	1947–1952	Case series	14	4/7	1/7	3/7	–	–	–	–
Gvertic-Ahmetasevic 2008	Croatia	1998–2006	Case control	49	25/49	–	40/49	27/49	–	–	–
Ducci 2016	Brazil	1990–2015	Case series	34	8/24	2/34	16/17	19/34	7/34	1/34	0
Eden 1983	United States	1958–1981	Case series	12	12/12	0	9/12	4/12	0	0	0
Cheng 2007	Taiwan	1997–2002	Case series	13	7/12	–	6/13	–	–	–	–
Almeida 2010	Portugal	1985–2007	Case series	15	–	–	9/15	8/15	0/15	5/15	0/15
Djelimis 2002	Croatia	1972–1999	Case series	69	25/65	16/69	53/69	23/69	3/69	0/69	9/69
Banner 2020	Canada	2000–2019	Case series	48	17/24	9/48	34/48	11/24	11/48	7/48	6/48

(continued)

Table 2. Continued.


Author Year	Mode of delivery				MG course in pregnancy					Pregnancy complications				
	CS: MG as indication (%)	CS: Obstetric indication (%)	Operative vaginal delivery (%)	SVD (%)	Exacerbation (%)	Asymptomatic/improvement (%)	Unchanged (%)	Myasthenic crisis in pregnancy (%)	Myasthenic crisis postpartum (%)	PTB (<34 weeks)	PTB (<37 weeks)	PPROM	SGA (<10% ile)	TNMG
Hoff 2003	0/127 (0%)	22/127 (17.3%)	11/127 (8.7%)	94/127 (74%)	—	—	—	—	—	—	7/127 (5.5%)	—	—	5/127 (3.9%)
Braga 2016	10/28 (40%)	8/28 (28.6%)	5/28 (17.9%)	5/28 (17.9%)	17/30 (56.7%)	—	—	0/30 (0%)	1/30 (3.3%)	—	—	—	0/28 (0%)	2/28 (7.1%)
Assis 1983	0/37 (0%)	13/37 (35.1%)	1/37 (2.7%)	23/37 (62.2%)	15/39 (38.5%)	23/39 (59%)	1/39 (2.6%)	—	—	—	1/37 (2.7%)	—	—	1/37 (2.7%)
Wen 2009	0/163 (0%)	73/163 (44.8%)	9/163 (5.5%)	81/163 (49.7%)	—	—	—	—	—	—	13/163 (8.0%)	—	29/163 (17.8%)	—
Tellez-Zenteno 2004	0/18 (0%)	8/18 (44.4%)	1/18 (5.6%)	9/18 (50%)	7/18 (38.9%)	2/18 (11.1%)	9/18 (50%)	1/18 (5.6%)	—	—	—	—	—	1/17 (5.9%)
Tanacan 2019	0/23 (0%)	18/23 (78.3%)	0/23 (0%)	5/23 (21.7%)	11/27 (40.7%)	7/27 (25.9%)	9/27 (33.3%)	—	7/27 (25.9%)	—	3/23 (13.0%)	4/23 (17.4%)	—	6/23 (26.1%)
Slama 2008	0/5 (0%)	1/5 (20%)	2/5 (40%)	2/5 (40%)	—	—	—	0/8 (0%)	—	—	—	—	—	—
Shi 2018	1/8 (12.5%)	2/8 (25%)	0/8 (0%)	5/8 (62.5%)	2/8 (25%)	0/8 (0%)	6/8 (75%)	0/8 (0%)	1/8 (12.5%)	0/8 (0%)	1/8 (12.5%)	—	—	1/8 (12.5%)
Ramirez 2006	0/27 (0%)	7/27 (25.9%)	6/27 (22.2%)	14/27 (51.9%)	7/27 (25.9%)	4/27 (14.8%)	16/27 (59.3%)	2/27 (7.4%)	4/27 (14.8%)	0/27 (0%)	3/27 (11.1%)	0/27 (0%)	3/27 (11.1%)	3/27 (11.1%)
Qi 2012	0/7 (0%)	5/7 (71.4%)	0/7 (0%)	2/7 (28.6%)	4/7 (57.1%)	0/7 (0%)	3/7 (42.9%)	—	—	1/7 (14.3%)	1/7 (14.3%)	—	3/7 (42.9%)	0/7 (0%)
Picone 2003	1/13 (7.7%)	4/13 (30.8%)	2/13 (15.4%)	6/13 (46.2%)	5/13 (38.5%)	0/13 (0%)	8/13 (61.5%)	3/13 (23.1%)	2/13 (15.4%)	—	—	—	—	3/14 (21.4%)
Garcia 1989	0/6 (0%)	4/6 (66.7%)	2/6 (33.3%)	0/6 (0%)	3/6 (50%)	0/6 (0%)	3/6 (50%)	1/6 (16.7%)	1/6 (16.7%)	0/6 (0%)	0/6 (0%)	0/6 (0%)	—	0/6 (0%)
Tsurane 2019	0/6 (0%)	0/6 (0%)	4/6 (66.7%)	2/6 (33.3%)	3/6 (50%)	2/6 (33.3%)	1/6 (16.7%)	2/6 (33.3%)	2/6 (33.3%)	0/6 (0%)	0/6 (0%)	0/6 (0%)	—	0/6 (0%)
Thoulon 1978	0/22 (0%)	0/22 (0%)	6/22 (27.3%)	16/22 (72.7%)	7/23 (30.4%)	3/23 (13%)	13/23 (56.6%)	—	—	—	3/22 (13.6%)	—	4/22 (18.2%)	1/22 (4.6%)
Roth 2006	0/12 (0%)	1/12 (8.3%)	3/12 (25%)	8/12 (66.7%)	3/15 (20%)	3/15 (20%)	9/15 (60%)	0/15 (0%)	0/15 (0%)	—	—	—	—	2/12 (16.7%)
Podciechowski 2005	0/13 (0%)	6/13 (46.2%)	1/13 (7.7%)	6/13 (46.2%)	—	—	—	—	—	2/13 (15.4%)	2/13 (15.4%)	—	1/14 (7.1%)	1/14 (7.1%)
Drazanac 1978	0/5 (0%)	0/5 (0%)	1/5 (20%)	4/5 (80%)	3/5 (60%)	0/5 (0%)	3/5 (60%)	0/5 (0%)	0/5 (0%)	0/5 (0%)	0/5 (0%)	0/5 (0%)	0/5 (0%)	1/5 (20%)
Kostera-Pruszczyk 2005	0/38 (0%)	6/38 (15.8%)	0/38 (0%)	32/38 (84.2%)	4/38 (10.5%)	12/38 (31.6%)	22/38 (57.9%)	—	—	1/38 (2.6%)	3/38 (7.9%)	—	—	10/38 (26.3%)
Mitchell 1992	0/11 (0%)	1/11 (9.1%)	0/11 (0%)	10/11 (90.9%)	5/11 (45.5%)	0/11 (0%)	6/11 (54.5%)	4/11 (36.4%)	0/11 (0%)	1/11 (9.1%)	2/11 (18.2%)	0/11 (0%)	0/11 (0%)	1/11 (9.1%)
Perucca 2006	0/19 (0%)	3/19 (15.8%)	1/19 (5.3%)	15/19 (79%)	4/19 (21.1%)	0/19 (0%)	15/19 (79%)	3/19 (15.8%)	1/19 (5.3%)	0/19 (0%)	1/19 (5.3%)	0/19 (0%)	6/19 (31.6%)	3/19 (15.8%)
Miller 1968	0/7 (0%)	0/7 (0%)	1/7 (14.3%)	6/7 (85.7%)	2/7 (28.6%)	0/7 (0%)	5/7 (71.4%)	0/7 (0%)	1/7 (14.3%)	—	3/7 (42.9%)	—	—	—
Chabert 2004	0/10 (0%)	3/10 (30%)	5/10 (50%)	2/10 (20%)	6/10 (60%)	1/10 (10%)	3/10 (30%)	0/10 (0%)	0/10 (0%)	0/10 (0%)	0/10 (0%)	0/10 (0%)	—	—

(continued)

**Table 2. Continued.**

Author Year	Mode of delivery			MG course in pregnancy						Pregnancy complications				
	CS: MG as indication (%)	CS: Obstetric indication (%)	Operative vaginal delivery (%)	SVD (%)	Exacerbation (%)	Asymptomatic/improvement (%)	Unchanged (%)	Myasthenic crisis in pregnancy (%)	Myasthenic crisis postpartum (%)	PTB <34 weeks	PTB <37 weeks	PPROM	SGA (<10% ile)	TNMG
Castro 1995	0/9 (0%)	3/9 (33.3%)	0/9 (0%)	6/9 (66.7%)	2/9 (22.2%)	0/9 (0%)	7/9 (77.8%)	0/9 (0%)	0/9 (0%)	0/9 (0%)	0/9 (0%)	-	-	1/9 (11.1%)
Gordon 1968	0/11 (0%)	0/11 (0%)	0/11 (0%)	11/11 (100%)	5/11 (45.5%)	5/11 (45.5%)	1/11 (9.1%)	0/11 (0%)	1/11 (9.1%)	-	-	-	-	-
Gomez 2017	0/5 (0%)	3/5 (60%)	0/5 (0%)	2/5 (40%)	0/5 (0%)	0/5 (0%)	5/5 (100%)	0/5 (0%)	0/5 (0%)	0/5 (0%)	0/5 (0%)	-	3/5 (60.0%)	0/5 (0%)
Fraser 1953	0/7 (0%)	0/7 (0%)	1/7 (14.3%)	6/7 (85.7%)	5/14 (35.7%)	2/14 (14.3%)	7/14 (50%)	-	2/14 (14.3%)	0/7 (0%)	2/7 (28.6%)	1/7 (14.3%)	-	2/14 (14.3%)
Gveric-Ahmetasevic 2008	-	-	-	-	-	-	-	-	-	-	-	-	-	16/49 (32.7%)
Ducci 2016	5/30 (16.7%)	15/30 (50%)	2/30 (6.7%)	8/30 (26.7%)	15/30 (50%)	9/30 (30%)	6/30 (20%)	2/30 (6.7%)	1/30 (3.3%)	-	7/30 (23.3%)	7/30 (23.3%)	-	4/31 (12.9%)
Eden 1983	0/12 (0%)	2/12 (16.7%)	2/12 (16.7%)	8/12 (66.7%)	4/12 (33.3%)	3/12 (25%)	5/12 (41.7%)	1/12 (8.3%)	1/12 (8.3%)	3/12 (25%)	5/12 (41.7%)	1/12 (8.3%)	-	4/12 (33.3%)
Cheng 2007	4/13 (30.8%)	7/13 (53.8%)	0/13 (0%)	2/13 (15.4%)	5/13 (38.5%)	1/13 (7.7%)	7/13 (53.9%)	1/13 (7.7%)	3/13 (23.1%)	1/13 (7.7%)	1/13 (7.7%)	1/13 (7.7%)	-	1/14 (7.1%)
Almeida 2010	3/15 (20%)	7/15 (46.7%)	0/15 (0%)	5/15 (33.3%)	4/15 (26.7%)	3/15 (20%)	8/15 (53.3%)	0/15 (0%)	3/15 (20%)	0/15 (0%)	2/15 (13.3%)	2/15 (13.3%)	-	0/15 (0%)
Djelmis 2002	1/69 (1.5%)	11/69 (15.9%)	10/69 (14.5%)	47/69 (68%)	21/69 (30.4%)	17/69 (24.6%)	31/69 (44.9%)	1/69 (1.5%)	1/69 (1.5%)	0/68 (0%)	5/68 (7.4%)	0	3/68 (4.4%)	21/68 (30.9%)
Banner 2020	0/48 (0%)	16/48 (33.3%)	10/48 (20.8%)	22/48 (45.8%)	12/48 (25.0%)	3/48 (6.3%)	33/48 (68.8%)	4/48 (8.3%)	0/48 (0%)	5/48 (10.4%)	11/48 (22.9%)	4/48 (8.3%)	6/48 (12.5%)	1/48 (2.1%)
Total <sup>a</sup>	25/824 (3.0%)	249/824 (30.2%)	86/824 (10.4%)	464/824 (56.3%)	181/535 (33.8%)	100/505 (19.8%)	242/505 (47.9%)	23/357 (6.4%)	31/380 (8.2%)	14/327 (4.3%)	69/579 (11.9%)	27/404 (6.7%)	55/390 (14.1%)	89/686 (13.0%)

MG: myasthenia gravis; Tx: treatment; IVIG: intravenous immunoglobulin; PLEX: plasmapheresis; CS: caesarean section; SVD: spontaneous vaginal delivery; PTB: preterm birth; PPROM: preterm prelabour rupture of membranes; SGA: small for gestational age; TNMG: transient neonatal myasthenia gravis.  
<sup>a</sup>Data combined were available from all 34 data sets.



**Figure 2.** A) Summary of risks of bias with the use of the Quality Assessment Tool for Case Series (NIH.gov). Green represents low risk of bias. B) Risk of bias items from the Quality Assessment Tool for Case Series represented by percentages.

significant subset experiencing potentially life-threatening myasthenic crisis in pregnancy or postpartum. MG does not appear to significantly increase the rate of other pregnancy complications, with the exception of PPRM, which occurs at a rate roughly double that of the general population. Neonates should be observed for signs of TNMG, regardless of antenatal risk factors or maternal status.

### Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

### Funding

The authors received no financial support for the research, authorship and/or publication of this article.

### Ethical approval

The Research Ethics Board of Mount Sinai Hospital approved this study (#19-0183-C, July 26, 2019).

### Trial registration

The review was registered in the PROSPERO database (2019; registration number CRD42019143072) prior to study initiation.

### Guarantor

KEM is the guarantor of the present work.

### Contributorship

HB and KEM conceived, designed and oversaw the conduct of the study. HB wrote the initial draft of the manuscript, conducted abstract and full-text screening, conducted risk of bias assessment and synthesized the data. KMN conducted screening of abstracts and full texts and conducted risk of bias assessment. MR developed and carried out the search strategies. MS, VB and KEM provided clinical and methodological input. All authors critically revised the article and approved the final version for publication.

### ORCID iD

Harrison Banner  <https://orcid.org/0000-0002-2106-4711>

### Clinical Summary

The chance of spontaneous vaginal birth is greater than 50%. An assisted second stage of labour should be considered for women with recent exacerbation or those experiencing a myasthenic crisis.

The chance of disease exacerbation during pregnancy is 34%.

The chance of myasthenic crisis is 6.4% during pregnancy and 8.2% during the postpartum period. This complication is serious and can be life-threatening.

The risk of preterm birth and other pregnancy complications is similar to that of the general population.

The incidence of transient neonatal myasthenia gravis is 13%. This complication is unpredictable and should be monitored for during the early neonatal period.

**Figure 3.** Clinical Summary for the clinician counselling about myasthenia gravis in pregnancy.



## Supplemental material

Supplemental material for this article is available online.

## References

- Gilhus N and Verschuuren JJ. Myasthenia gravis: subgroup classification and therapeutic strategies. *Lancet Neurol* 2015; 14: 1023-1036.
- Deenan JC, Horlings CG, Verschuuren JJ, et al. The epidemiology of neuromuscular disorders: a comprehensive overview of the literature. *J Neuromuscul Dis* 2015; 2: 73-85.
- Carr AS, Cardwell CR, McCarron PO, et al. A systematic review of population based epidemiological studies in Myasthenia Gravis. *BMC Neurol* 2010; 10.
- Sanders DB, Wolfe GI, Benatar M, et al. International consensus guidance for management of myasthenia gravis. *Neurology* 2016; 87: 419-425.
- Sussman J, Farrugia ME, Maddison P, et al. The association of British Neurologists' myasthenia gravis guidelines. *Ann N Y Acad Sci* 2018; 1412: 166-169.
- Ahmed S, Kirmani JF, Janjua N, et al. An update on myasthenic crisis. *Curr Treat Options Neurology* 2005; 7: 129-141.
- Melzer N, Ruck T, Fuhr P, et al. Clinical features, pathogenesis and treatment of myasthenia gravis: a supplement to the guidelines of the German neurological society. *J Neurology* 2016; 263: 1473-1494.
- Tanacan A, Fadiloglu E, Ozten G, et al. Myasthenia gravis and pregnancy: retrospective evaluation of 27 pregnancies in a tertiary center and comparison with previous studies. *Ir J Med Sci* 2019; 188: 1261-1267.
- Djelmis J, Sostarko M, Mayer D, et al. Myasthenia gravis in pregnancy: report on 69 cases. *Eur J Obstetrics Gynecology Reproductive Biol* 2002; 104: 21-25.
- Téllez-Zenteno JF, Hernández-Ronquillo L, Salinas V, et al. Myasthenia gravis and pregnancy: clinical implications and neonatal outcome. *BMC Musculoskelet Disord* 2004; 5.
- Braga AC, Pinto C, Santos E, et al. Myasthenia gravis in pregnancy: experience of a Portuguese center. *Muscle Nerve* 2016; 54: 715-720.
- Qi Q-W, Wang D, Liu J-T, et al. [Management of pregnancy with myasthenia gravis: 7 cases report]. *Zhonghua fu chan ke za zhi* 2012; 47: 241-244.
- Kostera-Pruszczyk AE-SB. Macierzynstwo chorych na miastenie. *Gin Pol* 2005; 76: 122-125.
- Ducci RD, Lorenzoni PJ, Kay CSK, et al. Clinical follow-up of pregnancy in myasthenia gravis patients. *Neuromuscul Disord, NMD* 2017; 27: 352-357.
- Hoff JM, Daltveit AK and Gilhus NE. Myasthenia gravis consequences for pregnancy, delivery, and the newborn. *Neurology* 2003; 61: 1362-1366.
- Wen JC, Liu TC, Chen YH, et al. No increased risk of adverse pregnancy outcomes for women with myasthenia gravis: a nationwide population-based study. *Eur J Neurol* 2009; 16: 889-894.
- Perucca PE, Cuellar JE, Ricci AP, et al. Miastenia gravis: embarazo e impacto perinatal. *Rev Chil Obstet Ginecol* 2006; 71: 201-206.
- de Assis JL, Marchiori PE, Zambon AA, et al. Scaff M miastenia grave na gravidez e parto. *Neurobiol Recife* 1983; 46: 3-12.
- Eden RD and Gall SA. Myasthenia gravis and pregnancy: a reappraisal of thymectomy. *Obstet Gynecol* 1983; 62: 328-333.
- Kramer MS, Platt RW, Wen SW, et al., Fetal/Infant Health Study Group of the Canadian Perinatal Surveillance System. A new and improved population-based Canadian reference for birth weight for gestational age. *Pediatrics* 2001; 108: e35-e42.
- Tsurane K, Tanabe S, Miyasaka N, et al. Management of labor and delivery in myasthenia gravis: a new protocol. *J Obstet Gynaecol Res* 2019; 45: 974-980.
- Thoulong JM, Galopin G, Seffert P, et al. [Myasthenia and pregnancy (author's transl)]. *Myasthenie et grossesse* 1978; 7: 1395-1403.
- Tellez-Zenteno JF, Hernandez-Ronquillo L, Salinas V., et al. Myasthenia gravis and pregnancy: clinical implications and neonatal outcome. *BMC Musculoskelet Disord* 2004; 5: 42.
- Slama A, Tourni A, Tarmiz K, et al. Regional anaesthesia for labour and delivery in the myasthenic patient: report of five cases. *Annales Francaises D Anesthesie Et De Reanimation* 2008; 27: 180-181.
- Shi BB and Zeng LC. Management of myasthenia gravis during pregnancy: a report of eight cases. *Open Life Sci* 2018; 13: 28-33.
- Roth TC, Raths J, Carboni G, et al. Effect of pregnancy and birth on the course of myasthenia gravis before or after transsternal radical thymectomy. *Eur J Cardio-Thoracic Surgery, Off J Eur Assoc Cardio-Thoracic Surg* 2006; 29: 231-235.
- Ramirez C, de Seze J, Delrieu O, et al. [Myasthenia gravis and pregnancy: clinical course and management of delivery and the postpartum phase]. *Myasthenie Auto-immune Et grossesse: Evolution Clinique, Accouchement et Post-Partum* 2006; 162: 330-338.
- Podciechowski L, Brocka-Nitecka U, Dabrowska K, et al. Pregnancy complicated by Myasthenia gravis - twelve years experience. *Neuro Endocrinol Lett* 2005; 26: 603-608.
- Picone O, Audibert F, Gajdos P, et al. [Myasthenia gravis and pregnancy. Report on 13 cases]. *Myasthenie et grossesse: a propos de 13 grossesses* 2003; 32: 654-659.
- Mitchell PJ and Bebbington M. Myasthenia gravis in pregnancy. *Obstet Gynecol* 1992; 80: 178-181.
- Miller HJ, Sellmann AH and Mickal A. Myasthenia gravis of pregnancy at charity hospital. *J La Med Soc* 1968; 120: 231-234.
- Kostera-Pruszczyk A and Emeryk-Szajewska B. [Myasthenia gravis and pregnancy]. *Macierzynstwo Chorych Na Miastenie* 2005; 76: 122-126.
- Hoff JM, Daltveit AK and Gilhus NE. Myasthenia gravis: consequences for pregnancy, delivery, and the newborn. *Neurology* 2003; 61: 1362-1366.
- Gveric-Ahmetasevic S, Colic A, Elvedji-Gasparovic V, et al. Can neonatal myasthenia gravis be predicted? *J Perinat Med* 2008; 36: 503-506.
- Gordon SV and Peven PS. Myasthenia gravis and pregnancy. *Grace Hospital Bull* 1968; 46: 44-50.
- Garcia SA, Ogata AJ, Patriota RG, et al. Myasthenia gravis and pregnancy: review of the literature and report of 6 cases occurring in 5 patients. *Rev Paul Med* 1989; 107: 144-148.
- Gamez J, Salvado M, Casellas M, et al. Intravenous immunoglobulin as monotherapy for myasthenia gravis during pregnancy. *J Neurol Sci* 2017; 383: 118-122.
- Fraser D and Turner JWA. Myasthenia gravis and pregnancy. *Lancet* 1953; 2: 417-419.
- Ducci RD, Lorenzoni PJ, Kay CSK, et al. Clinical follow-up of the pregnancy in myasthenia gravis patients. *J Neuromuscul Dis* 2016; 3: S194.
- Drazancic A, Jusic A and Jurenic D. [Myasthenia gravis and pregnancy]. *Myasthenia Gravis I Trudnoca* 1978; 26: 41-46.
- Djelmis J, Sostarko M, Mayer D, et al. Myasthenia gravis in pregnancy: report on 69 cases. *Eur J Obstet Gynecol Reprod Biol* 2002; 104: 21-25.
- Cheng I, Lin C-H, Lin M-I, et al. Outcome of myasthenia gravis mothers and their infants. *Acta Paediatrica TAIWANICA = TAIWAN Er Ke Yi Xue Hui Za Zhi* 2007; 48: 141-145.
- Chabert L and Benhamou D. Myasthenia gravis, pregnancy and delivery: a series of ten cases. *Ann Fr Anesth Reanim* 2004; 23: 459-464.

44. Castro R, Cunha S, Duarte G, et al. Miastenia gravis e gravidez: análise de 9 casos. *Rev Bras Ginecol Obstet* 1995; 17: 34-42.
45. Assis JL, Marchiori PE, Zambon AA, et al. Miastenia grave na gravidez e parto. *Neurobiologia* 1983; 46: 3-12.
46. Almeida C, Coutinho E, Moreira D, et al. Myasthenia gravis and pregnancy: anaesthetic management—a series of cases. *Eur J Anaesthesiol* 2010; 27: 985-990.
47. Chabert L and Benhamou D. [Myasthenia gravis, pregnancy and delivery: a series of ten cases]. *Myasthenie, Grossesse Et Accouchement: a Propos De Dix Cas* 2004; 23: 459-464.
48. Uncu G, Küçükdoğan I, Ozan H, et al. Pregnancy and myasthenia gravis. A case report. *Clin Exp Obstet Gynecol* 1995; 22: 145-147.
49. Smith GN, Rafuse C, Anand N, et al. Prevalence, management, and outcomes of preterm prelabour rupture of membranes of women in Canada. *J Obstet Gynaecol Can* 2005; 27: 547-553.
50. Blencowe H, Cousens S, Oestergaard MZ, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet* 2012; 379: 2162-2172.